

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 March 2004 (18.03.2004)

PCT

(10) International Publication Number
WO 2004/022561 A1

(51) International Patent Classification?: **C07D 487/04**,
A61K 31/519 // (C07D 487/04, 239:00, 231:00)

(21) International Application Number:
PCT/US2003/027555

(22) International Filing Date:
3 September 2003 (03.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/408,027 4 September 2002 (04.09.2002) US
60/421,959 29 October 2002 (29.10.2002) US

(71) Applicants: SCHERING CORPORATION [US/US];
2000 Galloping Hill Road, Kenilworth, NJ 07033-0530
(US). PHARMACOPEIA, INC. [US/US]; 3000 Eastpark
Boulevard, Cranbury, NJ 08512 (US).

(72) Inventors: GUZI, Timothy, J.; 48 Red Road, Chatham,
NJ 07928 (US). PARUCH, Kamil; 20 Third Avenue,
Garwood, NJ 07027 (US). DWYER, Michael, P.; 235
Katherine Street, Scotch Plains, NJ 07076 (US). DOLL,
Ronald, J.; 8 Concord Lane, Convent Station, NJ 07960
(US). GIRIJAVALLABHAN, Viyyoor, Moopill; 10
Maplewood Drive, Parsippany, NJ 07054 (US). MAL-
LAMS, Alan; 147 Kings Highway, RR3, Hackettstown,
NJ 07840 (US). ALVAREZ, Carmen, S.; 121 Walnut
Street, Livingston, NJ 07039 (US). KEERTIKAR, Kar-
tik, M.; 24 Columbia Avenue, East Windsor, NJ 08520
(US). RIVERA, Jocelyn; 72 Regal Drive, Monmouth
Junction, NJ 08852 (US). CHAN, Tin-Yau; 26 Barlow
Road, Edison, NJ 08817 (US). MADISON, Vincent;
12 Ronarm Drive, Mountain Lakes, NJ 07046 (US).
FISCHMANN, Thierry, O.; 2700 Sky Top Drive, Scotch
Plains, NJ 07076 (US). DILLARD, Lawrence, W.; 278
Spring Hill Road, Skillman, NJ 08558 (US). TRAN, Vinh,
D.; 17374 Winemast Street, Fountain Valley, CA 92708
(US). HE, Zhen Min; 5 Pennyroyal Court, Princeton,

NJ 08540 (US). JAMES, Ray, Anthony; 1036 Radcliffe
Street, Bristol, PA 19007 (US). PARK, Haengsoon; 8910
Tamarron Drive, Plainsboro, NJ 08536 (US). PARAD-
KAR, Vidyadhar, M.; 3 Pine Ridge Drive, Somerville, NJ
08876 (US). HOBBS, Douglas, Walsh; 1330 University
Drive, Yardley, PA 19067 (US).

(74) Agent: KALYANARAMAN, Palaiyur, S.; Schering Cor-
poration, Patent Department K-6-1 1990, 2000 Galloping
Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,
MA, MD, MG, MK, MN, MX, MZ, NI, NO, NZ, PG, PH,
PL, PT, RO, RU, SC, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UZ, VC, VN, YU, ZA, ZM.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: PYRAZOLOPYRIMIDINES AS CYCLIN-DEPENDENT KINASE INHIBITORS

(57) Abstract: In its many embodiments, the present invention provides a novel class of pyrazolo[1,5-a]pyrimidine compounds as inhibitors of cyclin dependent kinases, methods of preparing such compounds, pharmaceutical compositions containing one or more such compounds, methods of preparing pharmaceutical formulations comprising one or more such compounds, and methods of treatment, prevention, inhibition, or amelioration of one or more diseases associated with the CDKs using such compounds or pharmaceutical compositions.

WO 2004/022561 A1

PYRAZOLOPYRIMIDINES AS CYCLIN-DEPENDENT KINASE INHIBITORS

Field of the Invention


The present invention relates to pyrazolo[1,5-a]pyrimidine compounds useful as protein kinase inhibitors (such as for example, the inhibitors of the cyclin-dependent kinases, mitogen-activated protein kinase (MAPK/ERK), glycogen synthase kinase 3 (GSK3beta) and the like), pharmaceutical compositions containing the compounds, and methods of treatment using the compounds and compositions to treat diseases such as, for example, cancer, inflammation, arthritis, viral diseases, neurodegenerative diseases such as Alzheimer's disease, cardiovascular diseases, and fungal diseases. This application claims benefit of priority from U.S. provisional patent applications, Serial No. 60/408,027 filed September 4, 2002, and Serial No. 60/421,959 filed October 29, 2002.

Background of the Invention

Protein kinase inhibitors include kinases such as, for example, the inhibitors of the cyclin-dependent kinases (CDKs), mitogen activated protein kinase (MAPK/ERK), glycogen synthase kinase 3 (GSK3beta), and the like. Protein kinase inhibitors are described, for example, by M. Hale *et al* in WO02/22610 A1 and by Y. Mettley *et al* in *J. Med. Chem.*, (2003) 46 222-236. The cyclin-dependent kinases are serine/threonine protein kinases, which are the driving force behind the cell cycle and cell proliferation. Individual CDK's, such as, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6 and CDK7, CDK8 and the like, perform distinct roles in cell cycle progression and can be classified as either G1, S, or G2M phase enzymes. Uncontrolled proliferation is a hallmark of cancer cells, and misregulation of CDK function occurs with high frequency in many important solid tumors. CDK2 and CDK4 are of particular interest because their activities are frequently misregulated in a wide variety of human cancers. CDK2 activity is required for progression through G1 to the S phase of the cell cycle, and CDK2 is one of the key components of the G1 checkpoint. Checkpoints serve to maintain the proper sequence of cell cycle events and allow the cell to

respond to insults or to proliferative signals, while the loss of proper checkpoint control in cancer cells contributes to tumorigenesis. The CDK2 pathway influences tumorigenesis at the level of tumor suppressor function (e.g. p52, RB, and p27) and oncogene activation (cyclin E). Many reports have demonstrated that both the coactivator, cyclin E, and the inhibitor, p27, of CDK2 are either over – or underexpressed, respectively, in breast, colon, nonsmall cell lung, gastric, prostate, bladder, non-Hodgkin's lymphoma, ovarian, and other cancers. Their altered expression has been shown to correlate with increased CDK2 activity levels and poor overall survival. This observation makes CDK2 and its regulatory pathways compelling targets for the development years, a number of adenosine 5'-triphosphate (ATP) competitive small organic molecules as well as peptides have been reported in the literature as CDK inhibitors for the potential treatment of cancers. U.S. 6,413,974, col. 1, line 23- col. 15, line 10 offers a good description of the various CDKs and their relationship to various types of cancer.


CDK inhibitors are known. For example, flavopiridol (Formula I) is a nonselective CDK inhibitor that is currently undergoing human clinical trials, A. M. Sanderowicz *et al*, *J. Clin. Oncol.* (1998) 16, 2986-2999.



Formula I

Other known inhibitors of the CDKs include, for example, olomoucine (J. Vesely *et al*, *Eur. J. Biochem.*, (1994) 224, 771-786) and roscovitine (I. Meijer *et al*, *Eur. J. Biochem.*, (1997) 243, 527-536). U.S. 6,107,305 describes certain pyrazolo[3,4-b] pyridine compounds as CDK inhibitors. An illustrative compound from the '305 patent has the Formula II:

3



Formula II

K. S. Kim *et al*, *J. Med. Chem.* 45 (2002) 3905-3927 and WO 02/10162 disclose certain aminothiazole compounds as CDK inhibitors.

5 Pyrazolopyrimidines are known. For Example, WO92/18504, WO02/50079, WO95/35298, WO02/40485, EP94304104.6, EP0628559 (equivalent to US Patents 5,602,136, 5,602,137 and 5,571,813), U.S. 6,383,790, *Chem. Pharm. Bull.*, (1999) 47 928, *J. Med. Chem.*, (1977) 20, 296, *J. Med. Chem.*, (1976) 19 517 and *Chem. Pharm. Bull.*, (1962) 10 620 disclose various
10 pyrazolopyrimidines.


There is a need for new compounds, formulations, treatments and therapies to treat diseases and disorders associated with CDKs. It is, therefore, an object of this invention to provide compounds useful in the treatment or prevention or amelioration of such diseases and disorders.

15

Summary of the Invention

In its many embodiments, the present invention provides a novel class of pyrazolo[1,5-a]pyrimidine compounds as inhibitors of cyclin dependent kinases, methods of preparing such compounds, pharmaceutical compositions comprising
20 one or more such compounds, methods of preparing pharmaceutical formulations comprising one or more such compounds, and methods of treatment, prevention, inhibition or amelioration of one or more diseases associated with the CDKs using such compounds or pharmaceutical compositions.


In one aspect, the present application discloses a compound, or pharmaceutically acceptable salts or solvates of said compound, said compound having the general structure shown in Formula III:



Formula III

wherein:

R is H, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, cycloalkyl, cycloalkylalkyl, alkenylalkyl, alkynylalkyl, heterocyclyl, heterocyclalkyl, heteroarylalkyl (including N-oxide of said heteroaryl), $-(CHR^5)_n$ -aryl, $-(CHR^5)_n$ -





wherein each of said alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocyclyl, and heteroaryl can be unsubstituted or optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, heterocyclalkyl, CF_3 , OCF_3 , CN, $-OR^5$, $-NR^5R^{10}$, $-C(R^4R^5)_p-R^9$, $-N(R^5)Boc$, $-(CR^4R^5)_pOR^5$, $-C(O_2)R^5$, $-C(O)R^5$, $-C(O)NR^5R^{10}$, $-SO_3H$, $-SR^{10}$, $-S(O_2)R^7$, $-S(O_2)NR^5R^{10}$, $-N(R^5)S(O_2)R^7$, $-N(R^5)C(O)R^7$ and $-N(R^5)C(O)NR^5R^{10}$;

20 R^2 is selected from the group consisting of R^9 , alkyl, alkenyl, alkynyl, CF_3 , heterocyclyl, heterocyclalkyl, halogen, haloalkyl, aryl, arylalkyl, heteroarylalkyl, alkynylalkyl, cycloalkyl, heteroaryl, alkyl substituted with 1-6 R^9 groups which can be the same or different and are independently selected from the list of R^9 shown below, aryl substituted with 1-3 aryl or heteroaryl groups which can be the same or different and are independently selected from phenyl, pyridyl,

25


thiophenyl, furanyl and thiazolo groups, aryl fused with an aryl or heteroaryl group, heteroaryl substituted with 1-3 aryl or heteroaryl groups which can be the same or different and are independently selected from phenyl, pyridyl, thiophenyl, furanyl and thiazolo groups, heteroaryl fused with an aryl or



wherein one or more of the aryl and/or one or more of the heteroaryl in the above-noted definitions for R^2 can be unsubstituted or optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, $-CN$, $-OR^5$, $-SR^5$, $-S(O_2)R^6$, $-S(O_2)NR^5R^6$, $-NR^5R^6$, $-C(O)NR^5R^6$, CF_3 , alkyl, aryl and OCF_3 ;

R^3 is selected from the group consisting of H, halogen, $-NR^5R^6$, $-OR^6$, $-SR^6$, $-C(O)N(R^5R^6)$, alkyl, alkynyl, cycloalkyl, aryl, arylalkyl, heterocyclyl,

15 heterocyclylalkyl, heteroaryl and heteroarylalkyl,



wherein each of said alkyl, cycloalkyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl for R^3 and the heterocyclyl

20 moieties whose structures are shown immediately above for R^3 can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, CF_3 , CN , $-OCF_3$,

25 $-(CR^4R^5)_pOR^5$, $-OR^5$, $-NR^5R^6$, $-(CR^4R^5)_pNR^5R^6$, $-C(O_2)R^5$, $-C(O)R^5$, $-C(O)NR^5R^6$, $-SR^6$, $-S(O_2)R^6$, $-S(O_2)NR^5R^6$, $-N(R^5)S(O_2)R^7$, $-N(R^5)C(O)R^7$ and

-N(R⁵)C(O)NR⁵R⁶, with the proviso that no carbon adjacent to a nitrogen atom on a heterocyclyl ring carries a -OR⁵ moiety;

R⁴ is H, halo or alkyl;

R⁵ is H, alkyl, aryl or cycloalkyl;

5 R⁶ is selected from the group consisting of H, alkyl, alkenyl, aryl, arylalkyl, arylalkenyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl, wherein each of said alkyl, aryl, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl can be unsubstituted or optionally substituted with one or more moieties which can be the same or
10 different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, heterocyclylalkyl, CF₃, OCF₃, CN, -OR⁵, -NR⁵R¹⁰, -C(R⁴R⁵)_p-R⁹, -N(R⁵)Boc, -(CR⁴R⁵)_pOR⁵, -C(O₂)R⁵, -C(O)R⁵, -C(O)NR⁵R¹⁰, -SO₃H, -SR¹⁰, -S(O₂)R⁷, -S(O₂)NR⁵R¹⁰, -N(R⁵)S(O₂)R⁷, -N(R⁵)C(O)R⁷ and -N(R⁵)C(O)NR⁵R¹⁰;

15 R¹⁰ is selected from the group consisting of H, alkyl, aryl, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl, wherein each of said alkyl, aryl, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl can be unsubstituted or optionally substituted with one or more moieties which can be the same or
20 different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, heterocyclylalkyl, CF₃, OCF₃, CN, -OR⁵, -NR⁴R⁵, -C(R⁴R⁵)_p-R⁹, -N(R⁵)Boc, -(CR⁴R⁵)_pOR⁵, -C(O₂)R⁵, -C(O)NR⁴R⁵, -C(O)R⁵, -SO₃H, -SR⁵, -S(O₂)R⁷, -S(O₂)NR⁴R⁵, -N(R⁵)S(O₂)R⁷, -N(R⁵)C(O)R⁷ and -N(R⁵)C(O)NR⁴R⁵;

25 or optionally (i) R⁵ and R¹⁰ in the moiety -NR⁵R¹⁰, or (ii) R⁵ and R⁶ in the moiety -NR⁵R⁶, may be joined together to form a cycloalkyl or heterocyclyl moiety, with each of said cycloalkyl or heterocyclyl moiety being unsubstituted or optionally independently being substituted with one or more R⁹ groups;

30 R⁷ is selected from the group consisting of alkyl, cycloalkyl, aryl, arylalkenyl, heteroaryl, arylalkyl, heteroarylalkyl, heteroarylalkenyl, and heterocyclyl, wherein each of said alkyl, cycloalkyl, heteroarylalkyl, aryl, heteroaryl and arylalkyl can be unsubstituted or optionally independently

substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, CF_3 , OCF_3 , CN , $-\text{OR}^5$, $-\text{NR}^5\text{R}^{10}$, $-\text{CH}_2\text{OR}^5$, $-\text{C}(\text{O}_2)\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{C}(\text{O})\text{R}^5$, $-\text{SR}^{10}$, $-\text{S}(\text{O}_2)\text{R}^{10}$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$,
 5 $-\text{N}(\text{R}^5)\text{S}(\text{O}_2)\text{R}^{10}$, $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^{10}$ and $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{NR}^5\text{R}^{10}$;

R^8 is selected from the group consisting of R^6 , $-\text{OR}^6$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(=\text{N}-\text{CN})-\text{NH}_2$, $-\text{C}(=\text{NH})-\text{NHR}^5$, heterocyclyl, and $-\text{S}(\text{O}_2)\text{R}^7$;

R^9 is selected from the group consisting of halogen, $-\text{CN}$, $-\text{NR}^5\text{R}^{10}$,
 10 $-\text{C}(\text{O}_2)\text{R}^6$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{OR}^6$, $-\text{SR}^6$, $-\text{S}(\text{O}_2)\text{R}^7$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$, $-\text{N}(\text{R}^5)\text{S}(\text{O}_2)\text{R}^7$, $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^7$ and $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{NR}^5\text{R}^{10}$;

m is 0 to 4;

n is 1 to 4; and

p is 1 to 4,

15 with the proviso that when R^2 is phenyl, R^3 is not alkyl, alkynyl or halogen, and

that when R^2 is aryl, R is not $\begin{array}{c} \text{~} \\ \text{~} \\ \text{~} \end{array} \text{---} (\text{CHR}^5)_n \text{---} \text{NR}^5\text{R}^8$, and with the further proviso that when R is arylalkyl, then any heteroaryl substituent on the aryl of said arylalkyl contains at least three heteroatoms.

The compounds of Formula III can be useful as protein kinase inhibitors
 20 and can be useful in the treatment and prevention of proliferative diseases, for example, cancer, inflammation and arthritis. They may also be useful in the treatment of neurodegenerative diseases such Alzheimer's disease, cardiovascular diseases, viral diseases and fungal diseases.

25


Detailed Description

In one embodiment, the present invention discloses pyrazolo[1,5-a]pyrimidine compounds which are represented by structural Formula III, or a pharmaceutically acceptable salt or solvate thereof, wherein the various moieties are as described above.

30


In another embodiment, R is $-(\text{CHR}^5)_n\text{-aryl}$, $-(\text{CHR}^5)_n\text{-heteroaryl}$,

-(CHR⁵)_n-heteroaryl (with said heteroaryl being substituted with an additional, same or different, heteroaryl), -(CHR⁵)_n-heterocyclyl (with said heterocyclyl being substituted with an additional, same or different, heterocyclyl), or



5 In another embodiment, R² is halogen, CF₃, CN, lower alkyl, alkyl substituted with -OR⁶, alkynyl, aryl, heteroaryl or heterocyclyl.

In another embodiment, R³ is H, lower alkyl, aryl, heteroaryl, cycloalkyl, -NR⁵R⁶,



10

wherein said alkyl, aryl, heteroaryl, cycloalkyl and the heterocyclyl structures shown immediately above for R³ are optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, CF₃, OCF₃, lower alkyl, CN, -C(O)R⁵, -S(O₂)R⁵, -C(=NH)-NH₂, -C(=CN)-NH₂, hydroxyalkyl, alkoxy carbonyl, -SR⁵, and OR⁵, with the proviso that no carbon adjacent to a nitrogen atom on a heterocyclyl ring carries a -OR⁵ moiety.

15

In another embodiment, R⁴ is H or lower alkyl.

20

In another embodiment, R⁵ is H, lower alkyl or cycloalkyl.


In another embodiment, n is 1 to 2.

In an additional embodiment, R is -(CHR⁵)_n-aryl, -(CHR⁵)_n-heteroaryl.

In an additional embodiment, R^2 is halogen, CF_3 , CN, lower alkyl, alkynyl, or alkyl substituted with $-OR^6$.

In an additional embodiment, R^2 is lower alkyl, alkynyl or Br.

In an additional embodiment, R^3 is H, lower alkyl, aryl,



wherein said alkyl, aryl and the heterocyclyl moieties shown immediately above for R^3 are optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, CF_3 , lower alkyl, hydroxyalkyl, alkoxy, $-S(O_2)R^5$, and CN.

10 In an additional embodiment, R^4 is H.


In an additional embodiment, R^5 is H, ethyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

In an additional embodiment, R^8 is alkyl or hydroxyalkyl.


In an additional embodiment, n is 1.

15 In an additional embodiment, p is 1 or 2.


Another embodiment discloses the inventive compounds shown in **Table 1**, which exhibited CDK2 inhibitory activity of about $0.0001\mu M$ to $>$ about $5\mu M$. The assay methods are described later (from page 333 onwards).


Table 1


11




13












18




5

19






21




5

22




23



5


Another embodiment of the invention discloses the following compounds, which exhibited CDK2 inhibitory activity of about 0.0001 μ M to about 0.5 μ M:

24




5

26



5


27




5


Another embodiment of the invention discloses the following compounds, which exhibited CDK2 inhibitory activity of about 0.0001 μ M to about 0.1 μ M:


28





30





As used above, and throughout this disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

"Patient" includes both human and animals.

"Mammal" means humans and other mammalian animals.

"Alkyl" means an aliphatic hydrocarbon group which may be straight or branched and comprising about 1 to about 20 carbon atoms in the chain.

- 5 Preferred alkyl groups contain about 1 to about 12 carbon atoms in the chain. More preferred alkyl groups contain about 1 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. "Lower alkyl" means a group having about 1 to about 6 carbon atoms in the chain which may be straight or
- 10 branched. The term "substituted alkyl" means that the alkyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, carboxy and -C(O)O-alkyl. Non-limiting examples of
- 15 suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl and t-butyl.

- "Alkynyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably
- 20 about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkynyl chain. "Lower alkynyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butylnyl and 3-methylbutynyl. The
- 25 term "substituted alkynyl" means that the alkynyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of alkyl, aryl and cycloalkyl.

- "Aryl" means an aromatic monocyclic or multicyclic ring system
- 30 comprising about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as

defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl.

"Heteroaryl" means an aromatic monocyclic or multicyclic ring system comprising about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the ring atoms is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. Preferred heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. A nitrogen atom of a heteroaryl can be optionally oxidized to the corresponding N-oxide. Non-limiting examples of suitable heteroaryls include pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridone (including N-substituted pyridones), isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, 1,2,4-thiadiazolyl, pyrazinyl, pyridazinyl, quinoxaliny, phthalazinyl, oxindolyl, imidazo[1,2-a]pyridinyl, imidazo[2,1-b]thiazolyl, benzofurazanyl, indolyl, azaindolyl, benzimidazolyl, benzothienyl, quinoliny, imidazolyl, thienopyridyl, quinazolinyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, isoquinoliny, benzoazaindolyl, 1,2,4-triazinyl, benzothiazolyl and the like. The term "heteroaryl" also refers to partially saturated heteroaryl moieties such as, for example, tetrahydroisoquinolyl, tetrahydroquinolyl and the like.

"Aralkyl" or "arylalkyl" means an aryl-alkyl- group in which the aryl and alkyl are as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, 2-phenethyl and naphthalenylmethyl. The bond to the parent moiety is through the alkyl.

"Alkylaryl" means an alkyl-aryl- group in which the alkyl and aryl are as previously described. Preferred alkylaryls comprise a lower alkyl group. Non-limiting example of a suitable alkylaryl group is tolyl. The bond to the parent moiety is through the aryl.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms.

The cycloalkyl can be optionally substituted with one or more "ring system

5 substituents" which may be the same or different, and are as defined above.

Non-limiting examples of suitable monocyclic cycloalkyls include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable multicyclic cycloalkyls include 1-decalinyl, norbornyl, adamantyl and the like, as well as partially saturated species such as, for example, indanyl,


10 tetrahydronaphthyl and the like.

"Halogen" means fluorine, chlorine, bromine, or iodine. Preferred are fluorine, chlorine and bromine.

"Ring system substituent" means a substituent attached to an aromatic or non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, alkylaryl, heteroaralkyl, heteroarylalkenyl, heteroarylalkynyl, alkylheteroaryl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halo, nitro, cyano, carboxy, alkoxycarbonyl,


20 aryloxycarbonyl, aralkoxycarbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, heterocyclyl, $-C(=N-CN)-NH_2$, $-C(=NH)-NH_2$, $-C(=NH)-NH(alkyl)$, Y_1Y_2N- , $Y_1Y_2N-alkyl-$, $Y_1Y_2NC(O)-$, $Y_1Y_2NSO_2-$ and $-SO_2NY_1Y_2$, wherein Y_1 and Y_2 can be the same or different and are independently selected from the group

25 consisting of hydrogen, alkyl, aryl, cycloalkyl, and aralkyl. "Ring system substituent" may also mean a single moiety which simultaneously replaces two available hydrogens on two adjacent carbon atoms (one H on each carbon) on a ring system. Examples of such moiety are methylene dioxy, ethylenedioxy, $-C(CH_3)_2-$ and the like which form moieties such as, for example:



"Heterocyclyl" means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. Any -NH in a heterocyclyl ring may exist protected such as, for example, as an -N(Boc), -N(CBz), -N(Tos) group and the like; such protections are also considered part of this invention. The heterocyclyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, lactam, lactone, and the like.

It should be noted that in hetero-atom containing ring systems of this invention, there are no hydroxyl groups on carbon atoms adjacent to a N, O or S, as well as there are no N or S groups on carbon adjacent to another heteroatom. Thus, for example, in the ring:



there is no -OH attached directly to carbons marked 2 and 5.

It should also be noted that tautomeric forms such as, for example, the moieties:



are considered equivalent in certain embodiments of this invention.

"Alkynylalkyl" means an alkynyl-alkyl- group in which the alkynyl and alkyl are as previously described. Preferred alkynylalkyls contain a lower alkynyl and a lower alkyl group. The bond to the parent moiety is through the alkyl. Non-limiting examples of suitable alkynylalkyl groups include propargylmethyl.

5 "Heteroaralkyl" means a heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain a lower alkyl group. Non-limiting examples of suitable aralkyl groups include pyridylmethyl, and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkyl.

10 "Hydroxyalkyl" means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

15 "Acyl" means an H-C(O)-, alkyl-C(O)- or cycloalkyl-C(O)-, group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl and propanoyl.

"Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1- naphthoyl.

20 "Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy, n-propoxy, isopropoxy and n-butoxy. The bond to the parent moiety is through the ether oxygen.

25 "Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond to the parent moiety is through the ether oxygen.

30 "Aralkyloxy" means an aralkyl-O- group in which the aralkyl group is as previously described. Non-limiting examples of suitable aralkyloxy groups include benzyloxy and 1- or 2-naphthalenemethoxy. The bond to the parent moiety is through the ether oxygen.

"Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkylthio groups include methylthio and ethylthio. The bond to the parent moiety is through the sulfur.

5 "Arylthio" means an aryl-S- group in which the aryl group is as previously described. Non-limiting examples of suitable arylthio groups include phenylthio and naphthylthio. The bond to the parent moiety is through the sulfur.

"Aralkylthio" means an aralkyl-S- group in which the aralkyl group is as previously described. Non-limiting example of a suitable aralkylthio group is benzylthio. The bond to the parent moiety is through the sulfur.

10 "Alkoxycarbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxycarbonyl groups include methoxycarbonyl and ethoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aryloxycarbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxycarbonyl groups include phenoxycarbonyl and
15 naphthoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aralkoxycarbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxycarbonyl group is benzyloxycarbonyl. The bond to the parent moiety is through the carbonyl.

20 "Alkylsulfonyl" means an alkyl-S(O₂)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

"Arylsulfonyl" means an aryl-S(O₂)- group. The bond to the parent moiety is through the sulfonyl.

25 The term "substituted" means that one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. By "stable compound" or "stable
30 structure" is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties.

The term "isolated" or "in isolated form" for a compound refers to the physical state of said compound after being isolated from a synthetic process or natural source or combination thereof. The term "purified" or "in purified form" for a compound refers to the physical state of said compound after being obtained from a purification process or processes described herein or well known to the skilled artisan, in sufficient purity to be characterizable by standard analytical techniques described herein or well known to the skilled artisan.

It should also be noted that any heteroatom with unsatisfied valences in the text, schemes, examples and Tables herein is assumed to have the hydrogen atom(s) to satisfy the valences.

When a functional group in a compound is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for example, T. W. Greene *et al*, *Protective Groups in organic Synthesis* (1991), Wiley, New York.

When any variable (e.g., aryl, heterocycle, R^2 , etc.) occurs more than one time in any constituent or in Formula III, its definition on each occurrence is independent of its definition at every other occurrence.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

Prodrugs and solvates of the compounds of the invention are also contemplated herein. The term "prodrug", as employed herein, denotes a compound that is a drug precursor which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of Formula III or a salt and/or solvate thereof. A discussion of prodrugs is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems* (1987) 14 of the A.C.S. Symposium Series, and in *Bioreversible*

Carriers in Drug Design, (1987) Edward B. Roche, ed., American Pharmaceutical Association and Pergamon Press, both of which are incorporated herein by reference thereto.

"Solvate" means a physical association of a compound of this invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid.

"Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanولات, methanولات, and the like.

"Hydrate" is a solvate wherein the solvent molecule is H₂O.

"Effective amount" or "therapeutically effective amount" is meant to describe an amount of compound or a composition of the present invention effective in inhibiting the CDK(s) and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect.

The compounds of Formula III can form salts which are also within the scope of this invention. Reference to a compound of Formula III herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of Formula III contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein.

Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compounds of the Formula III may be formed, for example, by reacting a compound of Formula III with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates,

camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides, lactates, maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates,) and the like.

- 5 Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by S. Berge *et al*, *Journal of Pharmaceutical Sciences* (1977) 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33 201-217; Anderson *et al*, *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

- Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as dicyclohexylamines, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, and dibutyl sulfates), long chain halides (e.g. decyl, lauryl, and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

- All such acid salts and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.

Compounds of Formula III, and salts, solvates and prodrugs thereof, may exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

- 30 All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates and prodrugs of the compounds as well as the salts and solvates of the prodrugs),

such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this invention, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the *IUPAC* 1974 Recommendations.

The use of the terms "salt", "solvate" "prodrug" and the like, is intended to equally apply to the salt, solvate and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the inventive compounds.

The compounds according to the invention have pharmacological properties; in particular, the compounds of Formula III can be inhibitors of protein kinases such as, for example, the inhibitors of the cyclin-dependent kinases, mitogen-activated protein kinase (MAPK/ERK), glycogen synthase kinase 3(GSK3beta) and the like. The cyclin dependent kinases (CDKs) include, for example, CDC2 (CDK1), CDK2, CDK4, CDK5, CDK6, CDK7 and CDK8. The novel compounds of Formula III are expected to be useful in the therapy of proliferative diseases such as cancer, autoimmune diseases, viral diseases, fungal diseases, neurological/neurodegenerative disorders, arthritis, inflammation, anti-proliferative (e.g., ocular retinopathy), neuronal, alopecia and cardiovascular disease. Many of these diseases and disorders are listed in U.S. 6,413,974 cited earlier, the disclosure of which is incorporated herein.

More specifically, the compounds of Formula III can be useful in the treatment of a variety of cancers, including (but not limited to) the following: carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma; hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T- cell

lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma and Burkett's lymphoma;

hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic

5 leukemia;

tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma;

tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma and schwannomas; and

10 other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma.

Due to the key role of CDKs in the regulation of cellular proliferation in general, inhibitors could act as reversible cytostatic agents which may be useful
15 in the treatment of any disease process which features abnormal cellular proliferation, e.g., benign prostate hyperplasia, familial adenomatosis polyposis, neuro-fibromatosis, atherosclerosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis following angioplasty or vascular surgery, hypertrophic scar formation, inflammatory bowel disease, transplantation
20 rejection, endotoxic shock, and fungal infections.

Compounds of Formula III may also be useful in the treatment of Alzheimer's disease, as suggested by the recent finding that CDK5 is involved in the phosphorylation of tau protein (*J. Biochem*, (1995) 117, 741-749).

Compounds of Formula III may induce or inhibit apoptosis. The apoptotic
25 response is aberrant in a variety of human diseases. Compounds of Formula III, as modulators of apoptosis, will be useful in the treatment of cancer (including but not limited to those types mentioned hereinabove), viral infections (including but not limited to herpesvirus, poxvirus, Epstein- Barr virus, Sindbis virus and adenovirus), prevention of AIDS development in HIV-infected individuals,
30 autoimmune diseases (including but not limited to systemic lupus, erythematosus, autoimmune mediated glomerulonephritis, rheumatoid arthritis, psoriasis, inflammatory bowel disease, and autoimmune diabetes mellitus),

neurodegenerative disorders (including but not limited to Alzheimer's disease, AIDS-related dementia, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, spinal muscular atrophy and cerebellar degeneration), myelodysplastic syndromes, aplastic anemia, ischemic injury associated with myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol related liver diseases, hematological diseases (including but not limited to chronic anemia and aplastic anemia), degenerative diseases of the musculoskeletal system (including but not limited to osteoporosis and arthritis) aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple sclerosis, kidney diseases and cancer pain.

Compounds of Formula III, as inhibitors of the CDKs, can modulate the level of cellular RNA and DNA synthesis. These agents would therefore be useful in the treatment of viral infections (including but not limited to HIV, human papilloma virus, herpesvirus, poxvirus, Epstein-Barr virus, Sindbis virus and adenovirus).

Compounds of Formula III may also be useful in the chemoprevention of cancer. Chemoprevention is defined as inhibiting the development of invasive cancer by either blocking the initiating mutagenic event or by blocking the progression of pre-malignant cells that have already suffered an insult or inhibiting tumor relapse.

Compounds of Formula III may also be useful in inhibiting tumor angiogenesis and metastasis.

Compounds of Formula III may also act as inhibitors of other protein kinases, e.g., protein kinase C, her2, raf 1, MEK1, MAP kinase, EGF receptor, PDGF receptor, IGF receptor, PI3 kinase, wee1 kinase, Src, Abl and thus be effective in the treatment of diseases associated with other protein kinases.

Another aspect of this invention is a method of treating a mammal (e.g., human) having a disease or condition associated with the CDKs by administering a therapeutically effective amount of at least one compound of Formula III, or a pharmaceutically acceptable salt or solvate of said compound to the mammal.

A preferred dosage is about 0.001 to 500 mg/kg of body weight/day of the

compound of Formula III. An especially preferred dosage is about 0.01 to 25 mg/kg of body weight/day of a compound of Formula III, or a pharmaceutically acceptable salt or solvate of said compound.

The compounds of this invention may also be useful in combination

- 5 (administered together or sequentially) with one or more of anti-cancer treatments such as radiation therapy, and/or one or more anti-cancer agents selected from the group consisting of cytostatic agents, cytotoxic agents (such as for example, but not limited to, DNA interactive agents (such as cisplatin or doxorubicin)); taxanes (e.g. taxotere, taxol); topoisomerase II inhibitors (such as
- 10 etoposide); topoisomerase I inhibitors (such as irinotecan (or CPT-11), camptostar, or topotecan); tubulin interacting agents (such as paclitaxel, docetaxel or the epothilones); hormonal agents (such as tamoxifen); thymidilate synthase inhibitors (such as 5-fluorouracil); anti-metabolites (such as methotrexate); alkylating agents (such as temozolomide (TEMODAR™ from
- 15 Schering-Plough Corporation, Kenilworth, New Jersey), cyclophosphamide); Farnesyl protein transferase inhibitors (such as, SARASAR™(4-[2-[4-[(11R)-3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-yl]-1-piperidiny]-2-oxoehtyl]-1-piperidinecarboxamide, or SCH 66336 from
- 20 Schering-Plough Corporation, Kenilworth, New Jersey), tipifarnib (Zarnestra® or R115777 from Janssen Pharmaceuticals), L778,123 (a farnesyl protein transferase inhibitor from Merck & Company, Whitehouse Station, New Jersey), BMS 214662 (a farnesyl protein transferase inhibitor from Bristol-Myers Squibb Pharmaceuticals, Princeton, New Jersey); signal transduction inhibitors (such as,
- 25 Iressa (from Astra Zeneca Pharmaceuticals, England), Tarceva (EGFR kinase inhibitors), antibodies to EGFR (e.g., C225), GLEEVEC™ (C-abl kinase inhibitor from Novartis Pharmaceuticals, East Hanover, New Jersey); interferons such as, for example, intron (from Schering-Plough Corporation), Peg-Intron (from Schering-Plough Corporation); hormonal therapy combinations; aromatase combinations; ara-C, adriamycin, cytoxan, and gemcitabine.
- 30 Other anti-cancer (also known as anti-neoplastic) agents include but are not limited to Uracil mustard, Chloromethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine,

Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, oxaliplatin, leucovirin, oxaliplatin (ELOXATIN™ from Sanofi-Synthelabo Pharmaceutics, France), Pentostatine, Vinblastine, 5 Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, Mithramycin, Deoxycoformycin, Mitomycin-C, L-Asparaginase, Teniposide 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrolacetate, Methylprednisolone, Methyltestosterone, 10 Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, goserelin, Cisplatin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, Anastrozole, Letrozole, Capecitabine, Reloxafine, Droloxafine, or 15 Hexamethylmelamine.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically active agent or treatment within its dosage range. For example, the CDC2 inhibitor olomucine has been found to act synergistically with 20 known cytotoxic agents in inducing apoptosis (*J. Cell Sci.*, (1995) 108, 2897. Compounds of Formula III may also be administered sequentially with known anticancer or cytotoxic agents when a combination formulation is inappropriate. The invention is not limited in the sequence of administration; compounds of Formula III may be administered either prior to or after administration of the 25 known anticancer or cytotoxic agent. For example, the cytotoxic activity of the cyclin-dependent kinase inhibitor flavopiridol is affected by the sequence of administration with anticancer agents. *Cancer Research*, (1997) 57, 3375. Such techniques are within the skills of persons skilled in the art as well as attending physicians.

30 Accordingly, in an aspect, this invention includes combinations comprising an amount of at least one compound of Formula III, or a pharmaceutically acceptable salt or solvate thereof, and an amount of one or more anti-cancer

treatments and anti-cancer agents listed above wherein the amounts of the compounds/ treatments result in desired therapeutic effect.

The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological assays. The exemplified

5 pharmacological assays which are described later have been carried out with the compounds according to the invention and their salts.

This invention is also directed to pharmaceutical compositions which comprise at least one compound of Formula III, or a pharmaceutically acceptable salt or solvate of said compound and at least one pharmaceutically acceptable
10 carrier.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and
15 tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g., magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture
20 for various compositions may be found in A. Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for
25 parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically
30 acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations for either oral or

parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols
5 and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

The compounds of this invention may also be delivered subcutaneously.

Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in a unit dosage form. In
10 such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 1 mg to about 100 mg, preferably from about 1 mg
15 to about 50 mg, more preferably from about 1 mg to about 25 mg, according to the particular application.

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the
20 skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

The amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as
25 age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 500 mg/day, preferably 1 mg/day to 200 mg/day, in two to four divided doses.

Another aspect of this invention is a kit comprising a therapeutically
30 effective amount of at least one compound of Formula III, or a pharmaceutically acceptable salt or solvate of said compound and a pharmaceutically acceptable carrier, vehicle or diluent.

Yet another aspect of this invention is a kit comprising an amount of at least one compound of Formula III, or a pharmaceutically acceptable salt or solvate of said compound and an amount of at least one anticancer therapy and/or anti-cancer agent listed above, wherein the amounts of the two or more ingredients result in desired therapeutic effect.

The invention disclosed herein is exemplified by the following preparations and examples which should not be construed to limit the scope of the disclosure. Alternative mechanistic pathways and analogous structures will be apparent to those skilled in the art.

Where NMR data are presented, ^1H spectra were obtained on either a Varian VXR-200 (200 MHz, ^1H), Varian Gemini-300 (300 MHz) or XL-400 (400 MHz) and are reported as ppm down field from Me_4Si with number of protons, multiplicities, and coupling constants in Hertz indicated parenthetically. Where LC/MS data are presented, analyses was performed using an Applied Biosystems API-100 mass spectrometer and Shimadzu SCL-10A LC column: Altech platinum C18, 3 micron, 33mm x 7mm ID; gradient flow: 0 min – 10% CH_3CN , 5 min – 95% CH_3CN , 7 min – 95% CH_3CN , 7.5 min – 10% CH_3CN , 9 min – stop. The retention time and observed parent ion are given.

The following solvents and reagents may be referred to by their abbreviations in parenthesis:

Thin layer chromatography: TLC

dichloromethane: CH_2Cl_2

ethyl acetate: AcOEt or EtOAc

methanol: MeOH

trifluoroacetate: TFA

triethylamine: Et_3N or TEA

butoxycarbonyl: n-Boc or Boc

nuclear magnetic resonance spectroscopy: NMR

liquid chromatography mass spectrometry: LCMS

high resolution mass spectrometry: HRMS

milliliters: mL

millimoles: mmol

microliters: μ l

grams: g

milligrams: mg

room temperature or rt (ambient): about 25°C.


5 dimethoxyethane: DME

EXAMPLES

In general, the compounds described in this invention can be prepared through the general routes described below in Scheme 1. Treatment of the

10


Scheme 1



starting nitrile with potassium t-butoxide and ethyl formate gives rise to the intermediate enol 2 which upon treatment with hydrazine gives the desired substituted 3-aminopyrazole. Condensation of compounds of type 3 with the

15 appropriately functionalized keto ester of type 5 gives rise to the pyridones 6 as shown in Scheme 3. The keto esters used in this general route are either commercially available or can be made as illustrated in Scheme 2.

Scheme 2




20

The chlorides of type 9 can be prepared by treatment of the pyridones 8 with POCl₃. When R² is equal to H, substitution in this position is possible on the compounds of type 9 by electrophilic halogenation, acylation, and various other electrophilic aromatic substitutions.

Introduction of the N7-amino functionality can be accomplished through

25 displacement of the chloride of compounds of type 9 by reaction with the appropriate amine as shown in Scheme 3.

50


Scheme 3

Condensation of compounds of type 7 with the appropriately functionalized malonate ester of type 11 gives rise to the pyridones 13 as shown in Scheme 4.


The chlorides of type 14 can be prepared by treatment of the pyridones 13 with POCl₃. When R² is H, substitution in this position is possible on compounds of type 9 by electrophilic halogenation, acylation, and various other electrophilic aromatic substitutions.

Incorporation of the N7-amino functionality can be accomplished through regioselective displacement of the chloride of compounds of type 14.

Incorporation of the N5-amino functionality by addition of an appropriate amine at higher temperature.


Scheme 4

Alternatively, condensations of the aminopyrazoles of type 7 with an appropriately functionalize keto ester as prepared in Scheme 5, leads to compounds of type 13 as shown in Scheme 4.

Scheme 5

The chlorides of type 14 can be prepared by treatment of the pyridones 13 with POCl_3 . When R^2 is equal to H, substitution in this position is possible on compounds of type 14 by electrophilic halogenation, acylation, and various other electrophilic aromatic substitutions.


Incorporation of the N7-amino functionality can be accomplished through displacement of the chloride of compounds of type 15.

Preparative Examples:**PREPARATIVE EXAMPLE 1:****Step A:**

A procedure in German patent DE 19834047 A1, p 19 was followed. To a solution of KOtBu (6.17 g, 0.055 mol) in anhydrous THF (40 mL) was added, dropwise, a solution of cyclopropylacetonitrile (2.0 g, 0.025 mol) and ethyl formate (4.07 g, 0.055 mol) in anhydrous THF (4 mL). A precipitate formed immediately. This mixture was stirred for 12 hr. It was concentrated under vacuum and the residue stirred with Et_2O (50 mL). The resulting residue was decanted and washed with Et_2O (2 x 50 mL) and Et_2O removed from the residue under vacuum. The residue was dissolved in cold H_2O (20 mL) and pH adjusted to 4 – 5 with 12 N HCl. The mixture was extracted with CH_2Cl_2 (2 x 50 mL). The

organic layers were combined, dried over MgSO_4 and concentrated under vacuum to give the aldehyde as a tan liquid.

Step B:




- 5 The product from Preparative Example 1, Step A (2.12 g, 0.0195 mol), $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ (1.95 g, 0.039 mol) and 1.8 g (0.029 mole) of glacial $\text{CH}_3\text{CO}_2\text{H}$ (1.8 g, 0.029 mol) were dissolved in EtOH (10 mL). It was refluxed for 6 hr and concentrated under vacuum. The residue was slurried in CH_2Cl_2 (150 mL) and the pH adjusted to 9 with 1N NaOH. The organic layer was washed with brine,
- 10 dried over MgSO_4 and concentrated under vacuum to give the product as a waxy orange solid.

PREPARATIVE EXAMPLES 2-4:


- By essentially the same procedure set forth in Preparative Example 1, only substituting the nitrile shown in Column 2 of Table 2, the compounds in
- 15 Column 3 of Table 2 were prepared:

TABLE 2

Prep. Ex.	Column 2	Column 3
2		
3		
3.10		

PREPARATIVE EXAMPLE 4


2-Carbomethoxycyclopentanone (6.6 ml, 0.05 mol) in THF (15 ml) was added dropwise to a vigorously stirred suspension of NaH (60% in mineral oil, 4 g, 0.1 mol) in THF (100 ml) at 0 – 10 °C. When bubbling ceased, the reaction mixture was treated at the same temperature with ClCOOMe (7.8 ml, 0.1 mol) in THF (15 ml). The resulted off-white suspension was stirred for 30 minutes at room temperature and 30 minutes under reflux. The reaction was monitored by TLC for disappearance of starting material. The reaction mixture was quenched with water carefully and partitioned between ethyl acetate and saturated solution of ammonium chloride in a funnel. Shaken and separated, the organic layer was washed with brine and dried over anhydrous sodium sulfate. Solvents were removed, and the residue was purified by flash chromatography, eluted with 5% and then 10% ethyl acetate in hexane. 9.4 g colorless oil was obtained with 94% yield. ¹H NMR (CDCl₃) δ 3.90(s, 3H), 3.73(s, 3H), 2.65(m, 4H), 1.98(m, 2H).

PREPARATIVE EXAMPLE 5

To lithium diisopropylamide solution in THF (2.0 N, 0.04 mol) at –65 °C, was added dropwise 2,2-dicarbomethoxycyclopentanone (4 g, 0.02 mol) in THF (60 ml). The resulted reaction mixture was stirred at the same temperature before adding methyl chloroformate (1.54 ml, 0.02 mol). Reaction mixture stirred for an hour and poured into saturated ammonium chloride solution with some ice. This solution was extracted three times with ether, and the combined ethereal layers were dried over sodium sulfate. Solvents were removed in vacuo, and the residue was purified by flash chromatography, eluted with 30%

increased to 50% ethyl acetate in hexane. 2.3 g yellowish oil was obtained with 58% yield. ^1H NMR (CDCl_3) δ 3.77(s, 6H), 3.32(t, 1H), 3.60-3.10(m, 4H).

PREPARATIVE EXAMPLE 6:

























- 5 The reactions were done as outlined in (K. O. Olsen, *J. Org. Chem.*, (1987) 52, 4531 – 4536). Thus, to a stirred solution of lithium diisopropylamide in THF at -65 to -70°C was added freshly distilled ethyl acetate, dropwise. The resulting solution was stirred for 30 min and the acid chloride was added as a solution in THF. The reaction mixture was stirred at -65 to -70°C for 30 min
10 and then terminated by the addition of 1 N HCl solution. The resulting two-phased mixture was allowed to warm to ambient temperature. The resulting mixture was diluted with EtOAc (100 mL) the organic layer was collected. The aqueous layer was extracted with EtOAc (100 mL). The organic layers were combined, washed with brine, dried (Na_2SO_4), and concentrated *in vacuo* to give
15 the crude β -keto esters, which were used in the subsequent condensations.

PREPARATIVE EXAMPLES 7-19:

By essentially the same procedure set forth in Preparative Example 6 only substituting the acid chlorides shown in Column 2 of Table 3, the β -keto esters shown in Column 3 of Table 3 were prepared:


TABLE 3

Prep. Ex.	Column 2	Column 3	DATA
7			LCMS: $\text{MH}^+ = 223$

8			LCMS: $MH^+ = 253$
9			LCMS: $MH^+ = 261$
10			$MH^+ = 199$
11			
12			
13			LCMS: $MH^+ = 271$
14			Yield = quant $MH^+ = 249$
15			Yield = quant $MH^+ = 237$
16			Yield = quant $MH^+ = 262$
17			Yield = 48 $MH^+ = 195$
18			Yield = 99 $MH^+ = 199$

56

19			Yield=77% ¹ H NMR (CDCl ₃) δ 7.42(t, 1H), 6.68(d, 2H), 4.29(q, 2H), 3.97(d, 2H), 3.95(s, 3H), 1.38(t, 3H).
----	--	--	--

PREPARATIVE EXAMPLE 20:

- To a solution of the acid in THF was added Et₃N, followed by isobutyl chloroformate at -20 to -30°C. After the mixture was stirred for 30 min at -20 to -30°C, triethylamine hydrochloride was filtered off under argon, and the filtrate was added to the LDA-EtOAc reaction mixture (prepared as outlined in Method A) at -65 to -70°C. After addition of 1 N HCl, followed by routine workup of the reaction mixture and evaporation of the solvents, the crude β-keto esters were isolated. The crude material was used in the subsequent condensations.











PREPARATIVE EXAMPLES 21 - 28:


- By essentially the same conditions set forth in Preparative Example 20 only substituting the carboxylic acid shown in Column 2 of Table 4, the compounds shown in Column 3 of Table 4 were prepared:

TABLE 4

Prep. Ex.	Column 2	Column 3	CMPD
21			Yield = 99% MH ⁺ = 213
22			Yield = 70% MH ⁺ = 275
23			Yield = quant MH ⁺ = 213

57

24			Yield = quant MH ⁺ = 211
25			Yield = 99 MH ⁺ = 334
26			Yield = 99 MH ⁺ = 334
27			Yield = 99 MH ⁺ = 334
28			Yield=77% ¹ H NMR (CDCl ₃) δ 4.21(q, 2H), 3.95(d, 2H), 3.93-3.79(m, 4H), 3.52(s, 2H), 2.65(m, 1H), 1.25(t, 3H), 1.23-1.2(m, 2H).

PREPARATIVE EXAMPLE 29:



















- 5 A solution of 3-aminopyrazole (2.0g, 24.07 mmol) and ethyl benzoylacetate (4.58 mL, 1.1 eq.) in AcOH (15 mL) was heated at reflux for 3 hours. The reaction mixture was cooled to room temperature and concentrated *in vacuo*. The resulting solid was diluted with EtOAc and filtered to give a white solid (2.04 g, 40% yield).

























10 **PREPARATIVE EXAMPLES 30-73:**




























By essentially the same procedure set forth in Preparative Example 29 only substituting the aminopyrazole shown in Column 2 of Table 5 and the ester




























shown in Column 3 of Table 5, the compounds shown in Column 4 of Table 5 were prepared:




























TABLE 5













Prep. Ex.	Column 2	Column 3	Column 4	Column 5
30				
31				
32				
33				
34				
35				


36				
37				
37.10				
38				
39				
40				
41				
42				

43				
44				
45				
46				
47				
48				
49				
50				
51				


52				
53				
54				
55				
56				
57				
58				Yield = 68 MH ⁺ = 152
59				Yield = 46 MH ⁺ = 268
60				Yield = 63 MH ⁺ = 255

61				Yield = 80 MH ⁺ = 280
62				Yield =72 MH ⁺ = 214
63				Yield = 51 MH ⁺ = 218
64				Yield = 82 MH ⁺ =218
65				Yield = 39 MH ⁺ =232
66				Yield = 30 MH ⁺ =230
67				Yield = 80 MH ⁺ =353
68				Yield = 49 MH ⁺ =353
69				Yield = 42 MH ⁺ =353

70				
71				
72				
73				

PREPARATIVE EXAMPLE 74:

5 Ethyl benzoylacetate (1.76 mL, 1.1 eq.) and 3-amino-4-cyanopyrazole (1.0 g, 9.25 mmol) in AcOH (5.0 mL) and H₂O (10 mL) was heated at reflux 72 hours. The resulting solution was cooled to room temperature, concentrated *in vacuo*, and diluted with EtOAc. The resulting precipitate was filtered, washed with EtOAc, and dried *in vacuo* (0.47 g, 21% yield).

PREPARATIVE EXAMPLE 75







10 A procedure in US patent 3,907,799 was followed. Sodium (2.3 g, 2 eq.) was added to EtOH (150 mL) portionwise. When the sodium was completely

dissolved, 3-aminopyrazole (4.2 g, 0.05 mol) and diethyl malonate (8.7 g, 1.1 eq.) were added and the resulting solution heated to reflux for 3 hours. The resulting suspension was cooled to room temperature and filtered. The filter cake was washed with EtOH (100 mL) and dissolved in water (250 mL). The resulting solution was cooled in an ice bath and the pH adjusted to 1-2 with concentrated HCl. The resulting suspension was filtered, washed with water (100 mL) and dried under vacuum to give a white solid (4.75 g, 63% yield).

PREPARATIVE EXAMPLES 76-78:

By essentially the same procedure set forth in Preparative Example 75 only substituting the compound shown in Column 2 of Table 6, the compounds shown in Column 3 of Table 6 are prepared:

TABLE 6

Prep. Ex.	Column 2	Column 3
76		
77		
78		

PREPARATIVE EXAMPLE 79:











A solution of the compound prepared in Preparative Example 29 (1.0 g, 4.73 mmol) in POCl₃ (5 mL) and pyridine (0.25 mL) was stirred at room

















temperature 3 days. The resulting slurry was diluted with Et₂O, filtered, and the solid residue washed with Et₂O. The combined Et₂O washings were cooled to 0°C and treated with ice. When the vigorous reaction ceased, the resulting mixture was diluted with H₂O, separated, and the aqueous layer extracted with Et₂O. The combined organics were washed with H₂O and saturated NaCl, dried over Na₂SO₄, filtered, and concentrated to give a pale yellow solid (0.86 g, 79% yield). LCMS: MH⁺=230.

















PREPARATIVE EXAMPLE 80-122:

By essentially the same procedure set forth in Preparative Example 79, only substituting the compound shown in Column 2 of Table 7, the compounds shown in Column 3 of Table 7 were prepared:

















TABLE 7

Prep. Ex.	Column 2	Column 3	CMPD
80			MS: MH ⁺ =248
81			
82			MS: MH ⁺ =298
83			MS: MH ⁺ =196

84			MS: $MH^+ = 210$
85			
86			MS: $MH^+ = 272$
87			
87.10			
88			MS: $MH^+ = 255$
89			
90			Yield = 65% MS: $MH^+ = 260$


91			Yield = 35% MS: MH ⁺ = 290
92			Yield = 32% MS: MH ⁺ = 298
93			Yield = 45% MS: MH ⁺ = 236
94			Yield = 100% LCMS: MH ⁺ = 250
95			Yield = 88% MS: MH ⁺ = 314
96			Yield=43% MS: MH ⁺ =223
97			Yield=30% MS: MH ⁺ =295
98			Yield=98% MS: MH ⁺ =244

99			
100			
101			
102			
103			
104			
105			
106			
107			45% yield; MS: MH ⁺ =226
108			MS: MH ⁺ = 308

109			Yield = quant MH ⁺ = 286
110			Yield = 50 MH ⁺ = 272
111			Yield = 85 MH ⁺ = 299
112			Yield = 97 MH ⁺ = 231
113			Yield = 45 MH ⁺ = 236
114			Yield = quant. MH ⁺ = 236
115			Yield = 57 MH ⁺ = 250
116			Yield = 89 MH ⁺ = 248

70

117			Yield = 96 MH ⁺ = 371
118			Yield = 99 MH ⁺ = 371
119			Yield = 50 MH ⁺ = 371
120			Yield=57% LCMS: MH ⁺ =224
121			Yield=34% LCMS: MH ⁺ =226
122			Yield=100% ¹ H NMR (CDCl ₃) δ 8.53(d, 1H), 7.66(t, 1H), 7.51(s, 1H), 7.45(d, 1H), 6.84(d, 2H).

PREPARATIVE EXAMPLE 123







5 POCl₃ (62 mL) was cooled to 5 °C under nitrogen and dimethylaniline (11.4 g, 2.8 eq.) and the compound prepared in Preparative Example 75 (4.75 g,

0.032 mol). The reaction mixture was warmed to 60 °C and stirred overnight. The reaction mixture was cooled to 30 °C and the POCl₃ was distilled off under reduced pressure. The residue was dissolved in CH₂Cl₂ (300 mL) and poured onto ice. After stirring 15 minutes, the pH of the mixture was adjusted to 7-8 with solid NaHCO₃. The layers were separated and the organic layer was washed with H₂O (3 x 200 mL), dried over MgSO₄, filtered, and concentrated. The crude product was purified by flash chromatography using a 50 : 50 CH₂Cl₂ : hexanes solution as eluent to elute the dimethyl aniline. The eluent was then changed to 75 : 25 CH₂Cl₂ : hexanes to elute the desired product (4.58 g, 77% yield). MS: MH⁺=188.

PREPARATIVE EXAMPLES 124-126


By essentially the same procedure set forth in Preparative Example 123 only substituting the compound in Column 2 of Table 8, the compounds shown in Column 3 of Table 8 are prepared:

TABLE 8

Prep. Ex.	<u>Column 2</u>	<u>Column 3</u>
124		
125		
126		

PREPARATIVE EXAMPLE 127:

72



















A solution of the compound prepared in Preparative Example 79 (0.10 g, 0.435 mmol) in CH₃CN (3 mL) was treated with NBS (0.085 g, 1.1 eq.). The reaction mixture was stirred at room temperature 1 hour and concentrated under reduced pressure. The crude product was purified by flash chromatography using a 20% EtOAc-in-hexanes solution as eluent (0.13 g, 100% yield). LCMS:MH⁺=308.



















PREPARATIVE EXAMPLES 128-164:



















By essentially the same procedure set forth in Preparative Example 127 only substituting the compounds shown in Column 2 of Table 9, the compounds shown in Column 3 of Table 9 were prepared:

















TABLE 9

Prep. Ex.	Column 2	Column 3	CMPD
128			MS: MH ⁺ = 326
129			MS: MH ⁺ = 342
130			MS: MH ⁺ = 376

131			MS: MH ⁺ =274
132			MS: MH ⁺ =288
133			
134			Yield = 75% MS: MH ⁺ = 338
135			Yield = 52% MS: MH ⁺ = 368
136			Yield = 87% MS: MH ⁺ = 376
137			Yield = 100% MS: MH ⁺ = 316
138			Yield = 92% MS: MH ⁺ = 330

139			Yield = 82% MS: MH ⁺ = 395
140			Yield=88% MS: MH ⁺ =308
141			Yield=100% MS: MH ⁺ =322
142			MH ⁺ =266
143			
144			
145			
146			
147			

148			
149			
150			
151			LCMS: MH ⁺ = 386
152			Yield = quant MH ⁺ = 364
153			Yield = quant MH ⁺ = 353
154			Yield = 95 MH ⁺ = 378
155			Yield = 77 MH ⁺ = 311
156			Yield = quant. MH ⁺ = 314

157			Yield = 99 MH ⁺ = 328
158			Yield = 98 MH ⁺ = 326
159			Yield = 99 MH ⁺ = 449
160			Yield = 95 MH ⁺ = 449
161			Yield = 72 MH ⁺ = 449
162			Yield=98% LCMS: MH ⁺ =302
163			Yield=95% LCMS: MH ⁺ =305
164			Yield=50% ¹ H NMR (CDCl ₃) δ 8.36(s, 1H), 7.72(d, 1H), 7.20(s, 1H), 6.82(d, 1H), 3.99(s, 3H), 3.90(s, 3H);

PREPARATIVE EXAMPLE 165:

77



A solution of the compound prepared in Preparative Example 80 (0.3 g, 1.2 mmol) in CH₃CN (15 mL) was treated with NCS (0.18 g, 1.1 eq.) and the resulting solution heated to reflux 4 hours. Additional NCS (0.032 g, 0.2 eq.) added and the resulting solution was stirred at reflux overnight. The reaction mixture was cooled to room temperature, concentrated *in vacuo* and the residue purified by flash chromatography using a 20% EtOAc in hexanes solution as eluent (0.28 g, 83% yield). LCMS: MH⁺ = 282.

PREPARATIVE EXAMPLE 166-167:

- 10 By essentially the same procedure set forth in Preparative Example 165 only substituting the compound shown in Column 2 of Table 10, the compound shown in Column 3 of Table 10 was prepared:

TABLE 10

Prep. Ex.	Column 2	Column 3	CMPD
166			Yield = 82% LCMS: MH ⁺ = 286
167			

15 PREPARATIVE EXAMPLE 167.10:




By essentially the same procedure set forth in Preparative Example 165 only substituting N-iodosuccinimide, the above compound was prepared.

78

PREPARATIVE EXAMPLE 168:

To a solution of the compound from Preparative Example 79 (1.0 g, 4.35 mmol) in DMF (6 mL) was added POCl₃ (1.24 mL, 3.05 eq.) and the resulting mixture was stirred at room temperature overnight. The reaction mixture was cooled to 0°C and the excess POCl₃ was quenched by the addition of ice. The resulting solution was neutralized with 1N NaOH, diluted with H₂O, and extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography using a 5% MeOH in CH₂Cl₂ solution as eluent (0.95 g, 85% yield). LCMS: MH⁺=258.

PREPARATIVE EXAMPLE 169:


By essentially the same procedure set forth in Preparative Example 168 only substituting the compound prepared in Preparative Example 80, the above compound was prepared (0.45 g, 40% yield).

PREPARATIVE EXAMPLE 170:

To a solution of the product of Preparative Example 169 (0.25 g, 0.97 mmol) in THF was added NaBH₄ (0.041 g, 1.1 eq.) and the resulting solution was stirred at room temperature overnight. The reaction mixture was quenched

by the addition of H₂O and extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude product was purified by flash chromatography using a 60: 40 hexanes : EtOAc mix as eluent (0.17 g, 69% yield). MS: MH⁺=260.


5 PREPARATIVE EXAMPLE 171:



A solution of the compound prepared in Preparative Example 170 (0.12 g, 0.462 mmol), dimethyl sulfate (0.088 mL, 2.0 eq), 50% NaOH (0.26 mL) and catalytic Bu₄NBr in CH₂Cl₂ (4 mL) was stirred at room temperature overnight.

- 10 The reaction mixture was diluted with H₂O and extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude product was purified by flash chromatography using a 30% EtOAc-in-hexanes solution as eluent (0.062 g, 48% yield).

PREPARATIVE EXAMPLE 172




15

To a solution of PPh₃ (4.07 g, 4.0 eq.) and CBr₄ (2.57 g, 2.0 eq.) in CH₂Cl₂ (75 mL) at 0 °C was added the compound prepared in Preparative Example 168 (1.0 g, 3.88 mmol). The resulting solution was stirred at 0 °C for 1 hour and concentrated under reduced pressure. The residue was purified by


20 flash chromatography using a 20% EtOAc in hexanes solution as eluent (1.07 g, 67% yield).

PREPARATIVE EXAMPLE 173:



By essentially the same procedure set forth in Preparative Example 172 only substituting the compound prepared in Preparative Example 169 the above compound was prepared (0.5 g, 70% yield).

PREPARATIVE EXAMPLE 174:



The compound prepared in Preparative Example 127 (3.08 g, 10.0 mmol), 2.0 M NH_3 in 2-propanol (50 mL, 100.0 mmol), and 37 % aqueous NH_3 (10.0 mL) were stirred in a closed pressure vessel at 50°C for 1 day. The solvent was evaporated and the crude product was purified by flash chromatography using 3:1 CH_2Cl_2 :EtOAc as eluent. Pale yellow solid (2.30 g, 80%) was obtained. LCMS: $\text{M}^+ = 289$.









PREPARATIVE EXAMPLES 175-180:


By essentially the same procedure set forth in Preparative Example 174 only substituting the compound shown in Column 2 of Table 11, the compounds shown in Column 3 of Table 11 were prepared.

TABLE 11

Prep. Ex.	Column 2	Column 3
175		
176		


81

177		
178		
179		
180		


PREPARATIVE EXAMPLE 181:

The compound prepared in Preparative Example 80 (0.3 g, 1.2 mmol),
 5 K_2CO_3 (0.33 g, 2 eq.), and 4-aminomethylpyridine (0.13 mL, 1.1 eq.) was heated to reflux overnight. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The residue was diluted with H_2O and extracted with CH_2Cl_2 . The combined organics were dried over Na_2SO_4 , filtered and, concentrated. The crude product was purified by flash chromatography
 10 using a 5% (10% NH_4OH in MeOH) solution in CH_2Cl_2 as eluent (0.051 g, 40% yield). LCMS: $MH^+=320$.

82

PREPARATIVE EXAMPLE 182:

By essentially the same procedure set forth in Preparative Example 181 only substituting the compound described in Preparative Example 92, the above compound was prepared. LCMS: $MH^+ = 370$.

PREPARATIVE EXAMPLE 183:











To a solution of the compound prepared in Preparative Example 123 (0.25 g, 1.3 mmol) in dioxane (5 mL) was added iPr_2NEt (0.47 mL, 2.0 eq.) and 3-aminomethylpyridine (0.15 mL, 1.1 eq.). The resulting solution was stirred at room temperature 72 hours. The reaction mixture was diluted with H_2O and extracted with EtOAc. The combined organics were washed with H_2O and saturated NaCl, dried over Na_2SO_4 , filtered, and concentrated in vacuo. The crude product was purified by flash chromatography using a 5% MeOH in CH_2Cl_2 solution as eluent (0.29 g, 83% yield). MS: $MH^+ = 260$.





PREPARATIVE EXAMPLES 184-187:


By essentially the same procedure set forth in Preparative Example 183 only substituting the compound shown in Column 2 of Table 12, the compounds shown in Column 3 of Table 12 are prepared.

TABLE 12

Prep. Ex.	<u>Column 2</u>	<u>Column 3</u>

184		
184.1		
185		
186		
187		

187.1		
187.11		


PREPARATIVE EXAMPLE 188 and PREPARATIVE EXAMPLE 189:

To a solution of the compound prepared in Preparative Example 185
 5 (1.18 g, 3.98 mmol) in THF (35 mL) at -78 °C was added LAH (4.78 mL, 1M in Et₂O, 1.0 eq.) dropwise. The reaction mixture was stirred at -78 °C for 3 hours at which time additional LAH (2.0 mL, 1M in Et₂O, 0.42 eq.) was added dropwise. The reaction mixture was stirred an additional 1.25 hours and quenched by the addition of saturated Na₂SO₄ (8.5 mL). The reaction mixture was diluted with
 10 EtOAc (23 mL), H₂O (2 mL), and CH₃OH (50 mL). The resulting slurry was filtered through a plug of Celite. The Celite was washed with CH₃OH and the filtrate dried with Na₂SO₄, filtered, and concentrated. The product was purified by flash chromatography using a CH₂Cl₂ : CH₃OH (93 : 7) solution as eluent to yield aldehyde as the first eluting product and alcohol as the second eluting
 15 product.

Preparative Example 188: (aldehyde): 0.4 g, 39% yield. MS: MH⁺ = 254.


Preparative Example 189: (alcohol): 0.25 g, 24% yield. MS: $MH^+ = 256$.

PREPARATIVE EXAMPLE 190:



- 5 To a solution of the compound prepared in Preparative Example 188 (0.075 g, 0.30 mmol) in THF (2.0 mL) at 0 °C was added CH_3MgBr (0.3 mL, 3.0M solution in Et_2O , 3.0 eq.) dropwise. The resulting solution was stirred at 0 °C an additional 1.5 hours, warmed to room temperature, and stirred overnight. Additional CH_3MgBr (0.15 mL, 3.0M in Et_2O , 1. eq.) was added and the resulting solution stirred an additional 1.5 hours. The reaction mixture was cooled to 0 °C and quenched by the addition of saturated NH_4Cl . The resulting solution was diluted with CH_2Cl_2 and H_2O and extracted with CH_2Cl_2 . The combined organics were washed with saturated $NaCl$ and dried over Na_2SO_4 , filtered, and concentrated. The crude product was purified by flash chromatography using a $CH_2Cl_2 : CH_3OH$ (90 : 10) solution as eluent (0.048 g, 60% yield). MS: $MH^+ = 270$.
- 10
- 15


PREPARATIVE EXAMPLE 191:



- By essentially the same procedure set forth in Preparative Example 190 only substituting the compound prepared in Preparative Example 185 and using excess $MeMgBr$ (5 eq.), the above compound was prepared.
- 20


PREPARATIVE EXAMPLE 192:

86



The compound prepared in Preparative Example 181 (0.29 g, 0.91 mmol), BOC₂O (0.22 g, 1.1 eq), and DMAP (0.13 g, 1.1 eq.) in dioxane (10 mL) was stirred at room temperature 3 days. Additional BOC₂O (0.10g, 0.5 eq.) was added and the reaction mixture was stirred 4 hours. The reaction mixture was concentrated *in vacuo*, diluted with saturated NaHCO₃ (15 mL), and extracted with CH₂Cl₂ (2 x 100 mL). The combined organics were dried over Na₂SO₄, filtered, and concentrated under reduce pressure. The crude product was purified by flash chromatography using a 5% (10% NH₄OH in MeOH) solution in CH₂Cl₂ as eluent (0.35 g, 91% yield). LCMS: MH⁺ = 420.


15 PREPARATIVE EXAMPLE 193:



By essentially the same procedure set forth in Preparative Example 192 only substituting the compound prepared in Preparative Example 183, the above compound was prepared. MS: MH⁺ = 360.

20 PREPARATIVE EXAMPLE 193.10:


87



By essentially the same procedure set forth in Preparative Example 192 only substituting the compound prepared in Preparative Example 184.1, the above compound was prepared. MS: $MH^+ = 454$.


5

PREPARATIVE EXAMPLE 194:




By essentially the same procedure set forth in Preparative Example 192 only substituting the above compound prepared in Preparative Example 187.11, the above compound was prepared (0.223 g, 88% yield). MS: $MH^+ = 528$.

PREPARATIVE EXAMPLE 195:




15


By essentially the same procedure set forth in Preparative Example 127 only substituting the compound prepared in Preparative Example 192, the above compound was prepared (0.38 g, 95% yield). LCMS: $MH^+ = 498$.

PREPARATIVE EXAMPLE 196:

By essentially the same procedure set forth in Preparative Example 195, only substituting the compound prepared in Preparative Example 193, the above compound was prepared (0.3 g, 83% yield). MS: $MH^+ = 438$.

PREPARATIVE EXAMPLE 197:


A solution of the compound prepared in Preparative Example 195 (0.15 g, 0.3 mmol), phenylboronic acid (0.073 g, 2.0 eq.), K_3PO_4 (0.19 g, 3.0 eq.), and $Pd(PPh_3)_4$ (0.017 g, 5 mol %) was heated at reflux in DME (16 mL) and H_2O (4 mL) 7 hours. The resulting solution was cooled to room temperature, diluted with H_2O (10 mL), and extracted with CH_2Cl_2 (3 x 50 mL). The combined organics were dried over Na_2SO_4 , filtered, and concentrated. The crude product was purified by flash chromatography using a 2.5% (10% NH_4OH in MeOH) in CH_2Cl_2 solution as eluent (0.16 g, 100% yield).

PREPARATIVE EXAMPLE 198:

To a solution of 4-aminomethylpyridine (1.41 mL, 13.87 mmol) in CH_2Cl_2 (50 mL) was added BOC_2O (3.3 g, 1.1 eq.) and TEA and the resulting solution


was stirred at room temperature 2 hours. The reaction mixture was diluted with H₂O (50 mL) and extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by flash chromatography using a 5% (10% NH₄OH in MeOH) solution in CH₂Cl₂ as eluent to give a yellow solid (2.62 g, 91% yield). LCMS: MH⁺ = 209.

PREPARATIVE EXAMPLE 199:



- By essentially the same procedure set forth in Preparative Example 198 only substituting 3-aminomethylpyridine, the above compound was prepared as a yellow oil (2.66 g, 92% yield). LCMS: MH⁺ = 209.

PREPARATIVE EXAMPLE 200:




- To a solution of the compound prepared in Preparative Example 198 (0.20 g, 0.96 mmol) in CH₂Cl₂ (5 mL) at 0°C was added *m*-CPBA (0.17 g, 1.0 eq) and the resulting solution stirred at 0°C 2 hours and stored at 4°C overnight at which time the reaction mixture was warmed to room temperature and stirred 3 hours. The reaction mixture was diluted with H₂O and extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by flash chromatography using a 10% (10% NH₄OH in MeOH) solution as eluent: LCMS: MH⁺ = 255.

PREPARATIVE EXAMPLE 201:




A solution of oxone (58.6 g) in H₂O (250 mL) was added dropwise to the compound prepared in Preparative Example 199 (27 g, 0.13 mol) and NaHCO₃ (21.8 g, 2.0 eq.) in MeOH (200 mL) and H₂O (250 mL). The resulting solution was stirred at room temperature overnight. The reaction mixture was diluted with CH₂Cl₂ (500 mL) and filtered. The layers were separated and the aqueous layer extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give a white solid (21.0 g, 72% yield). MS: MH⁺= 255.

PREPARATIVE EXAMPLE 202:



The compound prepared in Preparative Example 200 (0.29 g, 1.29 mmol) was stirred at room temperature in 4M HCl in dioxane (0.97 mL) 2 hours. The reaction mixture was concentrated *in vacuo* and used without further purification. LCMS: MH⁺= 125.


PREPARATIVE EXAMPLE 203:



By essentially the same procedure set forth in Preparative Example 202 only substituting the compound prepared in Preparative Example 201, the compound shown above was prepared. LCMS: MH⁺= 125.


PREPARATIVE EXAMPLE 204:

91



To 4-N-t-Butoxycarbonylaminopiperidine (0.8 g, 4.0 mmol) in CH₂Cl₂ (10 mL) at 0°C was added TEA (1.40 mL, 2.5 eq.) and 3-trifluoromethyl benzoyl chloride (1.05 g, 1.25 eq.). The resulting solution was stirred 15 minutes and warmed to room temperature and stirred 3 hours. The reaction mixture was diluted with CH₂Cl₂ and washed with 5% Na₂CO₃ (2 x 100 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated to yield a pale yellow solid (quantitative crude yield).


PREPARATIVE EXAMPLE 205:



To a solution of the compound prepared in Preparative Example 204 (1.0 g, 2.76 mmol) in CH₂Cl₂ (15 mL) at 0°C was added TFA (8 mL) and the resulting solution was stirred at 0°C for 30 minutes and room temperature 1 hour. The reaction mixture was poured onto Na₂CO₃ (40 g) and H₂O (400 mL) added and the resulting mixture was extracted with CH₂Cl₂. The combined organics were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude product was purified by flash chromatography using a 20% (7N NH₃ in MeOH) solution in CH₂Cl₂ as eluent (0.6 g, 82% yield).

PREPARATIVE EXAMPLES 206:

92

**STEP A:**

To a solution of 6-chloronicotinamide (1g, 6.39 mmol) in isoamyl alcohol (15 mL) at rt was added Na₂CO₃ (0.81g, 7.67 mmol) followed by methoxyethylamine (0.67 mL, 7.67 mmol). The mixture was heat at 130 °C for 16h, cooled to rt, and was filtered thru a medium glass-fritted filter. The resulting filtrate was concentrated under reduced pressure and the resultant solid was triturated with Et₂O (2 x 10 mL). The crude solid was placed under high vacuum to afford 1.2 g (96%) of a light yellow solid. M+H = 196.











STEP B:


To a solution of amide (1.2 g, 6.12 mmol) from Preparative Example 206, Step A in THF (5 mL) at 0 °C was added a solution of BH₃-THF (43 mL; 43 mmol) dropwise over 10 min. The resultant solution was warmed to rt and stirred for 14 h. The mixture was cooled to 0 °C and was sequentially treated with 6M HCl (35 mL), water (30 mL), and MeOH (150 mL). The mixture was stirred for 8 h and was concentrated under reduced pressure. The crude residue was triturated with MeOH, concentrated under reduced pressure, and placed under high vacuum to afford 1.6 g (82%) of a white solid as the dihydrochloride salt. M+H (free base) = 182.0. This material was used crude in the coupling with 7-Cl adducts.

PREPARATIVE EXAMPLES 207-211:


By essentially the same known procedure set forth in Preparative Example 206 only by utilizing the amines shown in Column 2 of Table 13 and the amines shown in Column 3 of Table 13 were prepared:

TABLE 13


Prep.Ex.	Column 2 (Amine)	Column 3 (Amine)	CMPD M+H (free base)
207		 · 2 HCl	M+H = 138
208		 · 2 HCl	M+H = 152
209		 · 2 HCl	M+H = 178
210		 · 3 HCl	M+H = 195
211		 · 3 HCl	M+H = 207

5 PREPARATIVE EXAMPLE 212:

The above compound was prepared accordingly to the methods
10 described in WO 91/18904.

PREPARATIVE EXAMPLE 213:


The above compound was prepared accordingly to the methods
15 described in US 6,180,627 B1.

PREPARATIVE EXAMPLE 214:

The known amine was prepared as described in *J. Med. Chem.* (2001), 44, 4505-4508.


PREPARATIVE EXAMPLE 215:

5



The known amine was prepared as described in *J. Med. Chem.* (1997), 40, 3726-3733.

10 PREPARATIVE EXAMPLE 216:



STEP A:

15 A solution of aldehyde (50 g, 0.41 mol) [WO 0232893] in MeOH (300 mL) was cooled to 0 °C and carefully treated with NaBH₄ (20g, 0.53 mol in 6 batches) over 20 minutes. The reaction was then allowed to warm to 20 °C and was stirred for 4 hours. The mixture was again cooled to 0 °C, carefully quenched with saturated aqueous NH₄Cl, and concentrated. Flash chromatography (5–
20 10% 7N NH₃-MeOH/CH₂Cl₂) provided the primary alcohol (31g, 62%) as a light yellow solid.


STEP B:

A slurry of alcohol (31 g, 0.25 mol) from Preparative Example 216, Step A in CH₂Cl₂ (500 mL) was cooled to 0 °C and slowly treated with SOCl₂ (55mL, 0.74 mol over 30 minutes). The reaction was then stirred overnight at 20 °C.
25 The material was concentrated, slurried in acetone, and then filtered. The resulting beige solid was dried overnight in vacuo (38.4g, 52%, HCl salt).

STEP C:


To a 15 mL pressure tube charged with a stir bar was added chloride (150 mg, 0.83 mmol) from Preparative Example 216, Step B followed by 7 M NH_3/MeOH (10 mL). The resulting solution was stirred for 48 h at rt where upon the mixture was concentrated under reduced pressure to afford a light yellow solid (0.146 g, 83%). $M+H$ (free base) = 140.

PREPARATIVE EXAMPLE 217:




The above compound was prepared accordingly to methods described in WO 00/26210.

PREPARATIVE EXAMPLE 218:




The above compound was prepared accordingly to methods described in WO 99/10325.

PREPARATIVE EXAMPLE 219:



The known amine dihydrochloride was prepared according to methods described in WO 02/64211.

PREPARATIVE EXAMPLE 220:



The above compound was prepared according to methods described in WO 02/64211.

PREPARATIVE EXAMPLE 221:


96



The known primary alcohol was prepared according to WO 00/37473 and was converted to the desired amine dihydrochloride in analogous fashion as

- 5 **Preparative Example 220** according to WO 02/064211.

PREPARATIVE EXAMPLE 222:



10 **STEP A:**

To a solution of aldehyde (WO 02/32893) (0.46 g, 2.07 mmol) in MeOH/THF (2 mL/2 mL) at 0 °C was added NaBH₄ (94 mg, 2.48 mmol) in one portion. The resulting mixture was stirred for 12 h at rt and was diluted with sat. aq. NH₄Cl (3 mL). The mixture was concentrated under reduced pressure and the resultant aqueous layer was extracted with CH₂Cl₂ (3 x 5 mL). The organic layers were combined, washed with brine (1 x 5 mL), dried (Na₂SO₄), and filtered. The organic layer was concentrated under reduced pressure to afford 417 mg (90% yield) of a white solid. M+H = 225.

20 **STEP B:**

The crude alcohol from Preparative Example 222, step A (0.4 g, 1.78 mmol) in CH₂Cl₂ (4 mL) was added SOCl₂ (0.65 mL, 8.91 mmol) and the mixture was stirred for 2 h at rt. The mixture was concentrated under reduced pressure to afford 407 mg (94%) of a light yellow solid. M+H = 243. The crude product was taken on without further purification.

25 **STEP C:**


To a solution of crude chloride from Preparative Example 222, Step B (0.33 g, 1.36 mmol) in a pressure tube charged with 7M NH_3/MeOH (35 mL) and the mixture was stirred for 72 h. The mixture was concentrated under reduced pressure to afford 257 mg (85%) of a yellow semisolid. $\text{M}+\text{H}$ (free base) = 224.

5 PREPARATIVE EXAMPLE 223:



10 To a round bottom flask charged with amine hydrochloride (0.24 g, 1.1 mmol) from Preparative Example 222 and a stir bar was added 4N HCl/dioxane (10 mL). The resulting solution was stirred for 12h at rt, concentrated under reduced pressure, and triturated with CH_2Cl_2 (3 x 5 mL). The crude product was filtered, washed with Et_2O (2 x 5 mL), and dried under high vacuum to afford 0.19g (91%) as the dihydrochloride salt. $\text{M}+\text{H}$ (free base) = 124.

15 PREPARATIVE EXAMPLE 224:











20 $\text{Pd}(\text{PPh}_3)_4$ (0.404 gm, 0.35 mmol) was added to a degassed solution of 4-cyanobenzene boronic acid (1.029 g, 7 mmol) and 2-bromopyridine (1.11 g, 7 mmol) in 75 mL acetonitrile. 0.4 M sodium carbonate solution (35 mL) was added to the reaction mixture and the resulting solution was refluxed at 90°C under Ar for 24 hours (progress of reaction was monitored by TLC). The reaction mixture was cooled and aqueous layer was separated. The organic layer containing the product and spent catalyst was mixed with silica gel (15 g) and concentrated to dryness. The 4-(2-pyridyl)-benzonitrile was isolated by column chromatography (0.850 g, 68%). LCMS: $\text{MH}^+ = 181$; ^1H NMR (CDCl_3) δ 8.85 (d, 1H), 8.7 (dd, 1H), 7.9 (dd, 1H), 7.75 (d, 2H), 7.7 (d, 2H), 7.4 (dd, 1H).

PREPARATIVE EXAMPLES 225-228:

By following essentially same procedure described in Preparative Example 224, only substituting the bromides in column 2 of Table 14,


5 compounds in column 3 of Table 14 were prepared.

Table 14

Prep. Ex.	Column 2	Column 3	Column 4
225			Yield = 70% LCMS: MH ⁺ = 187
226			Yield = 60% LCMS: MH ⁺ = 187
227			Yield = 70% LCMS: MH ⁺ = 186
228			Yield = 70% LCMS: MH ⁺ = 200

PREPARATIVE EXAMPLE 229:

99



5 BH_3 -THF solution (1 M, 24 mL, 5 eq) was added slowly to a stirring solution of 4-(2-pyridyl)-benzonitrile (0.85 g, 4.72 mmol) in anhydrous THF (25 mL) under Ar, and the resulting solution was refluxed for about 12 hr. The solution was cooled to 0°C using ice-water. Methanol (15 mL) was added drop-
 wise to the cold reaction mixture and stirred for 1 h to destroy excess BH_3 . Added HCl – methanol (1M, 10 mL) slowly to the reaction mixture and refluxed for 5 h. Concentrated the solution to dryness and the residue was dissolved in 25
 10 mL water and extracted with ether to remove any un-reacted material. The aqueous solution was neutralized with solid potassium carbonate to pH 10-11. The free amine, thus formed was extracted with ether, dried over potassium carbonate (0.45 g, 50%). LCMS: $\text{MH}^+ = 185$; ^1H NMR (CDCl_3) δ 8.85 (d, 1H), 8.7 (dd, 1H), 7.9 (dd, 1H), 7.75 (d, 2H), 7.7 (d, 2H), 7.4 (dd, 1H), 3.7 (t, 2H), 1.7
 15 (t, 2H).







PREPARATIVE EXAMPLES 230-233:


By following essentially the same procedure set forth in Preparative Example 229, compounds in column 3 of Table 15 were prepared.

Table 15

Prep. Ex.	Column 2	Column 3	Column 4
230			Yield = 60% LCMS: $\text{MH}^+ = 191$

100

231			Yield = 60% LCMS: MH ⁺ = 191
232			Yield = 70% LCMS: MH ⁺ = 190
233			Yield = 70% LCMS: MH ⁺ = 204

PREPARATIVE EXAMPLE 234:**5 Step A:**


A mixture 4-fluorobenzonitrile (3 g, 25 mmol) and imidazolyl sodium (2.48 g, 27.5 mmol) in DMF (50 mL) was stirred at 80°C under Ar for 12 h. Progress of reaction was monitored by TLC. The reaction mixture was concentrated *in vacuo* and the residue was diluted with 50 mL water and stirred. The aqueous mixture was extracted with EtOAc (2 x 50 mL). Combined EtOAc extracts was dried over anhydrous MgSO₄, concentrated, and the 4-(1-imidazolyl)-benzonitrile was isolated by column chromatography (3.6 g, 78%). LCMS: MH⁺ = 170; ¹H NMR (CDCl₃) δ 8.0 (s, 1H), 7.5 (d, 2H), 7.4 (m, 3H), 7.3 (d, 1H)

Step B:

15 4-(1-imidazolyl)-benzonitrile (1g, 5.92 mmol) was dissolved in anhydrous THF (10 mL) and added drop-wise to a stirring solution of LAH –THF (1 M in THF, 18 mL) at room temperature. The reaction mixture was refluxed under Ar


for 2 h and the progress was monitored by TLC. The mixture was cooled to 0°C and quenched by drop-wise addition of a saturated Na₂SO₄ – H₂O solution. The mixture was stirred for 1 h and filtered to remove lithium salts. The filtrate was dried over anhydrous MgSO₄ and concentrated to obtain 4-(1-imidazolyl)-benzylamine (0.8 g, 80%). LCMS: MH⁺ = 174.

PREPARATIVE EXAMPLE 235:



A mixture of 4-(5-oxazolyl)benzoic acid (1.0 g, 5.46 mmol) and Et₃N (552 mg, 5.46 mmol) in 25 mL of THF was cooled to 0 °C and ClCOOi-Bu (745 mg, 5.46 mmol) was added dropwise. After the addition was over, the reaction mixture was stirred for additional 5 min and then aq NH₄OH (0.63 mL of 28% solution, 10.46 mmol) was added. After overnight stirring, the solvent was evaporated, the residue was taken up in water and basified to pH 9. The precipitated solid was filtered, washed with water and dried over P₂O₅ in a vacuum desiccator to provide 500 mg (48%) of the 4-(5-oxazolyl)-benzamide: ¹H NMR (DMSO-d₆) δ 8.50 (s, 1H), 8.20-7.80 (m, 5H).

PREPARATIVE EXAMPLE 236:









A suspension of 4-(5-oxazolyl)benzamide (500 mg, 2.657 mmol) in 10 mL of dry THF was cooled to 0 °C and 10 mL of 1 M BH₃.THF (10.00 mmol) was added. The contents were refluxed overnight and the excess borane was destroyed by dropwise addition of methanol. The solvent was evaporated and the residue was treated with methanolic HCl to decompose the amine-borane

complex. After evaporation of the methanol, the residue was taken in water, basified to pH 10 and the product was extracted in to DCM. The DCM layer was dried (K_2CO_3) and the solvent was removed to provide 150 mg (32%) of 4-(5-oxazolyl)benzylamine: 1H NMR ($CDCl_3$) δ 7.90 (s, 1H), 7.60 (d, 2H), 7.40 (d, 2H), 7.30 (s, 1H), 3.90 (s, 2H).


PREPARATIVE EXAMPLES 237-239:

By essentially the same procedures set forth above, the compounds in Column 2 of Table 16 were reduced using the method indicated in Column 3 of Table 16 to give the amine indicated in Column 4 of Table 16.

Table 16

Prep. Ex.	Column 2	Column 3	Column 4	CMPD
237		BH_3		1H NMR ($CDCl_3$) δ 7.15-6.90 (m, 3H), 3.85 (s, 2H), 1.45 (s, 2H)
238		H_2		1H NMR ($CDCl_3$) δ 8.40 (s, 1H), 7.55 (dd, 1H), 7.10 (d, 1H), 3.85 (s, 2H), 2.50 (s, 3H), 1.70 (bs, 2 H)
239		BH_3		

PREPARATIVE EXAMPLE 240




Prepared by the literature procedure (PCT. Int. Appl, WO 0105783): 1H


NMR ($CDCl_3$) δ 7.35 (d, 1H), 7.24-7.10 (m, 2 H), 7.02 (d, 1H), 3.95 (t, 1H), 3.70 (d, 1H), 3.37 (d, 1H), 2.65 (m, 2H), 2.45 (s, 3H), 1.90 (bs, 2H)

PREPARATIVE EXAMPLE 241:

3-(AMINOMETHYL)PIPERIDINE-1-CARBOXAMIDE



5

A. 3-(*tert*-BUTOXYCARBONYLAMINOMETHYL)PIPERIDINE-1-CARBOXAMIDE

10


3(R/S)-(*tert*-Butoxycarbonylaminomethyl)piperidine (3g, 14.0mmoles) was dissolved in anhydrous dichloromethane (50mL) and trimethylsilylisocyanate (9.68g, 11.4mL, 84.0mmoles) was added. The mixture was stirred under argon at 25°C for 68h. Additional trimethylsilylisocyanate (4.84g, 5.7mL, 42.0mmoles) was added and the mixture was stirred at 25°C for a total of 90h. The mixture was evaporated to dryness and chromatographed on a silica gel column (30x5cm) using 2% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 3-(*tert*-

butoxycarbonylaminomethyl)piperidine-1-carboxamide (3.05g, 85%): FABMS:

m/z 258.1 (MH⁺); HRFABMS: m/z 258.1816 (MH⁺). Calcd. for C₁₂H₂₄O₃N₃: m/z 258.1818; δ_H (CDCl₃) 1.22 (9H, m, CH₃), 1.42 (9H, s, -COOC(CH₃)₃), 1.48 (1H, m, CH₂), 1.67 (2H, m, CH₂), 1.78 (1H, m, CH), 2.80 (1H, m, CH₂), 2.99 (3H, m, CH₂), 3.59 (1H, m, CH₂), 3.69 (1H, m, CH₂), 4.76 (2H, bm, CONH₂) and 4.98ppm (1H, bm, NH); δ_C (CDCl₃) CH₃: 28.5, 28.5, 28.5; CH₂: 24.0, 28.3, 43.2, 45.1,

47.8; CH: 36.5; C: 79.4, 156.3, 158.5.


B. 3-(AMINOMETHYL)PIPERIDINE-1-CARBOXAMIDE




- 5 3-(*tert*-Butoxycarbonylaminoethyl)piperidine-1-carboxamide (150mg, 0.583mmoles) (prepared as described in Preparative Example 241, Step A above) was dissolved in methanol (3mL). 10% conc. sulfuric acid in 1,4-dioxane (7.9mL) was added and the mixture was stirred at 25°C for 1h. The mixture was
- 10 diluted with methanol and BioRad AG1-X8 resin (OH⁻ form) was added until the pH was basic. The resin was filtered off, washed with methanol, evaporated to dryness and chromatographed on a silica gel column (15x2cm) using dichloromethane followed by 15% (10% conc, ammonium hydroxide in methanol)-dichloromethane as the eluant to give the 3-(aminomethyl)piperidine-1-carboxamide (80mg, 87%): FABMS: *m/z* 158.1 (MH⁺); HRFABMS: *m/z*
- 15 158.1294 (MH⁺). Calcd. for C₇H₁₆N₃O: *m/z* 158.1293; δ_H (CDCl₃ + drop CD₃OD) 1.20 (1H, m, CH₂), 1.48 (1H, m, CH₂), 1.60 (1H, m, CH), 1.68 (1H, m, CH₂), 1.83 (1H, m, CH₂), 2.64 (bm, 2H, -CH₂NH₂), 2.82 (1H, m, CH₂), 3.02 (1H, m, CH₂), 2.98 (2H, m, CH₂), 3.70 (1H, m, -CH₂NH₂), 3.78 (1H, m, -CH₂NH₂) and 5.24 ppm (1H, bs, NH); δ_C (CDCl₃ + drop CD₃OD) CH₂: 24.1, 28.6, 44.0, 44.8, 47.9; CH:
- 20 38.3; C: 159.0.

PREPARATIVE EXAMPLE 242:

3-(2-AMINOETHYL)PIPERIDINE-1-CARBOXAMIDE



A. 3-(2-*tert*-BUTOXYCARBONYLAMINOETHYL)PIPERIDINE-1-CARBOXAMIDE



5


3-(2-*tert*-Butoxycarbonylaminoethyl)piperidine (500mg, 2.19mmoles) was dissolved in anhydrous dichloromethane (10mL) and trimethylsilylisocyanate (2.96mL, 21.9mmoles) was added. The mixture was stirred under argon at 25°C for 3.35h. The mixture was diluted with dichloromethane and washed with

10 saturated aqueous sodium bicarbonate. The organic layer was dried (MgSO₄), filtered, evaporated to dryness and chromatographed on a silica gel column (15x5cm) using 5% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 3-(2-*tert*-

15 butoxycarbonylaminoethyl)piperidine-1-carboxamide (417.7mg, 70%): FABMS: m/z 272.0 (MH⁺); HRFABMS: m/z 272.1979 (MH⁺). Calcd. for C₁₃H₂₆O₃: m/z 272.1974; δ_H (CDCl₃) 1.16 (1H, m, CH₂), 1.30-1.60 (5H, m, CH/CH₂), 1.46 (9H, s, -COOC(CH₃)₃), 1.68 (1H, m, CH₂), 1.84 (1H, m, CH₂), 2.54 (1H, dd, CH₂), 2.73 (1H, m, CH₂), 3.08 (1H, m, CH₂), 3.42 (1H, m, CH₂), 4.02 (1H, m, CH₂), 4.10 (1H, m, CH₂), 4.84 (1H, m, NH) and 4.96 ppm (2H, bm, CONH₂); δ_C (CDCl₃)

20 CH₃: 28.5, 28.5, 28.5; CH₂: 25.2, 31.7, 34.9, 37.3, 44.6, 50.3; CH: 32.9; C: 79.5, 156.4, 158.2.

B. 3-(2-AMINOETHYL)PIPERIDINE-1-CARBOXAMIDE




25

3-(2-*tert*-Butoxycarbonylaminoethyl)piperidine-1-carboxamide (392.7mg, 1.45mmoles) (prepared as described in Preparative Example 242, Step A above) was dissolved in methanol (7.5mL) and 10% conc. sulfuric acid in 1,4-dioxane


(19.5mL) was added. The mixture was stirred at 25°C for 1.25h. The mixture was diluted with methanol and BioRad AG1-X8 resin (OH⁻ form) was added until the pH was basic. The resin was filtered off, washed with methanol, evaporated to dryness and chromatographed on a silica gel column (30x2.5cm) using 15% (10% conc, ammonium hydroxide in methanol)-dichloromethane as the eluant to give 3-(2-aminoethyl)piperidine-1-carboxamide (233mg, 94%): FABMS: m/z 172.1 (MH⁺); HRFABMS: m/z 172.1444(MH⁺). Calcd for C₈H₁₈N₃O requires: m/z 172.1450; δ_H (CDCl₃ + 3% CD₃OD) 1.14 (1H, m, CH₂), 1.40 (2H, m, CH₂), 1.49 (1H, m, CH), 1.58 (1H, m, CH₂), 1.69 (1H, m, CH₂), 1.85 (1H, m, CH₂), 2.55 (1H, m, CH₂), 2.67 (5H, m, CH₂/NH₂), 2.76 (1H, bm, CH₂), 2.84 (1H, m, CH₂) and 3.82 ppm (2H, m, CONH₂); δ_C (CDCl₃ + 3% CD₃OD) CH₂: 24.8, 30.9, 36.6, 38.9, 44.9, 50.0; CH: 33.4.

PREPARATIVE EXAMPLE 243:

4-(2-AMINOETHYL)PIPERIDINE-1-CARBOXAMIDE



A. 4-(2-*tert*-BUTOXYCARBONYLAMINOETHYL)PIPERIDINE-1-CARBOXAMIDE




4-(2-*tert*-Butoxycarbonylaminoethyl)piperidine (500mg, 2.19mmoles) was dissolved in anhydrous dichloromethane (10mL) and trimethylsilylisocyanate (2.96mL, 21.9mmoles) was added. The mixture was stirred under argon at 25°C for 3.25h. The mixture was diluted with

dichloromethane and washed with saturated aqueous sodium bicarbonate. The organic layer was dried (MgSO_4), filtered, evaporated to dryness and chromatographed on a silica gel column (15x5cm) using 5% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 4-(2-*tert*-butoxycarbonylaminoethyl)piperidine-1-carboxamide (308.2mg, 52%):

FABMS: m/z 272.0 (MH^+); HRFABMS: m/z 272.1965 (MH^+). Calcd. for $\text{C}_{13}\text{H}_{26}\text{O}_3\text{N}_3$: m/z 272.1974; δ_{H} (CDCl_3) 1.20 (2H, m, CH_2), 1.47 (9H, s, - $\text{COOC}(\text{CH}_3)_3$), 1.45-1.55 (3H, m, CH/CH_2), 1.75 (2H, m, CH_2), 2.82 (2H, m, CH_2), 3.19 (2H, m, CH_2), 3.96 (2H, m, CH_2), 4.64 (2H, m, CH_2) and 4.70 ppm (1H, bm, NH); δ_{C} (CDCl_3) CH_3 : 28.5, 28.5, 28.5; CH_2 : 31.8, 31.8, 36.7, 38.0, 44.5, 44.5; CH: 33.4; C: 79.2, 156.7, 158.1.


A. 3-(2-AMINOETHYL)PIPERIDINE-1-CARBOXAMIDE



4-(2-*tert*-Butoxycarbonylaminoethyl)piperidine-1-carboxamide (283.3mg, 1.04mmoles) (prepared as described in Preparative Example 243, Step A above) was dissolved in methanol (5.4mL) and 10% conc. sulfuric acid in 1,4-dioxane (14.2mL) was added and the mixture was stirred at 25°C for 1.25h. The mixture was diluted with methanol and BioRad AG1-X8 resin (OH^- form) was added until the pH was basic. The resin was filtered off, washed with methanol, evaporated to dryness and chromatographed on a silica gel column (30x2.5cm) using 15% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give the 3-(2-aminoethyl)piperidine-1-carboxamide (170mg, 95%): FABMS: m/z 172.1 (MH^+); HRFABMS: m/z 172.1442. Calcd for $\text{C}_8\text{H}_{18}\text{N}_3\text{O}$ requires: m/z 172.1450; δ_{H} ($\text{CDCl}_3 + 3\% \text{CD}_3\text{OD}$) 1.16 (2H, m, CH_2), 1.43 (2H, m, CH_2), 1.52 (1H, m, CH), 1.70 (2H, m, CH_2), 2.70-2.85 (8H, m, CH_2) and 3.92 ppm (2H, m, CONH_2); δ_{C} ($\text{CDCl}_3 + 3\% \text{CD}_3\text{OD}$) CH_2 : 31.9, 31.9, 39.0, 39.7, 44.4, 44.4; CH: 33.5; C: 158.7.

PREPARATIVE EXAMPLE 244:

3-(AMINOMETHYL)-1-METHYLPYPERIDINE



5


A. 3-(BROMOMETHYL)-1-METHYLPYPERIDINE



10 3-(Hydroxymethyl)-1-methylpyperidine (2g, 15.5mmoles) was dissolved in anhydrous acetonitrile (32mL) and anhydrous pyridine (2.02mL, 24.8mmoles) was added and the solution was cooled to 0°C. Dibromotriphenylphosphorane (8.49g, 20.2mmoles) was added at 0°C and the mixture was allowed to warm up to 25°C and was stirred for 94h. The mixture was evaporated to dryness and the

15 residue was chromatographed on a silica gel column (30x5cm) using gradient elution with dichloromethane, 35% diethyl ether in dichloromethane and 5-10% methanol in dichloromethane as the eluant to give 3-(bromomethyl)-1-methylpyperidine (3.13g, 100%): FABMS: m/z 192.1 (MH^+); δ_H ($CDCl_3$) 1.52 (1H, m, CH_2), 1.99 (2H, m, CH_2), 2.43 (1H, m, CH_2), 2.75 (2H, m, CH_2), 2.82 (1H, m, CH), 2.86/2.88 (3H, s, NCH_3), 3.42/3.49 (2H, dd, $-CH_2Br$) and 3.56 ppm (2H, m, CH_2); δ_C ($CDCl_3$) CH_3 : 44.3; CH_2 : 22.1, 26.6, 35.4, 54.8, 58.2; CH: 34.6.

20

A. 3-(Di-*tert*-BUTOXYCARBONYLAMINOMETHYL)-1-METHYLPYPERIDINE

25

3-(Bromomethyl)-1-methylpyperidine (1.5g, 7.81mmoles) (from Preparative Example 244, Step A above) and di-*tert*-butyliminodiacrylate (1.697g, 7.81mmoles) were dissolved in anhydrous acetonitrile (25mL). Cesium carbonate (5.1g, 15.6mmoles) and lithium iodide (52mg, 0.391mmoles) were

added and the mixture was stirred at 70°C for 20h. The mixture was evaporated to dryness and the residue was partitioned between dichloromethane and saturated aqueous sodium bicarbonate. The organic layer was dried (MgSO₄), filtered and evaporated to dryness. the residue was chromatographed on a silica gel column (30x5cm) using 3% methanol in dichloromethane as the eluant to give 3-(di-*tert*-butoxycarbonylamino)-1-methylpiperidine (1.331g, 52%): FABMS: m/z 329.2 (MH⁺); HRFABMS: m/z 329.2438 (MH⁺). Calcd. for C₁₇H₃₃N₂O₄: m/z 329.2440; δ_H (CDCl₃) 1.10 (1H, m, CH₂), 1.54 (18H, s, -COOC(CH₃)₃), 1.86 (2H, m, CH₂), 2.01 (1H, m, CH₂), 2.19 (1H m, CH), 2.34 (2H, bm, CH₂), 2.59 (3H, -NCH₃), 3.19 (2H, m, CH₂) and 3.52/3.52 ppm (2H, -CH₂N-); δ_C (CDCl₃) CH₃: 28.5, 28.5, 28.5, 28.5, 28.5, 28.5, 47.2; CH₂: 25.4, 28.3, 50.4, 56.8, 60.8; CH: 37.2; C: 83.0, 83.0, 153.5, 153.5.


A. 3-(AMINOMETHYL)-1-METHYLPIPERIDINE



3-(Di-*tert*-butoxycarbonylamino)-1-methylpiperidine (500mg, 1.52mmoles) (from Preparative Example 244, Step B above) was dissolved in methanol (7.5mL) and 10% (v/v) conc. sulfuric acid in 1,4-dioxane (19.75mL) was added. The solution was stirred at 25°C for 0.5h. Methanol (300mL) was added, followed by BioRad AG1-X8 resin (OH⁻ form) until the pH was ~10. The resin was filtered off and washed with methanol (2x200mL). The combined eluates were evaporated to dryness and the residue was chromatographed on a silica gel column (30x2.5cm) using 10% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 3-(aminomethyl)-1-methylpiperidine (69.2mg, 35%): FABMS: m/z 129.1 (MH⁺); HRFABMS: m/z 129.1392 (MH⁺). Calcd. for C₇H₁₇N₂: m/z 129.1392; δ_H (CDCl₃) 0.90 (2H, m, CH₂), 1.65 (2H, m, CH₂), 1.72 (1H, m, CH), 1.79 (1H, m, CH₂), 1.91 (1H, m, CH₂), 2.30 (3H, s, -NCH₃), 2.64 (2H, m, CH₂), 2.82 (1H, m, -CH₂NH₂) and 2.92 ppm (1H, m, -CH₂NH₂); δ_C (CDCl₃) CH₃: 46.7; CH₂: 25.2, 28.0, 46.3, 56.4, 60.3; CH: 39.9.

PREPARATIVE EXAMPLE 245:

4-(AMINOMETHYL)-1-METHYLPYPERIDINE




5

A. 1-METHYLISONIPECOTAMIDE



- 10 Isonipecotamide (10g, 78.0mmoles) was dissolved in distilled water (100mL) and 37% aqueous formaldehyde (7.6mL, equivalent to 2.81g HCHO, 93.6mmoles) was added. Wet 10% Pd-C (8 spoon spatulas) was added under argon and the mixture was hydrogenated at 25°C and 50psi for 43h. The catalyst was filtered off through Celite and the latter was washed with water and
- 15 methanol. The combined filtrates were evaporated to dryness and the residue was chromatographed on a silica gel column (60x5cm) using 8%-10%-20% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 1-methylisonipecotamide (7.15g, 64%): FABMS: m/z 143.1 (MH⁺); HRFABMS: m/z 143.1184 (MH⁺). Calcd. for C₇H₁₅N₂O: m/z 143.1184; δ_H (d₆-DMSO) 1.50/1.57 (4H, m, CH₂), 1.76/1.94 (4H, m, CH₂), 2.10 (3H, s, -NCH₃), 2.72 (1H, m, CH) and 6.68/7.18 ppm (2H, m, CONH₂); δ_C (d₆-DMSO) CH₃: 41.2; CH₂: 28.5, 28.5, 54.9, 54.9; CH: 46.2; C: 176.7.
- 20

B. 4-(AMINOMETHYL)-1-METHYLPYPERIDINE




25


1-Methylisonipecotamide (6.75g, 47.5mmoles) (prepared as described in Preparative Example 245, Step A above) was dissolved in anhydrous THF (350mL) and the resulting mixture was added in portions to a stirred slurry of lithium aluminum hydride (1.8g, 47.5mmoles) in anhydrous THF (100mL) at 0°C under nitrogen. The mixture was stirred at 0°C for 30min and then heated at 66°C for 25h under nitrogen. Distilled water (1.88mL) was added dropwise to the stirred mixture at 0°C, followed by 20% aqueous sodium hydroxide (1.42mL) and then distilled water (6.75mL) and the mixture was stirred for 15min. The mixture was filtered and the solids were washed with THF and dichloromethane. The combined filtrates were evaporated to dryness and chromatographed on a silica gel column (30x5cm) using 15%-20% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 4-(aminomethyl)-1-methylpiperidine (0.678g, 11%): FABMS: m/z 129.1 (MH⁺); HRFABMS: m/z 129.1389 (MH⁺). Calcd. for C₇H₁₇N₂: m/z 129.1392; δ_H (d₆-DMSO): 2.08ppm (3H, s, -NCH₃); δ_C (d₆-DMSO): CH₃: under DMSO peaks; CH₂: 29.6, 29.6, 46.7, 55.2, 55.2; CH: 46.2.

PREPARATIVE EXAMPLE 246:

3-(AMINOMETHYL)BENZONITRILE



A. 3-(Di-*tert*-BUTOXYCARBONYLAMINO)BENZONITRILE



3-(Bromomethyl)benzonitrile (5g, 25.5mmoles) and di-*tert*-butyliminodicarboxylate (5.54g, 25.5mmoles) were dissolved in anhydrous THF

(50mL) and cesium carbonate (16.62g, 25.5mmoles) and lithium iodide (170.5mg, 1.275mmoles) were added. The mixture was stirred at 70°C for 22h and the reaction was worked up as described in Preparative Example 89, Step B above. The residue was chromatographed on a silica gel column (60x5cm) using 5% ethyl acetate in hexane as the eluant to give 3-(di-*tert*-butoxycarbonylamino)benzonitrile (7.39g, 87%): FABMS: m/z 333.2 (MH^+); HRFABMS: m/z 333.1815 (MH^+); Calcd. for $C_{18}H_{25}N_2O_4$: m/z 333.1814; δ_H ($CDCl_3$) 1.52 (18H, s, $-COOC(CH_3)_3$), 4.84 (2H, s, CH_2), 7.48 (1H, m, Ar-H), 7.60 (2H, m, Ar-H) and 7.65 ppm (1H, m, Ar-H); δ_C ($CDCl_3$) CH_3 : 28.1, 28.1, 28.1, 28.1, 28.1, 28.1; CH_2 : 48.4; CH: 129.2, 131.0, 131.0, 131.9; C: 83.2, 83.2, 112.5, 118.8, 140.1, 152.5, 152.5.


B. 3-(AMINOMETHYL)BENZONITRILE




3-(Di-*tert*-butoxycarbonylamino)benzonitrile (2g, 6.0mmoles) (prepared as described in Preparative Example 246, Step A above) was dissolved in methanol (30mL) and 10%(v/v) (10% conc. sulfuric acid in 1,4-dioxane) (79mL) was added. The solution was stirred at 25°C for 0.25h and worked up as described in Preparative Example 89, Step C above). The residue was chromatographed on a silica gel column (15x5cm) using 3% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give the title compound (651.4mg, 82%): FABMS: m/z 133.1 (MH^+); HRFABMS: m/z 133.0762 (MH^+). Calcd. for $C_8H_9N_2$: m/z 133.0766 ; δ_H ($CDCl_3$) 2.57 (2H, s, $-CH_2NH_2$), 3.92 (2H, s, $-CH_2NH_2$), 7.46 (1H, m, Ar-H), 7.57 (2H, m, Ar-H) and 7.64 ppm (1H, m, Ar-H); δ_C ($CDCl_3$) CH_2 : 45.2; CH: 129.4, 130.7, 130.7, 131.8; C: 112.4, 118.8, 143.8.

PREPARATIVE EXAMPLE 247:

4-(AMINOMETHYL)BENZONITRILE




5

A. 3-(Di-*tert*-BUTOXYCARBONYLAMINOMETHYL)BENZONITRILE

4-(Bromomethyl)benzonitrile (5g, 25.5mmoles) and di-*tert*-butyliminodicarboxylate (5.54g, 25.5mmoles) were dissolved in anhydrous THF (50mL) and cesium carbonate (16.62g, 25.5mmoles) and lithium iodide (170.5mg, 1.275mmoles) were added. The mixture was stirred at 70°C for 23h and the reaction was worked up as described in Preparative Example 244, Step B above. The residue was chromatographed on a silica gel column (50x5cm) using 5% ethyl acetate in hexane as the eluant to give 4-(di-*tert*-


butoxycarbonylamino)methylbenzonitrile (7.07g, 83%); FABMS: m/z 333.2 (MH^+); HRFABMS: m/z 333.1816 (MH^+). Calcd. for $C_{18}H_{25}N_2O_4$: m/z 333.1814; δ_H ($CDCl_3$) 1.45 (18H, s, $-COOC(CH_3)_3$), 4.81 (2H, s, CH_2), 7.37 (2H, d, Ar-H) and 7.62 ppm (2H, d, Ar-H); δ_C ($CDCl_3$) CH_3 : 28.1, 28.1, 28.1, 28.1, 28.1, 28.1; CH_2 : 49.2; CH : 127.8, 127.8, 132.3, 132.3; C: 83.2, 83.2, 111.1, 118.9, 144.1, 152.4, 152.4.

B. 4-(AMINOMETHYL)BENZONITRILE



4-(Di-*tert*-butoxycarbonylaminomethyl)benzonitrile (2g, 6.0mmoles) (prepared as described in Preparative Example 247, Step A above) was dissolved in TFA (4mL) and the solution was stirred at 25°C for 0.25h. The reaction mixture was diluted with dichloromethane and extracted with 1N sodium hydroxide. The organic layer was dried (MgSO₄), filtered and evaporated to dryness. The residue was chromatographed on a silica gel column (15x5cm) using 3% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 4-(aminomethyl)benzonitrile (108mg, 68%): FABMS: m/z 133.1 (MH⁺); HRFABMS: m/z 133.0764 (MH⁺). Calcd. for C₈H₉N₂: m/z 133.0766; δ_H (CDCl₃) 2.04 (2H, s, -CH₂NH₂), 3.89 (2H, s, -CH₂NH₂), 7.40 (2H, d, Ar-H) and 7.59 ppm (2H, d, Ar-H); δ_C (CDCl₃) CH₂: 45.7; CH: 127.8, 127.8, 132.4, 132.4; C: 110.6, 118.9, 148.0.

PREPARATIVE EXAMPLE 248









To a solution of (1S,2S)-2-benzyloxycyclopentyl amine (1.5 g, 7.84 mmol) in MeOH (50 mL) at rt was added 10 % Pd/C (50% wet, 1.0 g) followed by dropwise addition of conc. HCl (0.7 mL). The mixture was stirred under a balloon of H₂ for 14 h and the catalyst was filtered off thru a pad of Celite. The pad of Celite was washed with MeOH (2 x 10 mL) and the resulting filtrate was concentrated under reduced pressure to afford 0.97 g (90%) of a yellow semisolid; M+H (free base) = 102

PREPARATIVE EXAMPLES 249-251

In an analogous fashion to Preparative Example 248, the benzyl protected cycloalkyl amines (Column 2) were converted to the desired aminocycloalkanol hydrochloride derivatives (Column 3) as listed in Table 17.

115


TABLE 17

Ex.	Column 2 (Amine)	Column 3 (Cleavage method)	CMPD M+H
249			M+H = 102 (free base)
250			M+H = 116 (free base)
251			M+H = 116 (free base)

5 PREPARATIVE EXAMPLE 252



To a solution of ester (prepared according to *J. Org. Chem.* (1999), 64, 330) (0.5 g, 2.43 mmol) in THF (8 mL) at 0 °C. was added LiAlH₄ (0.37 g, 9.74 mmol) in one portion. The resulting mixture was heated at reflux for 12h and was cooled to 0 °C. The mixture was treated sequentially with H₂O (1 mL), 1 M NaOH (1 mL), and H₂O (3 mL). CH₂Cl₂ (10 mL) was added to the mixture which was stirred vigorously for 30 min. The mixture was filtered thru a pad of Celite which was washed generously with CH₂Cl₂ (3 x 5 mL). The resulting filtrate was concentrated under reduced pressure to afford 0.41 g (85%) of a yellow/orange solid. M+H = 142.

PREPARATIVE EXAMPLE 253**5** STEP A:

To a solution of *L*-proline methyl ester hydrochloride (0.50 g, 3.0 mmol) in CH_2Cl_2 (15 mL) at 0 °C was added Et_3N (1.1 mL, 7.55 mmol) followed by TFAA (0.56 mL, 3.92 mmol). The mixture was stirred for 12 h at rt and 1N HCl (25 mL) was added. The layers were separated and the organic layer was washed sequentially with sat. aq. NaHCO_3 (1 x 25 mL), and brine (1 x 25 mL). The organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure to afford 0.72 g (100%) of a yellow oil. $M+H = 226$. The crude material was taken onto Step B without further purification.

15 STEP B:

To a solution of the compound prepared in Preparative Example 253, Step A (0.68 g, 3.0 mmol) in THF (20 mL) at 0 °C was added MeMgI (5.1 mL, 3.0M in Et_2O) dropwise over 10 min. The resulting solution was stirred for 16 h at rt whereupon the mixture was quenched by addition of sat. aq. NH_4Cl . The mixture was concentrated to dryness and the resultant residue was stirred with EtOAc (100 mL) for 45 min and filtered. The filtrate was concentrated under reduced pressure to afford 0.68g (100%) of a yellow/orange oil. $M+H = 226$. The crude material was taken onto Step C without further purification.


25 STEP C:

To a solution of the compound prepared in Preparative Example 253, Step B (0.68 g, 3.0 mmol) in MeOH (5 mL) was added a solution of KOH (0.68 g, 12.1 mmol) in MeOH (5 mL). The mixture was stirred at reflux for 12h and rt for 72h whereupon the mixture was concentrated to dryness. The crude residue

117


was suspended in EtOAc (50 mL) and was stirred vigorously for 30 min and was filtered. This procedure was repeated 2X more and the resultant filtrate was concentrated under reduced pressure to afford 128 mg (33%) of a maroon/orange oil. M+H = 130. This material was used without purification in the subsequent coupling step.

PREPARATIVE EXAMPLE 254:




The aldehyde was prepared according to the procedure of Gupton (*J. Heterocyclic Chem.* (1991), 28, 1281).

PREPARATIVE EXAMPLE 255



Using the aldehyde from Preparative Example 254, the procedure of Gupton (*J. Heterocyclic Chem.* (1991), 28, 1281) was employed to prepare the title aldehyde.


PREPARATIVE EXAMPLE 256



The title aldehyde was prepared according to the procedure of Ragan et. al *Synlett* (2000), 8, 1172-1174.


PREPARATIVE EXAMPLE 257

118



The reaction of known cyclopentyl guanidine hydrochloride (*Org. Lett.* (2003), 5, 1369-1372) under the conditions of Ragan (*Synlett* (2000), 8, 1172-1174) afforded the title aldehyde.


PREPARATIVE EXAMPLE 258



The title compound was prepared according to known literature
10 *Monatshefte fur Chemie* (1973), 104, 1372-1382.

EXAMPLES

EXAMPLE 1:












A solution of the product from Preparative Example 127 (0.27 g, 0.875
15 mmol), 4-aminomethylpyridine (0.12 g, 1.3 eq.), and K_2CO_3 (0.24 g, 2 eq.) in CH_3CN (5 mL) was stirred at room temperature 48 hours. The reaction mixture was diluted with H_2O and extracted with CH_2Cl_2 . The combined organics were dried over Na_2SO_4 , filtered and concentrated. The crude product was purified by
flash chromatography using a 4% MeOH in CH_2Cl_2 solution as eluent (0.28 g,
20 93% yield). LCMS: $MH^+ = 380$; mp = $>205^\circ C$ (dec).

EXAMPLES 2-210:













By following essentially the same procedure set forth in Example 1 only substituting the chlorides shown in Column 2 of Table 18 and the amines shown in Column 3 of Table 18, the compounds in Column 4 of Table 18 were prepared:













5













TABLE 18













Ex.	Column 2	Column 3	Column 4	Data
2				LCMS: MH ⁺ = 380; mp=175- 176°C
3				LCMS: MH ⁺ = 398; mp= 156- 157°C
4				LCMS: MH ⁺ = 398; mp= 45-49°C

120
















5				LCMS: MH ⁺ = 354; mp= 43-46°C
6				LCMS: MH ⁺ = 354; mp= 149- 150°C
7				LCMS: MH ⁺ = 414; mp= 86-92°C
8				LCMS: MH ⁺ = 414; mp= 185- 186°C













9				LCMS: MH ⁺ = 448; mp= 167- 168°C
10				LCMS: MH ⁺ = 346; mp= 57-58°C
11				LCMS: MH ⁺ = 347; mp=122. 9-125.3 °C
12				LCMS: MH ⁺ = 360; mp= 127- 128°C
















13				LCMS: MH ⁺ = 342; mp= 133- 135°C
14				LCMS: MH ⁺ = 344; mp= 152- 155°C
15				LCMS: MH ⁺ = 362; mp= 164- 167°C
16				LCMS: MH ⁺ = 327; mp= 146- 155°C
















17				LCMS: MH ⁺ = 332; mp= 71-82°C
17. 1				MS: MH ⁺ = 332.
18				LCMS: MH ⁺ = 346; mp= 58-65°C
19				LCMS: MH ⁺ = 414; mp= 211- 213°C













20				LCMS: MH ⁺ = 414; mp = 194- 197°C
21				MS: MH ⁺ = 414 m.p. 211 - 216°C
22				LCMS: MH ⁺ = 544; mp = 104- 107°C
23				Yield = 83% LCMS: MH ⁺ = 410.













24				Yield = 84% LCMS: MH ⁺ = 410.
25				Yield = 96% LCMS: MH ⁺ = 440.
26				Yield = 99% LCMS: MH ⁺ = 440.
27				Yield = 89% LCMS: MH ⁺ = 448.
28				Yield = 78% LCMS: MH ⁺ = 448.













30				Yield = 96% LCMS: MH ⁺ = 483.
31				Yield = 35% LCMS: MH ⁺ = 483.
32				Yield = 77% LCMS: MH ⁺ = 515.
33				Yield = 100% m.p. 179°C LCMS: MH ⁺ = 388













34				Yield = 99% m.p. 186°C LCMS: MH ⁺ = 456
35				Yield = 98% m.p. 181°C LCMS: MH ⁺ = 401
36				Yield = 63% m.p. 192°C LCMS: MH ⁺ = 480
37				Yield = 75% m.p. 126 – 127°C LCMS: MH ⁺ = 400
38				Yield = 94% m.p. 132 – 133°C LCMS: MH ⁺ = 400

39				Yield = 95% m.p. 121 - 122°C LCMS: MH ⁺ = 400
40				Yield = 98% LCMS: MH ⁺ = 460
41				Yield = 87% m.p. 170 - 171°C LCMS: MH ⁺ = 464
42				Yield = 84% m.p. 216 - 217°C LCMS: MH ⁺ = 464
43				Yield = 96% m.p. 214°C LCMS: MH ⁺ = 464













44				Yield = 95% m.p. 158°C LCMS: MH ⁺ = 522
45				Yield=90 % LCMS: MH ⁺ = 278
46				Yield=10 0% ; LCMS: MH ⁺ =394
47				LCMS: MH ⁺ = 473 m.p. 84 - 87°C









48				MS: MH ⁺ = 396 m.p. 91.5 - 93.3°C
49				MS: MH ⁺ = 396 m.p. 196 - 199°C
50				MS: MH ⁺ = 430 m.p. 242 - 244°C
51				MS: MH ⁺ = 430 m.p. 218°C

52				MS: MH ⁺ = 430 m.p. 230 - 233°C
54				MS: MH ⁺ = 405 m.p. 185 - 188°C
55				MS: MH ⁺ = 370 m.p. 229 - 232°C
56				MS: MH ⁺ = 370 m.p. 85 - 90°C













57				MS: MH ⁺ = 386 m.p. 227 - 230°C
58				MS: MH ⁺ = 372 m.p. 212 - 215°C
59				MS: MH ⁺ = 318 m.p. 169 - 171°C
60				MS: MH ⁺ = 332 m.p. 170 - 173°C

133










61				MS: MH ⁺ = 346 m.p. 156 – 159°C
62				MS: MH ⁺ = 360 m.p. 114 – 116°C
63				MS: MH ⁺ = 348 m.p. 197 – 200°C
64				1. mp = 230-232 2. M+H = 396













65		$\text{H}_2\text{N}-\text{CH}_2-\text{C}_5\text{H}_4\text{N}-\text{NH}_2$ 2 HCl		1. mp = 205-207 2. M+H = 402
66		$\text{H}_2\text{N}-\text{CH}_2-\text{C}_5\text{H}_4\text{N}-\text{NH}_2$ 2 HCl		1. mp = 220-223 2. M+H = 414
67		$\text{H}_2\text{N}-\text{CH}_2-\text{C}_5\text{H}_4\text{N}-\text{NH}_2$ 2 HCl		1. mp = 191-193 2. M+H = 431
68		$\text{HCl} \cdot \text{H}_2\text{N}-\text{CH}_2-\text{C}_5\text{H}_3\text{N}_2-\text{NH}_2$		1. mp = 235-237 2. M+H = 397













135













69		$\text{HCl} \cdot \text{H}_2\text{N}$ 		1. mp = >250 2. M+H = 403
70		$\text{HCl} \cdot \text{H}_2\text{N}$ 		1. mp = 230-232 2. M+H = 415
71		$2\text{HCl} \cdot \text{H}_2\text{N}$ 		1. mp = 235-238 2. M+H = 431
72		H_2N 2HCl 		1. mp = 186-188 2. M+H = 410

136













73				1. mp = 136-138 2. M+H = 424
74				1. mp = 192-195 2. M+H = 450
75				1. mp = 88-90 2. M+H = 454

76				1. mp = 230-232 2. M+H = 467
77				1. mp = 131-133 2. M+H = 479
78				1. mp = 85-88 2. M+H = 376
79				1. mp = 131-133 2. M+H = 388













80		 2 HCl		1. mp = 206-208 2. M+H = 408
81		 HCl		1. mp = 108-110 2. M+H = 502
82		 2 HCl		1. mp = 83-85 2. M+H = 402
83		 2 HCl		1. mp = 220 2. M+H = 414













84		 · 2 HCl		1. mp = 154-156 2. M+H = 426
85		 · 2 HCl		1. mp = 152-153 2. M+H = 438
86		 · 2 HCl		1. mp = 159-161 2. M+H = 420
87		 · 2 HCl		1. mp = >220 2. M+H = 455

140













88				1. mp = 223-225 2. M+H = 425
89				1. mp = 199-201 2. M+H = 419
90				1. mp = 184-186 2. M+H = 426
91				1. mp = 196-198 2. M+H = 420

141













92				1. mp = 156-159 2. M+H = 440
93				1. mp = 173-176 2. M+H = 434
94				1. mp = 173-175 2. M+H = 452
95				1. mp = 174-176 2. M+H = 469
















96		 · 2 HCl		1. mp = 230-234 2. M+H = 434
97				1. mp = 191-193 2. M+H = 441
98				1. mp = 202-205 2. M+H = 434
99				1. mp = 209-212 2. M+H = 453

143
















100				1. mp = 219-221 2. M+H = 469
101				1. mp = 64-66 2. M+H = 403
102				1. mp = 168-170 2. M+H = 420
103				1. mp = 213-216 2. M+H = 411













144

104		 · 2 HCl		1. mp = 98-100 2. M+H = 561
105		 · 3 HCl		1. mp 70-72 2. M+H = 608
106		 HCl · H ₂ N		1. mp 168-170 2. M+H = 538
107		 · 2 HCl		1. mp 189-191 2. M+H = 592
















108				LCMS: MH ⁺ = 458;
109				Yield = 89 LCMS: MH ⁺ = 418 m. p. = 131-132 °C
110				Yield=95% LCMS: MH ⁺ = 347
111				Yield=91% (3H); LCMS: MH ⁺ = 484
112				Yield=87% LCMS: MH ⁺ = 427

146













113				Yield=80% LCMS: MH ⁺ = 427
114				Yield=91% LCMS: MH ⁺ = 378
115				Yield=92% (3H); LCMS: MH ⁺ =520
116				Yield=98% LCMS: MH ⁺ =536
117				Yield=82% LCMS: MH ⁺ =410

118				Yield=95% LCMS: MH ⁺ = 347
121				Yield = 65% LCMS: MH ⁺ = 481.02
126				Yield=71% MH ⁺ = 486
127				Yield=71% MH ⁺ = 495.1













148

128				Yield=55% MH ⁺ = 463
129				Yield = 77% LCMS: MH ⁺ = 455
130				¹ H NMR (Yield = 75% LCMS: MH ⁺ = 379
131				Yield = 75% LCMS: MH ⁺ = 407
132				Yield = 75% LCMS: MH ⁺ = 421













149

133				Yield = 70% LCMS: MH ⁺ = 421
134				Yield = 78% LCMS: MH ⁺ = 475
135				Yield = 75% LCMS: MH ⁺ = 476
136				Yield = 65% LCMS: MH ⁺ = 455













150

137				Yield = 55% LCMS: MH ⁺ = 473)
138				Yield = 60% LCMS: MH ⁺ = 439
139				Yield = 65% LCMS: MH ⁺ = 441
140				Yield = 80% LCMS: MH ⁺ = 432













151

141				Yield = 60% LCMS: MH ⁺ = 429
142				LCMS: MH ⁺ =330 ; mp=109- 111°C
143				LCMS: MH ⁺ =346 ; mp=186- 188°C
144				LCMS: MH ⁺ =384 ; mp=148- 150°C










152

145				LCMS: MH ⁺ =400 ; mp=186-188°C
146				LCMS: M2H ⁺ =390; mp=192-194°C
147				LCMS: M ⁺ =404; mp=220-222°C
148				LCMS: MH ⁺ =369 ; mp>230°C













153

149				LCMS: MH ⁺ =364 ; mp=186-188°C
150				LCMS: MH ⁺ =312 ; mp=138-140°C
151				LCMS: M ⁺ =380; mp=172-174°C
152				LCMS: MH ⁺ =352 ; mp=201-203°C
















154

153				LCMS: MH ⁺ =348 ; mp=166-168°C
154				LCMS: M2H ⁺ =53 1; mp=78-80°C
155				LCMS: M2H ⁺ =47 4; mp=161-163°C
















155

156				LCMS: $M^+ = 444$; mp=48-51°C
157				$MH^+ = 542.1$
158				$MH^+ = 520.1$
159				$MH^+ = 542.1$













156













160				MH ⁺ = 480.1
161				MH ⁺ = 506.1
162				MH ⁺ = 480.1
163				MH ⁺ = 494.1
164				MH ⁺ = 466.1













157












165				MH ⁺ = 494.1
166				MH ⁺ = 508.1
167				MH ⁺ = 520.1
168				MH ⁺ = 528.1
169				MH ⁺ = 520.1













158













170				MH ⁺ = 528.1
171				LCMS: MH ⁺ = 474;
172				LCMS: MH ⁺ = 437;
173				LCMS: MH ⁺ = 472;













174				LCMS: $MH^+ = 428.1$
175				LCMS: $MH^+ = 426.2$
176				LCMS: $MH^+ = 442.0$
177				LCMS: $MH^+ = 452.0$













178				Yield = 90 MH ⁺ = 436 m. pt. = 89.1 °C
179				MH ⁺ = 424 m. pt. = 188.2 °C
180				MH ⁺ = 448 m. pt. = 211.3 °C
181				Yield = quant. MH ⁺ = 464
















182				MH ⁺ = 382 m. pt. = 185.8 °C
183				MH ⁺ = 387 m. pt. = 181 – 182 °C
184				MH ⁺ = 453
185				MH ⁺ = 401 m. pt. = 178.3 °C













186				MH ⁺ = 402
187				Yield = 91 MH ⁺ = 386 m. pt. = 148.3 °C
188				Yield = 65 MH ⁺ = 402 m. pt. = 174.5 °C
189				MH ⁺ = 379 m. pt. = 82 – 83 °C







190				MH ⁺ = 379 m. pt. = 50.7 °C
191				Yield = 89 MH ⁺ = 469 m. pt. = 186.7°C
192				Yield = 93 MH ⁺ = 410 m. pt. = 86.7°C
193				Yield = 76 MH ⁺ = 333 m. pt. = 120.3°C

194				Yield = 86 MH ⁺ = 353 m. pt. = 188.9 °C
195				Yield=11% LCMS:374 MH ⁺ = 390
196				Yield=88% LCMS:374 MH ⁺ = 346
197				Yield=88% LCMS:374 MH ⁺ = 346

198				Yield = MH ⁺ = 400 m. pt. = 111.5 – 112.2 °C
199				MH ⁺ = 416
200				MH ⁺ = 415
201				MH ⁺ = 398 m.p. = 156.5 °C

202				MH ⁺ = 414 m.p. = 89.5 °C
203				MH ⁺ = 413
204				Yield = 86 MH ⁺ = 521 m.p. = 79.9 °C
204 .10				
204 .11				Yield = 87 MH ⁺ = 521 m.p. = 128.6 °C

205				Yield = 99 MH ⁺ = 537 m.p. = 83.5 °C
206				Yield = 94 MH ⁺ = 598 m.p. = 110.8 °C
207				Yield = quant. MH ⁺ = 545
208				Yield = 96 MH ⁺ = 468 m.p. = 69.2 °C

209				MH ⁺ = 498 m.p. = 226.5 °C
210				MH ⁺ = 564 m.p. = 174.2 °C

Additional data for select examples given below.

Example 23: ¹H NMR (CD₃OD) δ 8.63 (d, J = 5.7 Hz, 2H), 8.18 (s, 1H), 7.81 (dd, J = 8.1 Hz, 2.1 Hz, 1H), 7.58 (d, J = 6.0 Hz, 2H), 7.48 (m, 1H), 7.15-7.10 (m, 2H), 6.50 (s, 1H), 4.86 (s, 2H), 3.70 (s, 3H)

Example 24: ¹H NMR (CDCl₃) δ 8.82 (s, 1H), 8.73 (d, J = 4.2 Hz, 1H), 8.11 (s, 1H), 8.06 (dd, J = 7.8 Hz, 1.8 Hz, 1H), 7.91 (d, J = 8.1 Hz, 1H), 7.53-7.47 (m, 2H), 7.20 (m, 1H), 7.08 (d, J = 8.1 Hz, 1H), 6.75 (s, 1H), 4.81 (d, J = 4.5 Hz, 2H), 3.86 (s, 3H)

Example 25: ¹H NMR (CDCl₃) δ 8.75 (d, J = 5.7 Hz, 2H), 8.12 (s, 1H), 7.81 (d, J = 2.1 Hz, 1H), 7.53 (dd, J = 8.4, 2.1 Hz, 1H), 7.45 (d, J = 6.0 Hz, 2H), 6.96 (t, J = 6.0 Hz, 2H), 6.33 (s, 1H), 4.85 (d, J = 6.0 Hz, 2H), 4.09 (s, 3H), 4.03 (s, 3H)

Example 26: ¹H NMR (CDCl₃) δ 8.82 (s, 1H), 8.72 (s, 1H), 8.09 (m, 1H), 7.87-7.83 (m, 2H), 7.60 (m, 1H), 7.45 (m, 1H), 7.03 (d, J = 8.4 Hz, 1H), 6.87 (s, 1H), 6.43 (s, 1H), 4.83 (d, J = 4.5 Hz, 2H), 4.11 (s, 3H), 4.04 (s, 3H)

Example 27: ¹H NMR (CDCl₃) δ 8.75 (d, J = 4.5 Hz, 2H), 8.19 (s, 1H), 7.63 (d, J = 7.8 Hz, 2H), 7.44-7.40 (m, 3H), 7.07 (m, 1H), 6.26 (s, 1H), 4.83 (d, J = 5.1 Hz, 2H)

Example 28: ^1H NMR (CDCl_3) δ 8.86 (s, 1H), 8.74 (m, 1H), 8.17 (s, 1H), 7.97 (m, 1H), 7.66-7.63 (m, 2H), 7.62 (m, 1H), 7.41 (m, 1H), 7.07 (m, 1H), 6.35 (s, 1H), 4.87 (d, $J = 6.0$ Hz, 2H)

Example 30: ^1H NMR (CDCl_3) δ 8.16 (s, 1H), 7.66-7.62 (m, 2H), 7.41 (m, 1H),
5 7.33-7.22 (m, 3H), 6.96 (t, $J = 6.0$ Hz, 1H), 6.33 (s, 1H), 4.73 (d, $J = 6.0$ Hz, 2H)

Example 31: ^1H NMR (CDCl_3) δ 8.13 (s, 1H), 7.66 (d, $J = 7.8$ Hz, 2H), 7.45-7.40 (m, 2H), 7.10-7.04 (m, 2H), 6.93 (t, $J = 6.6$ Hz, 1H), 6.60 (s, 1H), 4.84 (d, $J = 6.6$ Hz, 2H)

Example 32: ^1H NMR (CDCl_3) δ 8.16 (s, 1H), 7.66-7.62 (m, 2H), 7.57-7.55 (m, 2H),
10 7.41 (t, $J = 7.8$ Hz, 1H), 7.31 (dd, $J = 7.8, 1.8$ Hz, 1H), 6.99 (t, $J = 6.0$ Hz, 1H), 6.32 (s, 1H), 4.73 (d, $J = 6.0$ Hz, 2H)

Example 40: ^1H NMR (CDCl_3) δ 8.01 (s, 1H), 7.31 – 7.24 (d, $J = 8.2$ Hz, 1H),
6.72 – 6.64 (br t, $J = 5.4$ Hz, 1H), 6.62 – 6.52 (m, 2H), 6.05 – 6.01 (s, 1H), 5.56 –
15 4.64 (d, $J = 6.0$ Hz, 2H), 4.03 – 3.93 (s, 3H), 3.94 – 3.86 (s, 3H), 2.79 – 2.70 (d, $J = 8.1$ Hz, 2H), 2.02 – 1.66 (m, 6H), 1.43 – 1.22 (m, 3H), 1.20 – 1.02 (m, 2H)

Example 45: ^1H NMR (CDCl_3) δ 8.73(d, 2H), 8.54(s, 1H), 7.41(d, 2H), 7.02(br, 1H),
5.90(s, 1H), 4.80(s, 2H), 4.48(q, 2H), 2.75(s, 2H), 1.50(t, 2H), 1.06(s, 9H);

Example 46: ^1H NMR (CDCl_3) δ 8.79(s, 1H), 8.72(d, 1H), 8.14(s, 1H), 7.84(d, 1H),
7.54-7.33(m, 4H), 6.97(t, 1H), 6.18(s, 1H), 4.79(d, 2H), 2.47(s, 3H)

Example 108: ^1H NMR (CDCl_3) δ 8.79 (s, 1H), 8.72 (d, $J = 3.0$ Hz, 1H), 8.16 (s, 1H),
20 7.84 (d, $J = 7.8$ Hz, 1H), 7.74 (d, $J = 7.5$ Hz, 2H), 7.55-7.35 (m, 3H), 6.92 (t, $J = 6.3$ Hz, 1H), 6.42 (s, 1H), 4.81 (d, $J = 6.3$ Hz, 2H)

Example 110: ^1H NMR (CDCl_3) δ 8.18(t, 1H), 8.03(s, 1H), 7.44(m, 1H), 7.30(t, 1H),
7.17(q, 1H), 6.66(s, 1H), 6.56(br, 1H), 4.28(d, 2H), 2.38(s, 1H)

Example 111: ^1H NMR (CDCl_3) δ 8.72(br, 1H), 8.59(d, 1H), 8.11(t, 1H), 8.06(s, 1H),
25 7.73(d, 1H), 7.44(d, 1H), 7.42-7.21(m, 3H), 7.07(q, 1H), 6.39(d, 1H), 5.21(q, 1H), 4.16(q, 2H), 3.08(d, 2H), 1.22(t, 3H)

Example 112: ^1H NMR (CDCl_3) δ 8.22(t, 1H), 8.15(s, 1H), 7.51-7.33(m, 7H),
7.21(q, 1H), 6.82(d, 1H), 6.51(s, 1H), 4.68(q, 1H), 2.18(m, 2H), 1.17(t, 3H)

Example 113: ^1H NMR (CDCl_3) δ 8.22(t, 1H), 8.14(s, 1H), 7.51-7.33(m, 7H),
30 7.21(q, 1H), 6.82(d, 1H), 6.51(s, 1H), 4.68(q, 1H), 2.18(m, 2H), 1.17(t, 3H)

Example 114: ^1H NMR (CDCl_3) δ 8.81(s, 1H), 8.75(d, 1H), 8.21(s, 1H), 7.84(d, 1H), 7.47(q, 1H), 6.96(s, 1H), 6.94(t, 1H), 4.85(d, 2H), 4.60(q, 2H), 1.58(t, 3H)

Example 115: ^1H NMR (CDCl_3) δ 8.77(s, 1H), 8.72(d, 1H), 8.14(s, 1H), 7.83(d, 1H), 7.65(d, 1H), 7.44(q, 1H), 7.80(t, 1H), 7.6(d, 1H), 6.18(s, 1H), 4.75(d, 2H),

5 3.91(s, 3H), 3.81(s, 3H)

Example 116: ^1H NMR (CDCl_3) δ 8.67(s, 1H), 8.55(d, 1H), 8.50(s, 1H), 7.92(d, 1H), 7.90(d, 1H), 7.78(t, 1H), 7.10(d, 1H), 6.97(s, 1H), 5.11(s, 2H), 3.77(s, 6H)

Example 117: ^1H NMR (CDCl_3) δ 8.38(s, 1H), 8.30(d, 1H), 8.17(s, 1H), 7.52-7.37(m, 6H), 6.97(t, 1H), 6.13(s, 1H), 4.77(d, 2H), 2.50(s, 3H)

10 **Example 118:** ^1H NMR (CDCl_3) δ 8.18(t, 1H), 8.03(s, 1H), 7.44(m, 1H), 7.30(t, 1H), 7.17(q, 1H), 6.66(s, 1H), 6.56(br, 1H), 4.28(d, 2H), 2.38(s, 1H);

Example 121: ^1H NMR (CDCl_3) δ 8.6 (s, 1H), 8.15 (dt, 1H), 8.1 (s, 1H), 8.0 (d, 2H), 7.5 (d, 2H), 7.4 (dd, 1H), 7.2 (d, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.6 (s, 1H), 4.75 (d, 2H).

15 **Example 126:** ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.5 (d, 1H), 7.42 – 7.35 (m, 2H), 7.3 – 7.2 (m, 2H), 7.15 (dd, 1H), 7.1 (dd, 1H), 7.0 (t, 1H), 6.6 (s, 1H), 4.8 (d, 2H).

Example 127: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 (dd, 1H), 7.3- 7.25 (m, 3H), 7.1 (dd, 1H), 6.9 - 6.85 (m, 2H), 6.7 (t, 1H), 6.6 (s, 1H), 4.6 (d, 2H), 3.2 (m, 4H), 2.6 (m, 4H), 2.3 (s, 3H)

20 **Example 128:** ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.1 (s, 1H), 8.0 (d, 2H), 7.5 (d, 2H), 7.4 (m, 2H), 7.25 (d, 1H), 7.2 (s, 1H), 7.15 (dd, 1H), 7.0 (s, 1H), 6.8 (t, 1H), 6.6 (s, 1H), 4.75 (d, 2H).

Example 129: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.05 (s, 1H), 8.0 (d, 2H), 7.5 (d, 2H), 7.4 (m, 1H), 7.3 (dd, 1H), 7.15 (dd, 1H), 6.9 (t, 1H), 6.5 (s, 1H), 4.75 (d, 2H), 3.85 (s, 3H)

Example 130: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 (dd, 1H), 7.3 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.4 (s, 1H), 4.2 (d, 2H), 3.8 (s, 3H).

Example 131: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 – 7.15 (m, 3H), 6.7 (t, 1H), 4.2 (q, 2H), 3.8 (dt, 2H), 2.8 (t, 2H), 1.2 (t, 3H)

30

Example 132: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 – 7.15 (m, 3H), 6.7 (t, 1H), 4.2 (q, 2H), 3.8 (dt, 2H), 2.8 (t, 2H), 2.05 (m, 2H) 1.2 (t, 3H)

Example 133: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.3 (dd 1H), 7.2 (dd, 1H), 6.5 (s, 1H), 6.4 (t, 1H), 3.7 (s, 3H), 3.5 (dd, 2H), 2.4 (t, 2H), 1.8 (m, 4H)

Example 134: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.95 (d, 2H), 7.6 (d, 2H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.9 (t, 1H), 6.5 (s, 1H), 4.8 (d, 2H), 3.0 (s, 3H)

Example 135: ^1H NMR ($\text{DMSO}-d_6$) δ 9.1 (bs, 2H), 8.4 (s, 1H), 8.0 (t, 1H), 7.85 (d, 2H), 7.7 (d, 2H), 7.6 (m, 1H), 7.4 (m, 2H), 6.6 (s, 1H), 4.8 (bs, 2H)

Example 136: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.9 (m, 3H), 6.7 (t, 1H), 6.5 (s, 1H), 4.5 (d, 2H), 4.2 (s, 4H)

Example 137: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.3 (dd, 1H), 7.2 (dd, 1H), 6.9 (dd, 1H), 6.8 (t, 1H), 6.7 (m, 1H), 6.6 (s, 1H), 5.3 (s, 2H), 4.85 (s, 2H), 4.6 (d, 2H).

Example 138: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.9 (d, 1H), 7.8 (d, 1H), 7.4 (m, 2H), 7.3 (dd, 1H), 7.1 (dd, 1H), 6.9 (t, 1H), 6.6 (s, 1H), 4.8 (d, 2H)

Example 139: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.3 (m, 2H), 7.2 (dd, 1H), 7.1 (dd, 1H), 6.8 (d, 1H), 6.7 (t, 1H), 6.6 (s, 1H), 4.6 (m, 4H), 3.2 (t, 2H)

Example 140: ^1H NMR (CDCl_3) δ 8.45 (s, 1H), 8.2 (dt, 1H), 8.0 (s, 1H), 7.7 (dd, 1H), 7.4 – 7.3 (m, 3H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.6 (s, 1H), 4.7 (d, 2H)

Example 141: ^1H NMR (CDCl_3) δ 8.2 (dt, 1H), 8.0 (s, 1H), 7.45 – 7.1 (m, 7H), 6.6 (s, 1H), 4.4 (dt, 2H), 2.6 (t, 2H), 1.8 (m, 2H), 1.4 (m, 2H)

Example 171: ^1H NMR (CD_3OD) δ 8.41 (s, 1H), 8.25 (d, $J = 6.3$ Hz, 1H), 8.15 (s, 1H), 7.67 (d, $J = 7.8$ Hz, 2H), 7.55–7.48 (m, 2H), 7.45 (dd, $J = 7.5, 1.2$ Hz, 1H), 7.34 (dd, $J = 7.5, 1.8$ Hz, 1H), 6.28 (s, 1H), 4.79 (s, 2H).

Example 172: ^1H NMR (CDCl_3) δ 8.64 (s, 1H), 7.68–7.64 (m, 2H), 7.52 (m, 1H), 7.43 (t, $J = 7.8$ Hz, 1H), 6.89 (t, $J = 6.0$ Hz, 1H), 6.51 (s, 1H), 6.48 (m, 2H), 4.74 (d, $J = 6.0$ Hz, 2H).

Example 173: ^1H NMR ($\text{DMSO}-d_6$) δ 8.86 (s, 1H), 8.46 (s, 1H), 8.32-8.28 (m, 2H), 7.97 (m, 1H), 7.87 (m, 1H), 7.52 (m, 1H), 7.35-7.24 (m, 2H), 6.57 (s, 1H), 6.46 (m, 1H), 3.65 (m, 4H).

Example 174: ^1H NMR (CDCl_3) δ 8.37 (s, 1H), 8.16 (t, $J = 7.5$ Hz, 1H), 7.45-7.35 (m, 1H), 7.32-7.20 (m, 3H), 7.17-7.07 (m, 1H), 6.92 (t, $J = 6$ Hz, 1H), 6.48 (s, 1H), 4.65 (d, 2H), 2.50 (s, 3H).

Example 175: ^1H NMR (CDCl_3) δ 8.16 (t, $J = 9$ Hz, 1H), 8.00 (s, 1H), 7.49 (d, $J = 9$ Hz, 1H), 7.46-7.36 (m, 1H), 7.18-7.08 (m, 1H), 7.00 (d, $J = 9$ Hz, 1H), 6.62-6.50 (m, 2H), 2.60 (s, 3H), 2.55 (s, 3H).

10 **Example 176:** ^1H NMR (CDCl_3) δ 8.15 (t, $J = 9$ Hz, 1H), 8.00 (s, 1H), 7.45-7.35 (m, 1H), 7.32-7.20 (m, 1H), 7.20-7.05 (m, 3H), 6.80 (t, 1H), 6.50 (s, 1H), 4.65 (d, 2H), 2.65 (s, 3H), 2.50 (s, 3H).

Example 177: ^1H NMR (CDCl_3) δ 8.20 (t, 1H), 7.90 (s, 1H), 7.50-7.05 (m, 8H), 6.80 (s, 1H), 5.05-4.90 (m, 2H), 3.80 (d, 1H), 3.45 (d, 1H), 3.00 (dd, 1H), 2.90 (dd, 1H), 2.50 (s, 3H).

Example 181: ^1H NMR (300MHz, CDCl_3) δ 8.41 (s, 1H), 8.28 – 8.23 (d, 1H), 8.15 (s, 1H), 7.69 – 7.60 (d, 1H), 7.62 – 7.50 (m, 3H), 7.50 – 7.47 (dd, 1H), 6.35 (s, 1H), 5.36 (s, 1H), 4.80 (s, 2H).

20 **Example 184:** ^1H NMR (300MHz, CDCl_3) δ 8.96 - 8.90 (s, 1H), 8.08 (s, 1H), 8.04 (d, 1H), 7.72 (d, 1H), 7.70 – 7.61 (dd, 1H), 7.24 – 7.20 (dd, 1H), 6.92 – 6.84 (t, 1H), 6.36 (s, 1H), 4.96 – 4.89 (d, 2H).

Example 186: ^1H NMR (300MHz, CDCl_3) δ 8.96 - 8.90 (s, 1H), 8.08 (s, 1H), 8.44 (s, 1H), 8.27 – 8.24 (d, 1H), 8.02 (s, 1H), 7.78 – 7.76 (d, 1H), 7.73 – 7.70 (d, 1H), 7.58 – 7.51 (m, 2H), 7.13 – 7.08 (dd, 1H), 5.51 (s, 2H).

25 **Example 195:** ^1H NMR (CD_3OD) δ 8.40(s, 1H), 8.27(d, 1H), 8.03(s, 1H), 7.75-7.50(m, 2H), 6.10(s, 1H), 4.76(s, 2H), 4.05(m, 2H), 3.88(m, 2H), 3.52(m, 1H), 2.33(m, 1H), 2.20(m, 1H).

Example 196: ^1H NMR (CD_3OD) δ 8.73(d, 1H), 8.58(q, 1H), 8.12(s, 1H), 8.00(d, 1H), 7.54(q, 1H), 6.19(s, 1H), 4.86(s, 2H), 4.22-4.08(m, 2H), 4.03-3.93(m, 2H), 3.63(m, 1H), 2.50-2.39(m, 1H), 2.32-2.21(m, 1H).

30


Example 197: ^1H NMR (CD_3OD) δ 8.73(d, 1H), 8.58(q, 1H), 8.12(s, 1H), 8.00(d, 1H), 7.54(q, 1H), 6.19(s, 1H), 4.86(s, 2H), 4.22-4.08(m, 2H), 4.03-3.93(m, 2H), 3.63(m, 1H), 2.50-2.39(m, 1H), 2.32-2.21(m, 1H).

Example 199: ^1H NMR (300MHz, CDCl_3) δ 8.29 (s, 1H), 8.15 (br s, 1H), 7.95 (s, 1H), 7.28 (d, 1H), 7.05 - 6.95 (appt t, 1H), 5.70 (s, 1H), 4.62 (d, 2H), 2.90 (m, 1H), 2.30 (m, 1H), 1.9 - 1.2 (m, 8H), 0.65 (d, 3H).

Example 200: ^1H NMR (300MHz, CDCl_3) δ 8.71 (s, 2H), 8.00 (s, 1H), 6.13 (s, 1H), 3.59 (s, 2H), 3.01 - 2.58 (m, 1H), 2.51 - 2.45 (m, 1H), 2.44 - 2.30 (m, 1H), 2.20 (s, 3H), 2.09 - 1.95 (m, 2H), 1.85 - 1.70 (m, 2H), 0.80 - 0.76 (d, 3H).

Example 203: ^1H NMR (300MHz, CDCl_3) δ 8.10 (s, 1H), 8.08 (s, 1H), 6.27 (s, 2H), 4.95 (s, 2H), 3.00 - 2.90 (dd, 2H), 2.60 (m, 2H), 2.48 (br s, 1H), 2.39 (s, 3h), 2.25 m, 1H), 1.95 - 1.70 (m, 3H).


EXAMPLE 211:




To a solution of the compound prepared in Example 156 (100 mg, 0.23 mmol) in dry THF (4 mL) was added LiAlH_4 (1.0 M in THF, 0.110 mL, 0.110 mmol) at 0°C under N_2 . The mixture was stirred at 0°C for 1hr, warmed to 25°C , then additional LiAlH_4 (1.0 M in THF, 0.400 mL) was added, the mixture was stirred for 20 min and then quenched with MeOH (2.0 mL). The solvent was evaporated and the crude product was purified by flash chromatography using 10:1 CH_2Cl_2 :MeOH as eluent. White solid (46 mg, 49%) was obtained. LCMS: $M^+ = 416$. $\text{Mp} = 71-72^\circ\text{C}$.

EXAMPLE 212:

174



To a solution of the compound prepared in Example 156 (70 mg, 0.16 mmol) in dry THF (3 mL) was added MeMgBr (3.0 M in Et₂O, 1.10 mL, 3.20 mmol) under N₂. The mixture was stirred at 25°C for 45 min and then quenched with saturated aqueous NH₄Cl (5.0 mL). The mixture was poured into saturated aqueous NH₄Cl (30 mL) and extracted with CH₂Cl₂ (3x20 mL). The extracts were dried over Na₂SO₄ and filtered. The solvent was evaporated and the crude product was purified by flash chromatography using 20:1 CH₂Cl₂:MeOH as eluent. White solid (25 mg, 36%) was obtained. LCMS: M⁺= 444. Mp=76-80°C.

EXAMPLE 213:

15










Anhydrous DMF (40 mL) was added under N₂ to the compound prepared in Preparative Example 174 (2.50 g, 8.65 mmol) and 60 % NaH in mineral oil (346 mg, 8.65 mmol). The mixture was stirred at 25°C for 1 hr, then 2-chloro-5-chloromethylpyridine N-oxide (1.54 g, 8.65 mmol) in anhydrous DMF (20 mL) was added slowly. The mixture was stirred at 25°C for 18 hr, the solvent was evaporated and the crude product was purified by flash chromatography using

30:1 CH₂Cl₂:MeOH as eluent. So obtained solid was triturated by 50 mL of 1:1 EtOAc: hexane. Pale yellow solid (1.25 g, 34%) was obtained. LCMS: MH⁺=432. Mp=224-226 °C.




EXAMPLES 214-217:


- 5 By essentially the same procedure set forth in Example 213 combining the compounds shown in Column 2 of Table 19 with compounds in Column 3 of Table 19, the compounds shown in Column 3 of Table 19 were prepared.

TABLE 19

Ex.	Column 2	Column 3	Column 4	CMPD
214				LCMS: MH ⁺ =380; mp=°C
215				LCMS: MH ⁺ =450; mp=218-222°C
216				LCMS: MH ⁺ =466; mp=126-128°C

176

217				LCMS: M ⁺ =523
-----	---	---	--	------------------------------

EXAMPLE 218:

5

CF₃CH₂OH (3.0 mL) was added under N₂ to 60% NaH in mineral oil (40 mg, 1.0 mmol), the mixture was stirred for 20 min, then the product prepared in Example 213 (50 mg, 0.12 mmol) was added. The mixture was refluxed for 20 hr, the solvent was evaporated, and the residue was purified by flash

10 chromatography using 20:1 CH₂Cl₂:MeOH as eluent to yield pale yellow solid (35 mg, 61%). LCMS: M₂H⁺=496. Mp=208-210 °C.









EXAMPLES 219-225:







By essentially the same procedure set forth in Example 218 combining the compounds shown in Column 1 of Table 20 with the appropriate alcohol, the compounds shown in Column 2 of Table 20 were prepared.

15

TABLE 20


Ex.	Column 1	Column 2	Data
-----	----------	----------	------

219			LCMS: $M^+ = 426$; mp=126-128°C
220			LCMS: $M^+ = 483$; mp=89-91°C
221			LCMS: $M2H^+ = 442$; mp=112-114°C
222			LCMS: $MH^+ = 462$; mp=121-123°C


223			LCMS: MH ⁺ =444; mp=112-114°C
224			LCMS: M ⁺ =376; mp=°C
225			LCMS: MH ⁺ =; mp=°C

EXAMPLE 226:

179



A mixture of the product prepared in Example 213 (100 mg, 0.23 mmol) and KOH (95 mg, 1.70 mmol) in 1,2-dimethoxyethane (3 mL) and H₂O (1.5 mL) was refluxed under N₂ for 20 hr, quenched with acetic acid (0.30 mL), and the solvent was evaporated. The residue was suspended in H₂O (15 mL), filtered and the solid was washed with H₂O (15 mL) and Et₂O (10 mL). Then it was mixed with CH₂Cl₂ (2 mL) and Et₂O (2 mL) and filtered. Et₂O (5 mL) was added to the filtrate and the mixture was allowed to stand overnight. The solid was removed by filtration, washed with Et₂O and then dissolved in MeOH (5 mL). The solution was filtered and the solvent from the filtrate was evaporated. Off-white solid (5 mg, 5%) was obtained. LCMS: M⁺=412. Mp= 206-208°C.

EXAMPLE 227:





A mixture of the product prepared in Example 213 (129 mg, 0.30 mmol), N,N-dimethylethylenediamine (0.165 mL, 1.50 mmol), and diisopropylethylamine (0.10 mL) in anhydrous N-methylpyrrolidinone (1.0 mL) was stirred at 100°C for

24 hr. The solvent was evaporated, and the residue was purified by flash chromatography using 20:1 CH₂Cl₂: 7N NH₃ in MeOH as eluent to yield pale yellow solid (110 mg, 76%). LCMS: M⁺=482. Mp=76-78 °C.







EXAMPLES 228-233:

- 5 By essentially the same procedure set forth in Example 227 combining the compounds shown in Column 1 of Table 21 with the appropriate amine, the compounds shown in Column 2 of Table 21 were prepared.



TABLE 21


Ex.	Column 1	Column 2	Data
228			LCMS: M ₂ H ⁺ =467 ; mp 126-128 °C
229			LCMS: M ⁺ =481; mp=128-130 °C

181


230			LCMS: $M^+ = 494$; mp=108-110°C
231			LCMS: $M2H^+ = 482$; mp=129-133°C
232			LCMS: $M2H^+ = 482$; mp=124-126°C

182


233			LCMS: $M2H^+ = 471$; mp=88-90°C
-----	---	--	--

EXAMPLE 234:

- 5 A mixture of the product prepared in Example 213 (80 mg, 0.19 mmol) and 2.0 M methylamine in THF was stirred in a closed pressure vessel at 50°C for 72 hr. The solvent was evaporated, and the residue was purified by flash chromatography using 10:1 CH₂Cl₂: MeOH as eluent to yield pale yellow solid (40 mg, 51%). LCMS: $M2H^+ = 427$. Mp=217-219°C.

10 **EXAMPLE 235:**

By essentially the same procedure set forth in Example 234, the compound shown above was prepared. LCMS: $M2H^+ = 441$. $Mp = 98-101^\circ C$.

EXAMPLE 236:

5

The compound prepared in Preparative Example 174 (140 mg, 0.48 mmol) and the aldehyde (71 mg, 0.58 mmol) were stirred in anhydrous THF (4 mL) at $50^\circ C$ under N_2 . $Ti(OiPr)_4$ (0.574 mL, 1.92 mmol) was added, the mixture was stirred at $50^\circ C$ 3 hr, and cooled to $25^\circ C$. $NaBH_3CN$ (181 mg, 2.88 mmol) was added, the mixture was stirred for 2 more hr, then poured into 10 % aqueous Na_2CO_3 (100 mL), and extracted with CH_2Cl_2 (3 x 50 mL). Combined extracts were dried over Na_2SO_4 , filtered, and the solvent was evaporated. The residue was purified by flash chromatography using 15:1 CH_2Cl_2 :MeOH as eluent to yield pale yellow solid (40 mg, 21%). LCMS: $MH^+ = 398$. $Mp > 230^\circ C$.

15













EXAMPLES 237-256:













By essentially the same procedure set forth in Example 236 combining the compounds shown in Column 2 and 3 of Table 22, the compounds shown in Column 4 of Table 22 were prepared.













20

TABLE 22













Ex.	<u>Column 2</u>	<u>Column 3</u>	<u>Column 4</u>	<u>Data</u>
-----	-----------------	-----------------	-----------------	-------------













237				LCMS: $M^+ = 381$; mp > 200°C
238				LCMS: $M^+ = 387$; mp = °C
239				LCMS: $MH^+ = 413$; mp = 157-159°C
240				LCMS: $M2H^+ = 419$; mp = 77-79°C

241				LCMS: M2H ⁺ =385 ; mp=214-216°C
242				LCMS: MH ⁺ =; mp=°C
243				LCMS: M ⁺ =416; mp=80-82°C
244				

245				
246				LCMS: M ⁺ =452; mp=54-56°C
247				LCMS: MH ⁺ = 401; mp>200°C
248				LCMS: M2H ⁺ = 474; mp>200.0. °C dec.


187

249				LCMS: MH ⁺ = 377; mp= 65-67°C
250				LCMS: M2H ⁺ =421; mp=87-93°C
251				LCMS: MH ⁺ =361; mp>225°C
252				LCMS: MH ⁺ =346; mp=270-271°C

253				LCMS: $MH^+ = 402$; mp=250-255°C
254				LCMS: $MH^+ = 416$; mp=210-215°C
255				LCMS: $MH^+ = 428$; mp=145°C
256				LCMS: $MH^+ =$; mp=°C

EXAMPLE 257:

189







A mixture of the compound prepared in Example 242 (100 mg, 0.24 mmol), conc. aqueous HCl (1.0 mL) and acetic acid (2.0 mL) were stirred at 100°C under N₂ for 2 hr, then poured onto Na₂CO₃ (15 g), and extracted with 1:1 acetone:CH₂Cl₂ (3 x 30 mL). Combined extracts were filtered, and the solvent was evaporated. The residue was purified by flash chromatography using 10:1 CH₂Cl₂:MeOH as eluent to yield pale yellow solid (36 mg, 37%). LCMS: M₂H⁺=398.


EXAMPLES 258-260:

By essentially the same procedure set forth in Example 257 starting from the compounds shown in Column 1 of Table 23, the compounds shown in Column 2 of Table 23 were prepared.


TABLE 23

Ex.	Column 1	Column 2	Data
258			LCMS: M ⁺ =402; mp=229-231°C

259			LCMS: MH ⁺ =416; mp=215- 218°C
260			LCMS: M2H ⁺ =398 mp>230°C

EXAMPLE 261:

- 5 To a stirred solution of the compound prepared in Example 239 (41 mg, 0.10 mmol) in CH₂Cl₂ was added 1.0 M BBr₃ (0.30 mL, 0.30 mmol) in CH₂Cl₂ at -78°C. The mixture was stirred at -78°C for 5 min, then at 24°C for 3 hr, then MeOH (2.0 mL) was added and the mixture was stirred for 10 min. The solvent was evaporated and the residue was purified by flash chromatography using
- 10 5:1:0.1 CH₂Cl₂:MeOH:conc. NH₄OH as eluent to yield white solid (39 mg, 99%). LCMS: M⁺=397. Mp>230 °C.

EXAMPLE 262:

- 5 A mixture of the product prepared in Example 217 (40 mg, 0.077 mmol) and 5.0 M aqueous NaOH (0.8 mL) in MeOH (3.0 mL) was refluxed under N₂ for 1 hr. NaHCO₃ (700 mg) was added, the solvent evaporated, and the residue was purified by flash chromatography using 10:1:0.1 CH₂Cl₂: MeOH: conc. NH₄OH as eluent to yield white solid (10 mg, 35%). LCMS: M₂H⁺=371. Mp=237-239 °C.



10 **EXAMPLES 263-264:**


By essentially the same procedure set forth in Example 262 starting from the compounds shown in Column 1 of Table 24, the compounds shown in Column 2 of Table 24 were prepared.

TABLE 24

Ex.	Column 1	Column 2	Data
263			LCMS: M ₂ H ⁺ =370 ; mp=166-168 °C

192

264			LCMS: $M2H^+ = 371$; mp=180- 182°C
-----	---	--	--


EXAMPLE 265:

- TFA (0.5 mL) was added to a solution of the compound prepared in
- 5 Preparative Example 197 (0.08 g, 0.16 mmol) in CH_2Cl_2 (2.0 mL) at 0°C and the resulting solution stirred 2.5 hours and stored at 4°C overnight at which time additional TFA (0.5 mL) was added. The resulting solution was stirred 4 hours and concentrated *in vacuo*. The residue was neutralized with 1N NaOH and extracted with CH_2Cl_2 . The combined organics were dried over Na_2SO_4 , filtered,
- 10 and concentrated under reduced pressure. The crude product was purified by flash chromatography using a 2.5% (10% NH_4OH in MeOH) in CH_2Cl_2 solution as eluent (0.009 g, 15% yield). LCMS: $MH^+ = 396$; mp= 53-54°C.

15

EXAMPLE 266:

193




A solution of the compound prepared in Preparative Example 182 (26 mg, 0.070 mmol) and potassium thiocyanate (13 mg, 0.14 mmol) in MeOH (1 mL)

5 was cooled in a cold water bath. To it was added a solution of bromine (22 mg, 0.14 mmol) in MeOH (0.7 mL) dropwise. The resulting reaction mixture was stirred for 4 h at room temperature and the volatiles were removed under reduced pressure. The residue obtained was suspended in a small amount of CH₂Cl₂. The potassium bromide was filtered off and pH of the filtrate was

10 adjusted to about 7 by the addition of aqueous ammonia. It was concentrated under reduced pressure and the residual oil was purified by preparative thin-layer chromatography using 15% MeOH in CH₂Cl₂ as eluent (26 mg, 87% yield). ¹H NMR (CDCl₃) δ 8.75 (d, J = 4.2 Hz, 2H), 8.38 (s, 1H), 7.68-7.64 (m, 2H), 7.46-7.39 (m, 3H), 7.22 (t, J = 6.3 Hz, 1H), 6.43 (s, 1H), 4.84 (d, J = 6.3 Hz, 2H);

15 LCMS: MH⁺ = 427.

EXAMPLE 267:




Boron tribromide (1 M in CH₂Cl₂, 0.60 mL, 0.60 mmol) was added dropwise to an ice-cold stirred solution of the compound prepared in Example 24 (50 mg, 0.12 mmol) in CH₂Cl₂ (1.5 mL) under an argon atmosphere. The

resulting reaction mixture was stirred at 0°C for 30 minutes, allowed to warm up to room temperature, and stirred overnight. The mixture was quenched by the addition of a small amount of water and extracted with CH₂Cl₂. The organic layer was dried over magnesium sulfate and concentrated *in vacuo* (45 mg, 94%

5 yield). ¹H NMR (CD₃OD) δ 9.16 (s, 1H), 8.95 (s, 1H), 8.88 (d, J = 8.1 Hz, 1H), 8.24 (t, J = 6.9 Hz, 1H), 8.18 (s, 1H), 7.95 (d, J = 7.8 Hz, 1H), 7.40 (t, J = 7.8 Hz, 1H), 7.00-6.96 (m, 2H), 6.86 (s, 1H), 5.28 (s, 2H); LCMS: MH⁺ = 396.

EXAMPLE 268:



10

A solution of the compound from Preparative Example 184 (0.05 g, 0.15 mmol), N-methylpiperazine (20 μL, 1.2 eq.) and iPr₂Et (52 μL, 2.0 eq.) in dioxane (1 mL) was heated to 70 °C overnight. The reaction mixture was cooled to room temperature and diluted with H₂O and saturated NaHCO₃. The resulting mixture was extracted with CH₂Cl₂, the combined organics dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude product was purified by Preparative TLC using a 5% (10% NH₄OH in MeOH) in CH₂Cl₂ solution as eluent (0.028 g, 47% yield). MS: MH⁺ = 402. mp = 210 °C (dec.)

15













EXAMPLES 269-275:










20 By essentially the same procedure set forth in Example 268 only substituting the amine in Column 2 of Table 25 and the chlorides in Column 3 of Table 25, the compounds shown in Column 4 of Table 25 are prepared:


TABLE 25

Ex.	Column 2	Column 3	Column 4	CMPD
-----	----------	----------	----------	------

195


269				MS: MH^+ = 387 m.p. 182 – 183 °C
270				MS: MH^+ = 373 m.p. 190 – 191 °C
271				MS: MH^+ = 403 m.p. 227 – 230 °C
272				MS: MH^+ = 388 m.p. 198 – 201 °C

273				MS: MH^+ = 430 m.p. 100-103 °C
274				MS: MH^+ = 456 m.p. 175-178 °C
275				MS: MH^+ = 403 m.p. 218 °C

EXAMPLE 276:**Step A:**


- 4-Fluorophenyl magnesium bromide (0.68 mL, 1.2 eq.) was added to the compound prepared in Preparative Example 193 (0.20 g, 0.55 mmol) and $\text{PdCl}_2(\text{dppf})_2$ (0.037 g, 10 mol%) in THF and the resulting solution was stirred at room temperature 72 hours. The reaction mixture was dilute with saturated NH_4Cl and extracted with EtOAc. The combined organics were washed with saturated NaCl, dried over Na_2SO_4 , filtered and concentrated. The crude product was purified by flash chromatography using neat EtOAc as eluent (0.15 g, 65% yield). MS: $\text{MH}^+ = 420$.

Step B:



By essentially the same procedure set forth in Preparative Example 127 only substituting the compound prepared in Example 276, Step A, the above compound was prepared (0.17 g, 94% yield).


Step C:




By essentially the same procedure set forth in Preparative Example 200 only substituting the compound prepared in Example 276, Step B, the above compound was prepared (0.1g, 100% yield).

Step D:

198




By essentially the same procedure set forth in Example 265 only substituting the compound prepared in Example 276, Step C, the above
 5 compound was prepared (0.049 g, 62% yield). MS: $MH^+ = 414$; mp = 110-115 °C.

EXAMPLE 277:Step A:

$Pd(PPh_3)_4$ (0.065 g, 10 mol%) was added to 3-cyanophenyl zinc iodide
 10 (2.2 mL, 0.5 M solution in THF, 2 eq.) and the compound prepared in Preparative Example 193 (0.2 g, 0.56 mmol) in DMF (2.0 mL) and the resulting solution heated to 80 °C for 144 hours. The reaction mixture was cooled to room temperature, diluted with saturated NH_4Cl and extracted with EtOAc. The combined organics were washed with H_2O and brine, dried over Na_2SO_4 , filtered,
 15 and concentrated under reduced pressure. The crude product was purified by flash chromatography using a neat EtOAc solution as eluent (0.07 g, 29% yield). MS: $MH^+ = 427$.


Step B through Step D:

199



By essentially the same procedures set forth in Example 276, Step B through Step D, the above compound was prepared (0.023 g, 53% yield). MS: $MH^+ = 421$; mp = 230 °C (dec.)


5 EXAMPLE 278:



By essentially the same procedure set forth in Example 276 only substituting the appropriate cyclopropylmagnesium bromide in Step A, the compound was prepared. MS: $MH^+ = 372$; m. p. = 96-98 °C.

10

EXAMPLE 279:



15




The palladium-catalyzed zinc cross-coupling reaction was carried out in a manner similar to the procedure described in *J. Org. Chem.* (1999), 453. A solution of the chloropyrazolopyrimidine (200 mg, 0.458 mmol), $Pd(PPh_3)_4$ (53













mg, 0.046 mmol), and *exo*-2-norbornylzinc bromide (0.5 M in THF, 0.95 mL, 0.47 mmol) in DMF (2 mL) was refluxed at 100°C (oil bath temp.) overnight. The reaction mixture was quenched with half-saturated NH₄Cl and extracted with CH₂Cl₂. The organic phase was dried over MgSO₄ and concentrated under reduced pressure. The residue was purified by flash chromatography using a 50% EtOAc in hexanes solution as eluent. A solution of the obtained *N*-Boc-protected product (121 mg, 53% yield, LCMS: MH⁺ = 498) and TFA (1 mL) in CH₂Cl₂ (2 mL) was stirred at room temperature for 2 hr. The volatiles were removed under reduced pressure. The residue was dissolved in CH₂Cl₂, neutralized with saturated NaHCO₃, and extracted with CH₂Cl₂. The organic phase was dried over MgSO₄ and concentrated *in vacuo* (96 mg, 99% yield). LCMS: MH⁺ = 398; ¹H NMR (CDCl₃) δ 8.78 (s, 1H), 8.71 (d, J = 4.2 Hz, 1H), 8.04 (d, J = 3.9 Hz, 1H), 7.80 (d, J = 7.8 Hz, 1H), 7.44 (m, 1H), 6.73 (m, 1H), 5.98 (d, J = 7.5 Hz, 1H), 4.74 (d, J = 5.4 Hz, 2H), 3.40-1.00 (m, 11H).

15 EXAMPLES 280-294:
















By following essentially the same procedure set forth in Example 279 only substituting the chlorides shown in Column 2 of Table 26 and the organozinc reagents shown in Column 3 of Table 26, the compounds in Column 4 of Table 26 were prepared:













20 Table 26




Ex.	Column 2	Column 3	Column 4	Data
280				LCMS: MH ⁺ = 395

281				LCMS: $MH^+ = 400$
282				LCMS: $MH^+ = 412$
283				LCMS: $MH^+ = 452$
284				LCMS: $MH^+ = 422$

202

285				LCMS: $MH^+ = 408$
286				LCMS: $MH^+ = 404$
287				LCMS: $MH^+ = 404$
288				LCMS: $MH^+ = 408$
289				LCMS: $MH^+ = 386$

290				LCMS: $MH^+ = 464$
291				LCMS: $MH^+ = 480$
292				LCMS: $MH^+ = 424$
293				LCMS: $MH^+ = 424$

294				LCMS: $MH^+ = 426$
-----	---	---	--	--------------------

Additional data for select compounds is shown below.

EXAMPLE 280: 1H NMR ($CDCl_3$) δ 8.65 (s, 1H), 8.57 (d, $J = 4.2$ Hz, 1H), 8.50 (d, $J = 4.5$ Hz, 1H), 8.01 (s, 1H), 7.69 (d, $J = 7.5$ Hz, 1H), 7.61 (d, $J = 7.8$ Hz, 1H), 7.31-7.22 (m, 2H), 6.77 (m, 2H), 4.71 (d, $J = 5.4$ Hz, 2H), 2.68 (s, 3H).

EXAMPLE 281: 1H NMR ($CDCl_3$) δ 8.80 (s, 1H), 8.72 (d, $J = 4.8$ Hz, 1H), 8.08 (s, 1H), 7.85-7.40 (m, 3H), 7.02 (d, $J = 5.1$ Hz, 1H), 6.90 (t, $J = 6.0$ Hz, 1H), 6.29 (s, 1H), 4.79 (d, $J = 6.0$ Hz, 2H), 2.61 (s, 3H).

EXAMPLE 282: 1H NMR ($CDCl_3$) δ 8.67 (s, 1H), 8.61 (d, $J = 3.9$ Hz, 1H), 8.03 (s, 1H), 7.72-7.31 (m, 3H), 7.22-7.00 (m, 2H), 6.81 (t, $J = 6.0$ Hz, 1H), 6.03 (s, 1H), 4.68 (d, $J = 6.0$ Hz, 2H), 2.28 (s, 3H).

EXAMPLE 283: 1H NMR ($CDCl_3$) δ 8.68 (s, 1H), 8.63 (d, $J = 4.0$ Hz, 1H), 8.00 (s, 1H), 7.80-7.72 (m, 2H), 7.54-7.47 (m, 3H), 7.35 (m, 1H), 6.74 (t, $J = 6.0$ Hz, 1H), 6.19 (s, 1H), 4.67 (d, $J = 6.0$ Hz, 2H), 4.21 (q, $J = 7.2$ Hz, 2H), 1.13 (t, $J = 7.2$ Hz, 3H).

EXAMPLE 284: 1H NMR ($CDCl_3$) δ 7.97 (s, 1H), 7.65 (d, $J = 7.2$ Hz, 1H), 7.33-7.15 (m, 5H), 6.73 (t, $J = 5.4$ Hz, 1H), 5.99 (s, 1H), 4.61 (d, $J = 5.4$ Hz, 2H), 3.09 (sept, $J = 6.9$ Hz, 1H), 1.11 (d, $J = 6.9$ Hz, 6H).

EXAMPLE 285: 1H NMR ($CDCl_3$) δ 8.56-8.55 (m, 2H), 7.94 (s, 1H), 7.54 (m, 1H), 7.30-7.22 (m, 6H), 6.59 (t, $J = 5.7$ Hz, 1H), 5.66 (s, 1H), 4.47 (d, $J = 5.7$ Hz, 2H), 4.26 (q, $J = 7.2$ Hz, 1H), 1.68 (d, $J = 7.2$ Hz, 3H).

EXAMPLE 286: 1H NMR ($CDCl_3$) δ 8.67 (m, 2H), 7.94 (s, 1H), 7.69 (d, $J = 7.8$ Hz, 1H), 7.34 (m, 1H), 6.63 (t, $J = 5.7$ Hz, 1H), 5.87 (s, 1H), 4.62 (d, $J = 5.7$ Hz, 2H), 3.64 (s, 3H), 3.13 (m, 2H), 2.82 (m, 1H), 1.22 (m, 3H).

EXAMPLE 287: ^1H NMR (CDCl_3) δ 8.66 (m, 2H), 7.94 (s, 1H), 7.68 (d, $J = 7.8$ Hz, 1H), 7.34 (m, 1H), 6.62 (t, $J = 6.0$ Hz, 1H), 5.87 (s, 1H), 4.62 (d, $J = 6.0$ Hz, 2H), 3.64 (s, 3H), 3.13 (m, 2H), 2.81 (m, 1H), 1.22 (m, 3H).

EXAMPLE 288: ^1H NMR (CDCl_3) δ 8.64 (s, 1H), 8.60 (d, $J = 3.6$ Hz, 1H), 8.04 (s, 1H), 7.68 (m, 1H), 7.31 (m, 1H), 7.16 (m, 1H), 7.07-7.05 (m, 2H), 6.80 (t, $J = 6.3$ Hz, 1H), 5.93 (s, 1H), 4.64 (d, $J = 6.3$ Hz, 2H), 2.08 (s, 6H).

EXAMPLE 289: ^1H NMR (CDCl_3) δ 8.72 (s, 1H), 8.62 (d, $J = 4.8$ Hz, 1H), 7.99-7.97 (m, 2H), 7.73-7.69 (m, 2H), 7.40-7.33 (m, 2H), 6.67 (t, $J = 6.0$ Hz, 1H), 6.29 (s, 1H), 4.71 (d, $J = 6.0$ Hz, 2H).

EXAMPLE 290: ^1H NMR (CDCl_3) δ 8.73 (s, 1H), 8.62 (d, $J = 4.5$ Hz, 1H), 8.01 (s, 1H), 7.76 (m, 1H), 7.41 (d, $J = 5.1$ Hz, 1H), 7.34 (dd, $J = 8.1, 5.1$ Hz, 1H), 7.05 (d, $J = 5.1$ Hz, 1H), 7.01 (s, 1H), 6.79 (t, $J = 6.0$ Hz, 1H), 4.74 (d, $J = 6.0$ Hz, 2H).


EXAMPLE 291: ^1H NMR ($\text{DMSO}-d_6$) δ 9.12 (s, 1H), 8.40 (s, 1H), 8.33 (s, 1H), 8.13 (m, 1H), 7.82 (d, $J = 5.1$ Hz, 1H), 7.40-7.39 (m, 2H), 7.22 (d, $J = 5.1$ Hz, 1H), 6.86 (s, 1H), 4.86 (s, 2H).

EXAMPLE 292: ^1H NMR (CDCl_3) δ 8.23 (s, 1H), 8.16 (d, $J = 6.0$ Hz, 1H), 8.06 (s, 1H), 7.31-7.05 (m, 5H), 6.86 (m, 1H), 5.87 (s, 1H), 4.62 (d, $J = 6.3$ Hz, 2H), 2.09 (s, 6H).


EXAMPLE 293: ^1H NMR (CDCl_3) δ 8.14 (s, 1H), 8.12 (d, $J = 6.3$ Hz, 1H), 7.94 (s, 1H), 7.29-7.16 (m, 6H), 7.07 (m, 1H), 6.78 (t, $J = 6.0$ Hz, 1H), 5.54 (s, 1H), 4.44 (d, $J = 6.0$ Hz, 2H), 4.24 (t, $J = 7.2$ Hz, 1H), 1.68 (d, $J = 7.2$ Hz, 3H).

EXAMPLE 294: ^1H NMR (CDCl_3) δ 8.67 (s, 1H), 8.59 (d, $J = 4.8$ Hz, 1H), 8.01 (s, 1H), 7.71 (m, 1H), 7.52 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.40-7.19 (m, 4H), 6.78 (t, $J = 6.0$ Hz, 1H), 6.32 (s, 1H), 4.67 (d, $J = 6.0$ Hz, 2H), 2.38 (s, 3H).

EXAMPLE 295:




To a suspension of lithium aluminum hydride (10 mg, 0.26 mmol) in anhydrous THF (2 mL) at 0°C was added dropwise a solution of the compound prepared in Example 283 (20 mg, 0.044 mmol) in anhydrous THF (2 mL). The resulting mixture was refluxed for 1 hr and stirred at room temperature overnight, neutralized with dilute sulfuric acid, and extracted with EtOAc. The organic phase was dried over MgSO₄ and concentrated under reduced pressure. The crude product was purified by preparative thin-layer chromatography using a 5% MeOH in EtOAc solution as eluent (15 mg, 83% yield). LCMS: MH⁺ = 410; ¹H NMR (CDCl₃) δ 8.69 (s, 1H), 8.61 (d, J = 3.9 Hz, 1H), 8.05 (d, J = 2.1 Hz, 1H), 7.74 (d, J = 7.8 Hz, 1H), 7.52-7.31 (m, 5H), 6.97 (t, J = 6.3 Hz, 1H), 6.55 (d, J = 2.7 Hz, 1H), 6.20 (s, 1H), 4.71 (d, J = 6.3 Hz, 2H), 4.52 (s, 2H).


EXAMPLE 296:

To a solution of the *N*-Boc-protected compound prepared in Example 294 (45 mg, 0.085 mmol) in CH₂Cl₂ (4 mL) at -50°C was added *m*-CPBA (18 mg, 0.10 mmol). After stirring for 1 hr at -50°C more *m*-CPBA (4 mg, 0.02 mmol) was added. The mixture was stirred for a further 2 hr, diluted with CH₂Cl₂ (20 mL), and washed with saturated NaHCO₃ (20 mL). The organic phase was dried over MgSO₄ and concentrated under reduced pressure. The residue was purified by preparative thin-layer chromatography using a 2.5% MeOH in CH₂Cl₂ solution as eluent. A solution of the obtained *N*-Boc-protected product (37 mg, 80% yield, LCMS: MH⁺ = 542) and TFA (1 mL) in CH₂Cl₂ (2 mL) was stirred at room temperature for 2 hr. The volatiles were removed under reduced pressure. The residue was dissolved in CH₂Cl₂, neutralized with saturated NaHCO₃, and

extracted with CH_2Cl_2 . The organic phase was dried over MgSO_4 and concentrated under reduced pressure. The crude product was purified by preparative thin-layer chromatography using a 5% MeOH in EtOAc solution as eluent (26 mg, 89% yield). LCMS: $\text{MH}^+ = 442$; ^1H NMR (CDCl_3) δ 8.71 (s, 1H), 8.64 (d, $J = 3.9$ Hz, 1H), 8.41 (m, 1H), 8.03 (s, 1H), 7.75-7.54 (m, 4H), 7.36 (dd, $J = 8.1, 5.1$ Hz, 1H), 6.81 (t, $J = 6.0$ Hz, 1H), 6.34 (s, 1H), 4.74 (d, $J = 6.0$ Hz, 2H), 3.25 (s, 3H).

EXAMPLE 297:

To a solution of the *N*-Boc-protected compound prepared in Example 294 (56 mg, 0.11 mmol) in CH_2Cl_2 (4 mL) at 0°C was added *m*-CPBA (42 mg, 0.24 mmol). After stirring for 2 hr at room temperature more *m*-CPBA (13 mg, 0.075 mmol) was added. The mixture was stirred at room temperature overnight, diluted with CH_2Cl_2 (20 mL), and washed with saturated NaHCO_3 (20 mL). The organic phase was dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by preparative thin-layer chromatography using a 2.5% MeOH in EtOAc solution as eluent. A solution of the obtained *N*-Boc-protected product (29 mg, 49% yield, LCMS: $\text{MH}^+ = 558$) and TFA (1 mL) in CH_2Cl_2 (2 mL) was stirred at room temperature for 2 hr. The volatiles were removed under reduced pressure. The residue was dissolved in CH_2Cl_2 , neutralized with saturated NaHCO_3 , and extracted with CH_2Cl_2 . The organic phase was dried over MgSO_4 and concentrated under reduced pressure. The crude product was purified by preparative thin-layer chromatography using a 2.5% MeOH in EtOAc solution as eluent (21 mg, 90% yield). LCMS: $\text{MH}^+ = 458$; ^1H NMR (CDCl_3) δ 8.64 (s, 2H), 8.20 (m, 1H), 8.01 (s, 1H), 7.73-7.60 (m, 3H), 7.46 (m, 1H), 7.35 (s, 1H), 6.82 (t, $J = 5.9$ Hz, 1H), 6.17 (s, 1H), 4.65 (d, $J = 5.7$ Hz, 2H), 3.60 (s, 3H).

EXAMPLE 298


- 5 By essentially the same procedure set forth in Preparative Example 127, only substituting the compound prepared in Preparative Example 189, the above compound was prepared. MS: $MH^+ = 334$; mp = 170-173 °C.

Examples 299-300:


- 10 By essentially the same procedure set forth in Example 298 only substituting the compound shown in Table 27, Column 2, the compounds shown in Table 27, Column 3 were prepared:

Table 27

Ex.	Column 2	Column 3	CMPD
299			MS: $MH^+ = 348$ m.p. = 73 – 83 °C
300			MS: $MH^+ = 362$ m.p. = 165 – 175 °C

EXAMPLE 301:


To a solution of the compound prepared in Preparative Example 186 (0.1
5 g, 0.21 mmol) in THF (4.0 mL) at -78 °C was added nBuLi (0.57 mL, 2.16M in
hexanes, 5.0 eq.) at -78 °C. The reaction mixture was stirred 2 hours at -78 °C,
quenched with H₂O, warmed to room temperature, and extracted with EtOAc.
The combined organics were dried over Na₂SO₄, filtered, and concentrated under
reduced pressure. The crude product was purified by Preparative TLC using a
10 2.5% (10%NH₄OH in CH₃OH) solution in CH₂Cl₂ as eluent (0.013 g, 20% yield).
MS: MH⁺ = 326; mp = 71-72 °C.

EXAMPLE 302:


15 By essentially the same procedure set forth in Example 301 only
substituting the compound from Preparative Example 187, the above compound
was prepared (0.049 g, 68% yield). MS: MH⁺ = 344; mp = 69-71 °C.

EXAMPLE 303:

210



To a solution of 3-H adduct from Preparative Example 187.1 (0.70 g, 2.32 mmol) in DMF (4.2 mL) at 0 °C was added POCl₃ (0.67 mL, 7.2 mmol) dropwise. The mixture was stirred for 14h at rt, cooled to 0 °C, and was quenched by addition of ice. 1N NaOH was carefully added to adjust pH to 8 and the mixture was extracted with CH₂Cl₂ (3 x 25 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was recrystallized from EtOAc to afford 0.43 g (56%) of a yellow solid. mp 181-183 °C; M+H = 330.

EXAMPLE 304:**STEP A:**

To a solution of aldehyde (100 mg, 0.30 mmol) from Example 303 in THF (1 mL) at 0 °C was added cyclohexyl magnesium bromide (0.46 mL, 2.0M in Et₂O) dropwise over 5 min. The resulting mixture was stirred at 0 °C for 2h and at rt for 12h. The mixture was cooled to 0 °C and was treated with sat. aq. NH₄Cl (3 mL) and CH₂Cl₂ (5 mL). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 5 mL). The organic layers were combined, washed with brine (1 x 5 mL), dried (Na₂SO₄), filtered and concentrated under

reduced pressure to afford 110 mg (89%) of a light yellow semisolid. M+H = 414. This material was carried on crude to Step B without further purification.



STEP B:

To a solution of alcohol (53 mg, 0.13 mmol) in CH₂Cl₂ (0.5 mL) at 0 °C was added Et₃SiH (24 μL, 0.15 mmol) followed by TFA (24 μL, 0.30 mmol). The mixture was stirred for 2 h at 0 °C and rt for 2 h whereupon additional portions of Et₃SiH (24 μL, 0.15 mmol) and TFA (24 μL, 0.30 mmol) were added and the mixture was stirred for 3 h at rt (until complete by TLC). The mixture was concentrated under reduced pressure and the crude residue was partitioned between CH₂Cl₂ (5 mL) and sat. aq. NaHCO₃ (2.5 mL). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 5 mL). The organic layers were combined, washed with brine (1 x 5 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude product was purified by prep TLC (8 x 1000 mM) eluting with CH₂Cl₂/MeOH (22:1) to afford 29 mg (56%) of a yellow semisolid. M+H = 398.









EXAMPLES 305-312:

By essentially the same procedure set forth in Example 304, utilizing the aldehyde from Example 303 and substituting the Grignard or organolithium reagents shown in Column 2 of Table 28, the compounds in Column 3 of Table 28 were prepared:







TABLE 28


Ex.	Column 2 (Organometallic)	Column 3 (Final Structure)	CMPD 1. mp (°C) 2. M+H
305			1. yellow oil 2. M+H = 392

212

306			1. red oil 2. M+H = 353
307			1. red oil 2. M+H = 398
308			1. yellow oil 2. M+H = 406
309			1. yellow semisolid 2. M+H = 384

213


310			1. semisolid 2. M+H = 340
311			1. mp = 141-143 2. M+H = 358
312			1. mp = 148-150 2. M+H = 372

EXAMPLE 313:

5

To solution of aldehyde (81 mg, 0.25 mmol) from Example 303 in benzene (2.5 mL) was added carboethoxymethylene triphenyl phosphorane


(0.12 g, 0.33 mmol) in one portion. The mixture was heated at reflux for 24h, cooled to rt, and concentrated under reduced pressure. The mixture was diluted CH₂Cl₂ (5 mL), brine (2 mL) was added, and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (2 x 4 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (8 x 1000 μ M) eluting with CH₂Cl₂/MeOH (20:1) to afford 98 mg (100%) of white solid. mp 151-153 °C; M+H = 400.

EXAMPLE 314:


To a mixture of benzyltriphenylphosphonium bromide (0.59 g, 1.37 mmol) in THF (3 mL) was added NaH (55 mg, 1.37 mmol) and the mixture was stirred for 30 min. The aldehyde (0.15 g, 0.46 mmol) from Example 303 was added in a single portion and the mixture was heated at reflux for 36h. The mixture was cooled to rt and was concentrated under reduced pressure. The mixture was diluted CH₂Cl₂ (5 mL), brine (2 mL) was added, and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (2 x 4 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (8 x 1000 μ M) eluting with CH₂Cl₂/MeOH (20:1) to afford 58 mg (32%) of yellow solid. mp 138-141 °C; M+H = 404.

EXAMPLE 315:

215




To a solution of aldehyde (0.20 g, 0.60 mmol) from Example 303 in THF (3 mL) was added Ti (i-OPr)₄ (0.36 mL, 1.21 mmol) dropwise followed by addition of (S)-(-)-2-methyl-2-propanesulfinamide (74 mg, 0.61 mmol). The resulting mixture was stirred for 18h at reflux, cooled to rt, and quenched with brine (2 mL). The mixture was filtered thru a pad of Celite which was washed with EtOAc (2 x 2 mL). The layers were separated and the aqueous layer was extracted with EtOAc (2 x 4 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (8 x 1000 μM) eluting with CH₂Cl₂/MeOH (20:1) to afford 0.21 g (80%) of yellow solid. mp 108-110 °C; M+H = 433.

EXAMPLE 316:

Prepared in the same fashion as Example 315 except substituting (R)-(-)-2-methyl-2-propanesulfinamide to afford 0.25 g (94%) as a yellow solid. mp 107-109 °C; M+H = 433.

EXAMPLE 317:

216

**STEP A:**


- 5 To a solution of sulfinimine (50 mg, 0.12 mmol) from Example 316 in CH₂Cl₂ (2.5 mL) at -40 °C was added MeMgBr (96 mL, 0.29 mmol) dropwise. The mixture was stirred for 5h at -40 °C and was stirred at rt for 12h. An additional portion of MeMgBr (96 mL, 0.29 mmol) and the mixture was stirred for 12 h. Sat. aq. NH₄Cl (2 mL) was added and the mixture was extracted with
- 10 EtOAc (3 x 4 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure to afford 30 mg (58%) of crude residue. This material was taken onto the next step without purification.

STEP B:

- The crude material from Step A (30 mg, 0.067 mmol) in MeOH (2 mL)
- 15 was added conc. HCl (2 mL). The mixture was stirred at rt for 12h and the mixture was concentrated to dryness. The crude material was partitioned between CH₂Cl₂ (3 mL) and sat. aq. NaHCO₃ (2 mL) and the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (2 x 3 mL) and the organic layers were combined. The organic layer was dried (Na₂SO₄), filtered,
- 20 and concentrated under reduced pressure to afford 6 mg (24%) of the title compound as a light yellow solid. mp 100-102 °C; M + H = 345.


EXAMPLE 318:

217



To a solution of aldehyde (75 mg, 0.23 mmol) from Example 300 in
 5 THF/CH₂Cl₂ (5 mL/1mL) at rt was added MeONH₂·HCl (38 mg, 0.46 mmol)
 followed by dropwise addition of pyridine (46 μL, 0.57 mmol). The mixture was
 stirred for 72h at rt whereupon the mixture was concentrated to dryness. The
 crude material was partitioned between CH₂Cl₂ (3 mL) and sat. aq. NaHCO₃ (2
 mL) and the layers were separated. The aqueous layer was extracted with
 10 CH₂Cl₂ (2 x 3 mL) and the organic layers were combined. The organic layer was
 dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude
 product was purified by preparative TLC (3 x 1000 μM) eluting with
 CH₂Cl₂/MeOH (22:1) to afford 90 mg (100%) of light yellow solid. mp 173-175 °C
 ; M + H = 359.

15


EXAMPLE 319:

To solution of aldehyde (60 mg, 0.18 mmol) from Example 303 at EtOH
 20 (2.5 mL) was added oxindole (48 mg, 0.37 mmol) followed by piperidine (3
 drops). The mixture was heated at reflux for 14h and the mixture was cooled to

rt. The resultant precipitate was filtered and washed with cold EtOH (2 x 2 mL). The product was dried under high vacuum to afford 81 mg (100%) of the title compound as an orange/brown solid. mp 182-185 °C; M+H = 445.

EXAMPLE 320:

5



To a solution of 3-H analog (106 mg, 0.35 mmol) from Preparative Example 187.10 in AcOH (2 mL) was added 37% aqueous formaldehyde (1.5 ml; 1.40 mmol) followed by piperidine (100 μ L; 0.37 mmol). The resulting mixture was stirred at rt for 24h and the AcOH was removed under reduced pressure. The mixture was diluted with water (2 mL) and neutralized with 2M NaOH until pH = 8. The aqueous layer was extracted with CH₂Cl₂ (3 x 7 mL) and the organic layers were combined. The organic layer was washed with brine (1 x 4 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to afford 96 mg (69%) of an off-white solid. mp 88-90 °C; M+H 399.

EXAMPLES 321-322:


By essentially the same procedure set forth in Example 320 only substituting the amines in Column 2 of Table 29 and employing the 3-H adduct from Preparative Example 187.10, the compounds in Column 3 of Table 29 were prepared:

TABLE 29

Ex.	<u>Column 2</u> (Amine)	<u>Column 3</u> (Final Structure)	<u>CMPD</u> 1. mp (°C) 2. M+H
321			1. mp = 178-180 2. M+H = 401

219

322			1. mp = 102-104 2. M+H = 414


EXAMPLE 323:

5

To a solution of 3-H analog (113 mg, 0.38 mmol) from Preparative Example 187.10 in CH₂Cl₂ (5 mL) at rt was added AlCl₃ (215 mg, 1.61 mmol) followed by AcCl (100 mL, 1.40 mmol). The mixture was heated at reflux for 12h and was cooled to rt. The mixture was treated sequentially with 3M HCl (3 mL) followed by sat. aq. NaHCO₃ (until pH = 8). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 5 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (8 x 1000 mM) eluting with CH₂Cl₂/MeOH (20:1) to afford 68 mg (52%) of white solid. mp 220-221 °C; M+H = 344.


15

EXAMPLE 324:



Utilizing the method described in Example 323, except employing benzoyl chloride, the title compound was prepared in 61% yield as a white solid. mp 172-175 °C; M+H = 406.

EXAMPLE 325:




10

To a solution of ketone (100 mg, 0.29 mmol) from Example 323 in CH₂Cl₂ (2.5 mL) at 0 °C was added MeMgBr (0.35 mL, 3.0M in Et₂O) dropwise. The resulting mixture was stirred for 18h at rt and was carefully quenched by addition of sat. aq. NH₄Cl (2 mL) and CH₂Cl₂ (2 mL) were added. The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 4 mL). The organic layers were combined, dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (8 x 1000 μM) eluting with CH₂Cl₂/MeOH (10:1) to afford 68 mg (52%) of a yellow solid. mp 160-162 °C; M+H = 360.

20


EXAMPLE 326:

221



To a solution of ketone (84 mg, 0.24 mmol) from Example 323 in
5 MeOH/THF (1:1; 2 mL total) at 0 °C was added NaBH₄ (12 mg, 0.30 mmol) in
one portion. The resulting mixture was stirred for 18h at rt whereupon and
additional portion of NaBH₄ (12 mg, 0.30 mmol) was added. The mixture was
stirred for 12h whereupon the mixture was quenched with ice followed by
addition of 1M NaOH to adjust the pH = 9. The mixture was diluted with CH₂Cl₂
10 (5 mL). The layers were separated and the aqueous layer was extracted with
CH₂Cl₂ (2 x 4 mL). The organic layers were combined, dried (Na₂SO₄), filtered,
and concentrated under reduced pressure. The crude product was purified by
preparative TLC (8 x 1000 μM) eluting with CH₂Cl₂/MeOH (10:1) to afford 25 mg
(30%) of a yellow solid. mp 148-150 °C; M+H = 346.


15 **EXAMPLE 327:**



Using the same procedure as outlined in Example 326, the ketone (84
20 mg, 0.21 mmol) was converted to 53 mg (62%) as a light yellow solid. mp 78-80
°C; M+H = 408.


EXAMPLE 328:

222



To a solution of 3-H adduct (1.3 g, 4.31 mmol) from Preparative Example 187.10 in CH_2Cl_2 (50 mL) was added Eschenmoser's salt (0.79 g, 4.31 mmol) followed by dropwise addition of TFA (0.56 mL, 7.33 mmol). The mixture was stirred at rt for 48 h and was diluted with CH_2Cl_2 (250 mL). The organic layer was washed with sat. aq. NaHCO_3 (2 x 125 mL) to afford 1.41 g (92%) of a yellow solid. mp 231-233 °C; M+H = 359.


10 **EXAMPLE 329:**




To a solution of tertiary amine adduct (100 mg, 0.28 mmol) from Example 328 in 50% aq. DMF (5 mL) in a pressure tube was added KCN (0.15 g, 2.32 mmol). The tube was capped and heated at 100 °C for 96h. The mixture was cooled to rt and was diluted with EtOAc (25 mL). The organic layer was washed with brine (1 x 5 mL) and water (1 x 5 mL). The organic layers was dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (4 x 1000 μM) eluting with EtOAc to afford 21 mg (30%) of brown solid. mp 152-155 °C; M+H = 341.

EXAMPLE 330:

223



- 5 To a solution of alcohol (45 mg, 0.14 mmol) from Example 17.10 in CH_2Cl_2 (0.7 mL) at 0 °C was added Et_3SiH (26 μL , 0.16 mmol) followed by TFA (25 μL , 0.33 mmol). The mixture was stirred for 2 h at 0 °C and rt for 2 h whereupon additional portions of Et_3SiH (26 μL , 0.16 mmol) and TFA (25 μL , 0.33 mmol) were added and the mixture was stirred for 4 h at rt (until complete by TLC). The mixture was concentrated under reduced pressure and the crude residue was partitioned between CH_2Cl_2 (3 mL) and sat. aq. NaHCO_3 (1.5 mL). The layers were separated and the aqueous layer was extracted with CH_2Cl_2 (2 x 4 mL). The organic layers were combined, washed with brine (1 x 5 mL), dried (Na_2SO_4), filtered and concentrated under reduced pressure. The crude product was purified by prep TLC (4 x 1000 mM) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (20:1) to afford 21 mg (48%) of a yellow solid. mp 146-148 °C; $\text{M}+\text{H} = 316$.

EXAMPLE 331:


20

To a solution of 3-H adduct (90 mg, 0.30 mmol) from Preparative Example 187.10 in conc. H_2SO_4 (2 mL) at 0 °C was added fuming HNO_3 (30 μL , 0.72 mmol) dropwise. The resulting mixture was stirred for 1 h at 0 °C whereupon ice

(~1g) was added to the mixture. The resulting precipitate was collected and was washed with water (2 x 2 mL) and CH₂Cl₂ (2 x 2 mL). The crude product was dried under high vacuum to afford 67 mg (60%) of the monosulfate salt as a yellow/orange solid. mp 250 °C; M+H (free base) = 392.


5 **EXAMPLE 332:**

Step A:



- 10 To a solution of aldehyde (0.10 g, 0.39 mmol) from Preparative Example 168 in THF (2.5 mL) at 0 °C was added CF₃TMS (64 mL, 0.43 mmol) followed by CsF (10 mg). The resulting mixture was stirred for 2 h at 0 °C and 2h at rt. 1M HCl (5 mL) was added and the mixture was diluted with CH₂Cl₂ (10 mL). The layers were separated, the aqueous layer was extracted with CH₂Cl₂ (2 x 10 mL),
15 and the organic layers were combined. The organic layer was washed with brine (1 x 10 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to afford 127 mg (99%) of a yellow semisolid. M+H =328. The crude product was carried on without further purification.


Step B:



- 20
25 By utilizing the general procedure set forth in Example 1, the 7-Cl adduct (127 mg, 0.39 mmol) from Example 332, Step A was reacted with 3-

225

(aminomethyl)pyridine (73 μ L, 0.43 mmol) to afford 80 mg (51%) of the title compound as a light yellow solid. mp 68-72 $^{\circ}$ C; M+H = 400.

EXAMPLE 333:

5

To a solution of aniline (200 mg, 0.69 mmol) from Preparative Example 174 in THF (6 mL) at rt was added aldehyde (114 mg, 0.83 mmol) from Preparative Example 256 followed by dropwise addition of $\text{Ti}(i\text{-OPr})_4$ (0.82 mL, 2.77 mmol). The mixture was stirred at reflux for 4 h and was cooled to rt.













- 10 NaCNBH_3 (347 mg, 5.53 mmol) was added and the mixture was stirred for 2 h at rt. The mixture was cooled to 0 $^{\circ}$ C, treated with 1M NaOH (4 mL) and brine (1 mL) and stirred for 30 min. The mixture was extracted with CH_2Cl_2 (3 x 10 mL) and the organic layers were combined. The organic layer was washed with brine (1 x 7 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure.
- 15 The crude product was purified by preparative thin-layer chromatography (8 x 1000 μ M plates) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (25:1) to afford 89 mg (31%) of the title compound as a yellow solid. mp 210-213 $^{\circ}$ C ; M+H = 411.

EXAMPLES 334-337:

- By essentially the same procedure set forth in Example 333 only by
 20 utilizing the anilines shown in Column 2 of Table 30 and the aldehydes shown in Column 3 of Table 30, the compounds in Column 4 of Table 30 were prepared:


TABLE 30

Ex.	Column 2 (Aniline)	Column 3 (Aldehyde)	Column 4 (Final Structure)	CMPD 1. mp ($^{\circ}$ C) 2. M+H
-----	-----------------------	------------------------	-------------------------------	---

334				1. mp = 85-87 2. M+H = 425
335				1. mp = 160-162 2. M+H = 451
336				1. mp = 117-119 2. M+H = 382
337				1. mp = 171-175 2. M+H = 400

EXAMPLE 338:

227

**STEP A:**

5 Reaction of aniline (0.20 g, 0.69 mmol) with aldehyde (0.13 g, 0.83 mmol) under the reaction conditions described in Example 333 afforded 70 mg (23%) of thiomethyl derivative as a yellow solid. $M+H = 428$.

STEP B:

10 To a solution of thiomethyl derivative (60 mg, 0.14 mmol) from Example 338, Step A in dioxane (2 mL) was added Boc_2O (61 mg, 0.28 mmol) followed by DMAP (21 mg, 0.17 mmol). The mixture was stirred for 14h at rt and was concentrated under reduced pressure. The crude product was purified by preparative thin-layer chromatography (6 x 1000 μM plates) eluting with hexanes/EtOAc (4:1) to afford 61 mg (83%) of the title compound as a yellow solid. $M+H = 528$.

STEP C:


15 To a solution of thiomethyl derivative from Example 338, Step B (41 mg, 0.078 mmol) in CH_2Cl_2 (2 mL) was added MCPBA (33 mg, 0.19 mmol) in one portion. The resulting mixture was stirred for 3h at rt and the mixture was diluted

with CH_2Cl_2 (5 mL) and sat. aq. NaHCO_3 (2.5 mL). The layers were separated, the aqueous layer was extracted with CH_2Cl_2 (2 x 5 mL), and the organic layers were combined. The organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure to afford 40 mg (92%) of the sulfone adduct as a light yellow solid. $M+H = 560$.

STEP D:

To a flask charged with sulfone from Example 338, Step C (75 mg, 0.13 mmol) and a stir bar was added morpholine (2 ml; 22 mmol). The mixture was heated at reflux for 12h, cooled to rt, and concentrated to dryness under high vacuum. The crude product was purified by preparative thin-layer chromatography (6 x 1000 μM plates) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (40:1) to afford 41 mg (68%) of the title compound as a yellow solid. mp 209-210 $^\circ\text{C}$; $M+H = 466$.


EXAMPLE 339:



The title compound was prepared according to the procedure outlined in Example 338 except using benzyl amine to afford 12 mg (70%) of a white solid. mp 194-196; $M+H = 487$.

EXAMPLE 340:


229

**STEP A:**

To a solution of 5-chloro adduct (0.15 g, 0.34 mmol) in dioxane/DIPEA (2.5mL/1.0mL) at rt was added cyclopentylamine (0.041 μ L, 0.41 mmol) dropwise. The resulting solution was stirred at reflux for 16h, cooled to rt, and concentrated under reduced pressure. The crude material was purified by preparative thin-layer chromatography (8 x 1000 μ M) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (25:1) to afford 148 mg (89%) of a yellow oil. $M+H = 489$.

STEP B: Removal of the t-butoxycarbonyl protecting group with TFA


To a solution of the compound prepared in Example 340, Step A (135 mg, 0.28 mmol) in CH_2Cl_2 (2 mL) at rt was added TFA (0.54 mL, 7.0 mmol) dropwise. The resulting solution was stirred for 18 h at rt and was concentrated under reduced pressure. The crude material was redissolved in CH_2Cl_2 (5 mL) and the organic layer was sequentially washed with sat. aq. NaHCO_3 (2 x 2 mL) and brine (1 x 2 mL). The organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The crude material was purified by preparative thin-layer chromatography (8 x 1000 μ M) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (20:1) to afford 105 mg (97%) of white solid. mp 120-122 $^\circ\text{C}$; $M+H = 389$.

EXAMPLE 341:**Step A:**

230

By essentially the same procedure set forth in Example 340 only substituting the appropriate amine, the above compound was prepared. MS: $MH^+ = 431$.

Step B: Removal to t-butoxycarbonyl protecting group with KOH.



5

To a mixture of the compound prepared in Example 341, Step A (0.14 g, 0.26 mmol) in EtOH : H₂O (3 mL, 2 : 1) was added KOH (0.29 g, 20 eq.) in one portion. The resulting solution was stirred at reflux 14 hours, cooled to room temperature, and concentrated under reduced pressure. The residue was taken up in CH₂Cl₂ (5 mL) and diluted with saturated NaHCO₃ (2 mL). The layers were separated and the aqueous layer extracted with CH₂Cl₂ (2 x 4 mL). The combined organics were washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude product was purified by preparative TLC (8 x 1000 μ M) eluting with 5% MeOH in CH₂Cl₂ solution (0.066 g, 59% yield). MS: $MH^+ = 432$; mp = 219-221°C.

10

15









EXAMPLES 342-397:

By essentially the same procedure set forth in Example 340 only substituting the chlorides in Column 2 of Table 31 and removing the t-butoxycarbonyl protecting group by the method shown in Column 3 of Table 31, the compounds shown in Column 4 of Table 31 were prepared.









20

25









Table 31

Ex.	Column 2	Column 3	Column 4	CMPD
342		HCl		MS: MH ⁺ = 403 m.p. 151 – 157 °C
343		HCl		MS: MH ⁺ = 466 m.p. 212 – 217 °C
344		HCl		MS: MH ⁺ = 405 m.p. 53 – 58 °C
345		HCl		MS: MH ⁺ = 405 m.p. 63 – 69 °C











232

346		HCl		MS: MH^+ = 363 m.p. 170 – 171 °C
347		HCl		MS: MH^+ = 407 m.p. 148 – 151 °C
348		HCl		MS: MH^+ = 435 m.p. 56 – 59 °C
349		HCl		MS: MH^+ = 445 m.p. 66 – 68 °C











233

350		KOH		MS: MH^+ = 417 m.p. 149 – 151 °C
351		KOH		MS: MH^+ = 431 m.p. 111 – 114 °C
352		KOH		MS: MH^+ = 417 m.p. 53 – 58 °C
353		KOH		MS: MH^+ = 456 m.p. 186 – 189 °C











234

354		KOH		MS: MH^+ = 416 m.p. 210 – 213 °C
355		TFA		1. mp = 68-70 2. M+H = 494
356		KOH		1. mp = 181-183 2. M+H = 404
357		TFA		1. mp = 69-71 2. M+H = 494
358		KOH		1. mp = 182-184 2. M+H = 404











235

359		KOH		1. mp = 202-204 2. M+H = 418
360		TFA		1. mp = 160-162 2. M+H = 402
361		TFA		1. mp = 151-153 2. M+H = 416
362		KOH		1. mp = 140-143 2. M+H = 418
363		KOH		1. mp = 139-142 2. M+H = 418











236

364	 (+/-)	KOH		1. mp = 115-117 2. M+H = 418
366	 (+/-)	TFA		1. mp = 102-104 2. M+H = 445
367	 (+/-)	TFA		1. mp = 118-120 2. M+H = 474
368	 (+/-)	TFA		1. mp = 106-108 2. M+H = 474
369	 (+/-)	TFA		1. mp = 160-161 2. M+H = 464











237

370	 (+/-)	TFA		1. mp = 93-95 2. M+H = 432
371	 (+/-)	KOH		1. mp = 108-110 2. M+H = 432
372		KOH		1. mp = 180-182 2. M+H = 418
373		TFA		1. mp = 169-170 2. M+H = 417
374		TFA		1. mp = 77-79 2. M+H = 479











238

375		TFA		1. mp = 76-79 2. M+H = 479
376		TFA		1. mp = 105-107 2. M+H = 389
377		TFA		1. mp = 105-107 2. M+H = 389
378		TFA		1. mp = 130-133 2. M+H = 389
379		TFA		1. mp = 132-135 2. M+H = 431










239

380		TFA		1. mp = 135-137 2. M+H = 372
381		KOH		1. mp = 78-82 2. M+H = 432
382		TFA		1. mp = 101-103 2. M+H = 432
383		TFA		1. mp = 92-95 2. M+H = 472
384		TFA		1. mp = 107-111 2. M+H = 444










240

384. 10		TFA		1. mp = 2. M+H = 417
384. 11		TFA		1. mp = 210-212 2. M+H = 391
385		TFA		1. mp = 122-124 2. M+H = 403
386		TFA		1. mp = 186-188 2. M+H = 491
387		TFA		1. mp = 173-175 2. M+H = 483

241

388		TFA		1. mp = 167-169 2. M+H = 450
389		TFA		1. mp = 90-92 2. M+H = 374
390		TFA		1. mp = 113-115 2. M+H = 404
391		TFA		1. mp = 114-116 2. M+H = 404
392	HNMe ₂	TFA		LCMS: MH ⁺ = 347;

242

393	<chem>H2NMe</chem>	TFA		LCMS: $MH^+ = 333$;
394		TFA		LCMS: $MH^+ = 359$;
395		TFA		LCMS: $MH^+ = 405$;
396		TFA		LCMS: $MH^+ = 405$;
397		TFA		LCMS: $MH^+ = 391$;

Additional data for select example shown below:

Example 392: ^1H NMR ($\text{DMSO}-d_6$) δ 8.65 (s, 1H), 8.46 (d, $J = 3.3$ Hz, 1H), 8.21 (t, $J = 6.6$ Hz, 1H), 7.90 (s, 1H), 7.80 (d, $J = 7.8$ Hz, 1H), 7.35 (dd, $J = 7.8, 4.8$ Hz, 1H), 5.46 (s, 1H), 4.61 (d, $J = 6.9$ Hz, 2H), 3.01 (s, 6H).

5 **Example 393:** ^1H NMR (CDCl_3) δ 8.65 (s, 1H), 8.60 (d, $J = 4.8$ Hz, 1H), 7.76 (s, 1H), 7.70 (m, 1H), 7.32 (dd, $J = 8.1, 4.8$ Hz, 1H), 6.43 (t, $J = 6.0$ Hz, 1H), 5.08 (s, 1H), 4.80 (m, 1H), 4.56 (d, $J = 6.0$ Hz, 2H), 2.96 (d, $J = 5.1$ Hz, 3H).

Example 394: ^1H NMR (CDCl_3) δ 8.68 (s, 1H), 8.60 (d, $J = 4.8$ Hz, 1H), 7.76 (s, 1H), 7.72 (m, 1H), 7.32 (dd, $J = 7.8, 5.4$ Hz, 1H), 6.55 (t, $J = 5.7$ Hz, 1H), 5.53 (s, 1H), 5.35 (s, 1H), 4.62 (d, $J = 5.7$ Hz, 2H), 2.49 (m, 1H), 0.75 (m, 2H), 0.51 (m, 2H).

10 **Example 395:** ^1H NMR (CDCl_3) δ 8.65 (s, 1H), 8.60 (d, $J = 4.0$ Hz, 1H), 7.75 (s, 1H), 7.69 (m, 1H), 7.33 (dd, $J = 8.1, 5.1$ Hz, 1H), 6.45 (t, $J = 6.0$ Hz, 1H), 5.07 (s, 1H), 4.69 (m, 1H), 4.54 (d, $J = 6.0$ Hz, 2H), 3.98 (m, 1H), 3.79 (dd, $J = 10.8, 2.4$ Hz, 1H), 3.59 (dd, $J = 11.1, 7.2$ Hz, 1H), 1.59-1.36 (m, 4H), 0.94 (t, $J = 6.9$ Hz, 3H).

Example 396: ^1H NMR (CDCl_3) δ 8.60 (s, 1H), 8.56 (d, $J = 4.2$ Hz, 1H), 7.73 (s, 1H), 7.66 (m, 1H), 7.31 (dd, $J = 7.8, 4.8$ Hz, 1H), 6.51 (t, $J = 6.0$ Hz, 1H), 5.05 (s, 1H), 4.86 (d, $J = 6.6$ Hz, 1H), 4.50 (d, $J = 6.0$ Hz, 2H), 3.94 (m, 1H), 3.78 (dd, $J = 11.1, 2.4$ Hz, 1H), 3.57 (dd, $J = 11.1, 7.2$ Hz, 1H), 1.57-1.34 (m, 4H), 0.91 (t, $J = 7.2$ Hz, 3H).

20 **Example 397:** ^1H NMR (CDCl_3) δ 8.65 (s, 1H), 8.59 (d, $J = 4.5$ Hz, 1H), 7.75 (s, 1H), 7.69 (m, 1H), 7.31 (m, 1H), 6.43 (t, $J = 6.0$ Hz, 1H), 5.06 (s, 1H), 4.88 (m, 1H), 4.55 (d, $J = 6.0$ Hz, 2H), 3.70 (m, 2H), 3.38 (m, 2H), 1.79-1.61 (m, 4H).

25









EXAMPLES 398-416:

By essentially the same conditions set forth in Example 341, Steps A and B only substituting the compound prepared in Preparative Example 193.10, the compounds in Column 4 of Table 32 were prepared.









30

244











Table 32

Ex.	Column 2	Column 3	Column 4	CMPD
398				MS: $MH^+ = 419$ m.p. 102 – 105 °C
399				MS: $MH^+ = 421$ m.p. 79 – 81 °C
400				MS: $MH^+ = 421$ m.p. 78 – 79 °C
401				MS: $MH^+ = 433$ m.p. 228-231 °C






245

402				MS: $MH^+ = 447$ m.p. 97-102 °C
403				MS: $MH^+ = 421$ m.p. °C
404				MS: $MH^+ = 421$ m.p. °C
405				MS: $MH^+ = 386$ m.p. °C

246

407		KOH		1. mp = 98-100 2. M+H = 390
408		TFA		1. mp = 170-173 2. M+H = 404
409		KOH		1. mp = 219-221 2. M+H = 420
410		KOH		1. mp = 110-112 2. M+H = 448
411		TFA		1. mp = 81-83 2. M+H = 448

247

412		TFA		1. mp = 136-138 2. M+H = 448
413	NaOMe	KOH		1. mp = 107-110 2. M+H = 351
414				LCMS: MH ⁺ = 375;

Additional data for select examples shown below:

Example 414: ¹H NMR (DMSO-*d*₆) δ 8.26 (s, 1H), 8.23 (m, 1H), 8.13 (m, 1H), 7.90 (s, 1H), 7.40-7.27 (m, 3H), 5.34 (s, 1H), 4.49 (d, J = 6.3 Hz, 2H), 2.56 (m, 1H), 0.67 (m, 2H), 0.35 (m, 2H).




Example 403: ¹H NMR (DMSO-*d*₆+CDCl₃) δ 8.08 (s, 1H), 7.90 (d, J = 6.3 Hz, 1H), 7.49 (s, 1H), 7.34 (t, J = 6.3 Hz, 1H), 7.16-7.09 (m, 2H), 5.65 (d, J = 6.6 Hz, 1H), 4.97 (s, 1H), 4.90 (s, 1H), 4.29 (d, J = 6.3 Hz, 2H), 3.70 (m, 1H), 3.46 (m, 1H), 3.34 (m, 1H), 1.35-1.17 (m, 4H), 0.71 (t, J = 7.2 Hz, 3H).

Example 404: ¹H NMR (DMSO-*d*₆) δ 8.21 (s, 1H), 8.12 (d, J = 6.6 Hz, 1H), 8.06 (m, 1H), 7.86 (s, 1H), 7.38 (t, J = 7.8 Hz, 1H), 7.30 (d, J = 7.5 Hz, 1H), 6.73 (d, J = 8.7 Hz, 1H), 5.28 (s, 1H), 4.70 (t, J = 5.1 Hz, 1H), 4.41 (d, J = 6.6 Hz, 2H), 4.00 (s, 1H), 3.39 (m, 1H), 1.53 (m, 1H), 1.36-1.25 (m, 3H), 0.86 (t, J = 7.0 Hz, 3H).



EXAMPLES 417—421:


- By the procedure set forth in *Chem. Pharm. Bull.* 1999, 47, 928-938.
utilizing the oxygen or sulfur nucleophiles shown in Column 2 as described of
5 Table 33 and by employing the cleavage method listed in Column 3 of Table 33,
the compounds in Column 4 of Table 33 were prepared:

TABLE 33

Ex.	Column 2 (Nucleophile)	Column 3 (Cleavage method)	Column 4 (Final Structure)	CMPD 1. mp. 2. M+H
417	NaSMe	TFA		1. mp = 172-175 2. M+H = 351
418	NaSt-Bu	TFA		1. mp = 165-168 2. M+H = 392
419	NaSPh	TFA		1. mp = 154-156 2. M+H = 412
420	NaOMe	TFA		1. mp = 161-163 2. M+H = 335

249

				
421	NaOPh	TFA		1. mp = 64-66 2. M+H = 397

EXAMPLE 422:

5





To a solution of amino compound (18 mg, 0.043 mmol) from Example 373 in CH_2Cl_2 (1 mL) at rt was added DIPEA (10 μL , 0.056 mmol) followed by MeSO_2Cl (4 mL, 0.052 mmol). The mixture was stirred at rt for 12 h and was diluted with CH_2Cl_2 (2 mL) and sat. aq. NaHCO_3 (2 mL). The layers were separated and the organic layer was extracted with brine (1 x 2 mL). The organic layer was dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The crude material was purified by preparative thin-layer chromatography (4 x 1000 μM) eluting with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (20:1) to afford 16 mg (75%) of white solid. mp 152-154 $^\circ\text{C}$; M+H = 495.


15 **EXAMPLES 423-424:**

Utilizing the procedure outlined in Example 422, the amino compounds (Column 2) were converted to the corresponding methylsulfonamides (Column 3) in Table 34.

5

TABLE 34

Ex.	Column 2 (Amine)	Column 3 (Final Structure)	CMPD 1. mp. 2. M+H
423			1. mp = 166-168 2. M+H = 467
424			1. mp = 165-168 2. M+H = 467

EXAMPLE 425:**STEP A:**


10

A mixture of the compound prepared in Preparative Example 194 (132 mg, 0.25 mmol), tributylvinyltin (95 mg, 0.30 mmol) and

251

tetrakis(triphenylphosphine) palladium (29 mg, 0.025 mmol) in anhydrous dioxane (5 mL) was refluxed under N₂ for 24 hr. The solvent was evaporated and the residue was purified by flash chromatography using 2:1 CH₂Cl₂:EtOAc as eluent to yield yellow waxy solid (53 mg, 50%). LCMS: MH⁺=428.


5 **STEP B:**



A mixture of the compound prepared in Example 425, Step A (50 mg, 0.12 mmol) and KOH (100 mg, 1.80 mmol) in ethanol (3 mL) and H₂O (0.6 mL) was stirred at 70°C under N₂ for 24 hr. NaHCO₃ (1.0 g), Na₂SO₄ (2.0g), and CH₂Cl₂ (20 mL) were added, the mixture was shaken and then filtered. The solvent was evaporated and the residue was purified by flash chromatography using 20:1:0.1 CH₂Cl₂:MeOH:conc.NH₄OH as eluent to yield yellow waxy solid (17 mg, 45%). LCMS: MH⁺=328. Mp=48-51 °C.

EXAMPLE 426:


STEP A:




By essentially the same procedure set forth in Example 425, Step A only using tributylmethylethynyltin, the compound shown above was prepared.

STEP B:

252




- A mixture of the compound prepared in Example 426, Step A (150 mg, 0.34 mmol) and PtO₂ (30 mg, 0.13 mmol) in glacial acetic acid (5 mL) was stirred under 1 atmosphere of H₂ for 20 hr. The mixture was filtered, fresh PtO₂ (30 mg, 0.13 mmol) was added and the mixture was stirred under 1 atmosphere of H₂ for 2.5 hr. The mixture was poured onto Na₂CO₃ (20 g) and H₂O (200 mL) and it was extracted with CH₂Cl₂ (4x20 mL). The combined extracts were dried over Na₂SO₄ and filtered. The solvent was evaporated and the residue was purified by flash chromatography using 1:1 CH₂Cl₂:EtOAc as eluent to yield yellow waxy solid (68 mg, 45%).

STEP C:


- By essentially the same procedure set forth in Example 425, Step B only substituting the compound prepared in Example 426, Step B, the compound shown above was prepared, MS: MH⁺=344. Mp=110-112 °C.

EXAMPLE 427:**STEP A:**

253




A mixture of the compound prepared in Preparative Example 194 (527 mg, 1.00 mmol), triethyl(trifluoromethyl)silane (666 mg, 3.60 mmol), potassium fluoride (210 mg, 3.60 mmol), and CuI (850 mg, 4.46 mmol) in anhydrous DMF (4 mL) was stirred in a closed pressure vessel at 80°C for 72 hr. CH₂Cl₂ (80 mL) was added and the mixture was filtered through Celite. The solvent was evaporated and the residue was purified by flash chromatography using 2:1 CH₂Cl₂: EtOAc as eluent to yield pale orange waxy solid (70 mg, 15%). LCMS: M⁺=470.

STEP B:


TFA (0.70 mL) was added at 0°C under N₂ to a stirred solution of the compound prepared in Example 427, Step A (70 mg, 0.15 mmol), in anhydrous CH₂Cl₂ (3 mL). The mixture was stirred at 0°C for 10 min, then at 25°C for 2 hr. It was poured into 10 % aqueous Na₂CO₃ (50 mL), extracted with CH₂Cl₂ (3x15 mL), dried over Na₂SO₄, and filtered. The solvent was evaporated and the residue was purified by flash chromatography using EtOAc as eluent to yield off-white solid (40 mg, 73%). LCMS: M⁺=370. Mp=156-158 °C.

EXAMPLE 428:**STEP A:**

254




A mixture of the compound prepared in Preparative Example 193 (100 mg, 0.28 mmol), tetracyclopropylltin (91 mg, 0.32 mmol), Pd₂dba₃ (8.0 mg, 0.009 mmol) and Pd(Pt-Bu₃)₂ (9.0 mg, 0.017 mmol) in anhydrous dioxane (3 mL) was refluxed under N₂ for 27 hr. The solvent was evaporated and the residue was purified by flash chromatography using 1:1 CH₂Cl₂:EtOAc as eluent to yield colorless waxy solid (38 mg, 38%). LCMS: MH⁺=366.

STEP B:

10

A mixture of the compound prepared in Example 428, Step A (36 mg, 0.10 mmol) and KOH (300 mg, 5.40 mmol) in ethanol (3 mL), 1,2-dimethoxyethane (3.0 mL) and H₂O (0.8 mL) was refluxed under N₂ for 4 hr. It was poured into saturated aqueous NaHCO₃ (100 mL), extracted with CH₂Cl₂ (5x10 mL), dried over Na₂SO₄, and filtered. The solvent was evaporated and the residue was purified by flash chromatography using 30:1 EtOAc:MeOH as eluent to yield colorless waxy (18 mg, 69%). LCMS: MH⁺=266.


STEP C:



N-bromosuccinimide (12 mg, 0.068 mmol) in anhydrous CH_3CN (2 mL) was added under N_2 to a stirred solution of the compound prepared in Example 428, Step B (18 mg, 0.068 mmol), in anhydrous CH_3CN (2 mL). The mixture was stirred at 25°C for 2 hr. The solvent was evaporated and the residue was purified by flash chromatography using EtOAc as eluent to yield 5 mg (17%) of the dibromo compound (white solid, LCMS: $\text{MH}^+=370$, mp= $150\text{--}152^\circ\text{C}$) and 8 mg (34%) of the monobromo compound (colorless solid, LCMS: $\text{M}^+=344$, mp= $196\text{--}198^\circ\text{C}$).

EXAMPLE 429:


STEP A:



1,3-propanesultam (72 mg, 0.60 mmol) in anhydrous DMF (3 mL) was added under N_2 to 60 % NaH in mineral oil (36 mg, 0.90 mmol). The mixture was stirred for 20 min, then the compound prepared in Preparative Example 196 (200 mg, 0.46 mmol) was added. The mixture was stirred at 100°C for 30 min, the solvent was evaporated and the residue was purified by flash chromatography using EtOAc as eluent to yield colorless solid (150 mg, 63%). LCMS: $\text{M}^+=523$.


STEP B:

256



TFA (1.5 mL) was added at 0°C under N₂ to a stirred solution of the compound prepared in Preparative Example 196 (140 mg, 0.27 mmol), in anhydrous CH₂Cl₂ (5 mL). The mixture was stirred at 0°C for 10 min, then at 25°C for 2 hr. It was poured onto Na₂CO₃ (10 g), extracted with CH₂Cl₂ (3x50 mL), and filtered. The solvent was evaporated and the residue was purified by flash chromatography using 40:1 EtOAc:MeOH as eluent to yield white solid (32 mg, 28%). LCMS: M⁺=423. Mp=218-220 °C.

EXAMPLE 430:











Where: R₂=H, or Cl

3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (1 equivalent) (prepared as described in Preparative Example 129), or 3-Bromo-7-chloro-5-phenylpyrazolo[1,5-a]pyrimidine (1 equivalent) (prepared as described in Preparative Example 127), R₁NH₂ (1.2 equivalents) and diisopropyl ethylamine (2 equivalents) were dissolved in anhydrous 1,4-dioxane and the mixture was heated at 75°C for the time given in Table 97. The solution was evaporated to dryness and the residue was chromatographed on a silica gel column as described in Table 97, to give the title compound.

Using the appropriate reactants and essentially the same procedure as described above, the products of Examples 431 to 438 were prepared. Variations in the reaction conditions are noted in Table 35.

TABLE 35

Ex.	Structure	MW	FABMS MH ⁺	Reaction Conditions	Yield	Chromatographic Data
431		463.8	463.0	75°C / 26h	52%	15x2.5cm 0.5-2% (10% Conc. ammonium hydroxide in methanol)- dichloromethane
432		429.3	429.2	75°C / 26h 25°C / 39h	53%	15x5cm Dichloromethane; 1.5% (10% Conc. ammonium hydroxide in methanol)- dichloromethane
433		477.8	477.1	75°C / 26h	48%	15x5cm Dichloromethane; 3.5-15% (10% Conc. ammonium hydroxide in methanol)- dichloromethane
434		477.8	477.0	75°C / 26h	50%	15x5cm Dichloromethane; 3.5-15% (10% Conc. ammonium hydroxide in methanol)- dichloromethane
435		434.8	434.1	75°C / 24h 25°C / 65h	53%	15x2.5cm 3% (10% Conc. ammonium hydroxide in methanol)- dichloromethane
436		434.8	434.2	75°C / 27h	31%	15x2.5cm 3% (10% Conc. ammonium hydroxide in methanol)- dichloromethane

						
437		438.7	438.1	75°C / 21h 25°C / 46h	97%	15x2.5cm 0.25% (10% Conc. ammonium hydroxide in methanol)- dichloromethane
438		438.7	438.1	75°C / 28h -20°C / 72h	95%	60x2.5cm 20% Ethyl acetate in hexane

Additional physical data for the compounds are given below:

- 5 **EXAMPLE 431:** Reactants: 3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (110mg, 0.318mmoles) (prepared as described in Preparative Example 129); 3-(aminomethyl)piperidine-1-carboxamide (60mg, 0.382mmoles) (prepared as described in Preparative Example 241 above); diisopropyl ethylamine (0.111mL, 0.636mmoles); anhydrous 1,4-dioxane (2.5mL). Physical
- 10 properties: HRFABMS: m/z 463.0628 (MH⁺). Calcd. for C₁₉H₂₁N₆OBrCl: m/z 463.0649: δ_H (CDCl₃) 1.38 (1H, m, CH₂), 1.52 (1H, m, CH₂), 1.73 (1H, m, CH), 1.93 (1H, m, CH₂), 2.02 (1H, m, CH₂), 2.98 (1H, m, CH₂), 3.06 (1H, m, CH₂), 3.37 (2H, m, CH₂), 3.58 (1H, m, CH₂), 3.82 (1H, m, CH₂), 4.87 (2H, bm, CONH₂), 6.28 (1H, s, H₆), 7.02 (1H, m, NH), 7.36 (2H, m, Ar-H), 7.45 (1H, m, Ar-H), 7.68
- 15 (1H, m, Ar-H) and 8.00 ppm (1H, s, H₂); δ_C (CDCl₃) CH₂: 23.7, 28.1, 44.6, 45.5, 47.2; CH: 35.2, 87.4, 127.2, 130.1, 130.3, 131.6, 143.9; C: 83.1, 132.1, 138.6, 145.5, 146.5, 158.0, 158.4.

EXAMPLE 432: Reactants: 3-Bromo-7-chloro-5-phenylpyrazolo[1,5-a]pyrimidine (500mg, 1.62mmoles) (prepared as described in Preparative Example 127); 3-(aminomethyl)piperidine-1-carboxamide (306mg, 1.944mmoles) (prepared as described in Preparative Example 241 above); diisopropyl ethylamine (0.566mL, 3.24mmoles); anhydrous 1,4-dioxane (13mL). Physical properties: HRFABMS: m/z 429.1031 (MH^+). Calcd. for $C_{19}H_{22}N_6OBr$: m/z 429.1038; δ_H ($CDCl_3$) 1.44 (1H, m, CH_2), 1.59 (1H, m, CH_2), 1.79 (1H, m, CH), 2.01 (1H, m, CH_2), 2.08 (1H, m, CH_2), 3.03 (1H, m, CH_2), 3.13 (1H, m, CH_2), 3.39 (1H, m, CH_2), 3.47 (1H, m, CH_2), 3.63 (1H, m, CH_2), 3.90 (1H, m, CH_2), 4.88 (2H, bm, $CONH_2$), 6.40 (1H, s, H_6), 6.90 (1H, m, NH), 7.53 (2H, m, Ar-H), 8.02 (1H, s, H_2) and 8.12 (1H, m, Ar-H); δ_C ($CDCl_3$) CH_2 : 23.7, 28.2, 44.7, 45.5, 47.3; CH: 35.2, 82.9, 127.5, 127.5, 128.7, 128.7, 130.0, 143.9; C: 83.0, 138.5, 145.8, 147.1, 158.3, 158.5.

EXAMPLE 433: Reactants: 3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (347mg, 1.01mmoles) (prepared as described in Preparative Example 129); 3-(aminoethyl)piperidine-1-carboxamide (208mg, 1.21mmoles) (prepared as described in Preparative Example 242 above); diisopropyl ethylamine (0.393mL, 2.02mmoles); anhydrous 1,4-dioxane (9mL). Physical properties: δ_H ($CDCl_3$) 1.24 (1H, m, CH_2), 1.55 (1H, m, CH), 1.72 (4H, m, CH_2), 1.93 (1H, m, CH_2), 2.69 (1H, m, CH_2), 2.94 (1H, m, CH_2), 3.55 (2H, m, CH_2), 3.73 (1H, m, CH_2), 3.98 (1H, m, CH_2), 4.83 (2H, bm, $CONH_2$), 6.55 (1H, s, H_6), 6.78 (1H, m, NH), 7.41 (2H, m, Ar-H), 7.50 (1H, m, Ar-H), 7.75 (1H, m, Ar-H) and 8.04 ppm (1H, s, H_2); δ_C ($CDCl_3$) CH_2 : 24.6, 30.7, 32.6, 39.9, 45.3, 49.3; CH: 33.3, 87.5, 127.4, 130.1, 130.2, 131.6, 143.8; C: 83.2, 132.1, 138.8, 145.7, 146.2, 158.1, 158.1.

EXAMPLE 434: Reactants: 3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (275mg, 0.803mmoles) (prepared as described in Preparative Example 129); 4-(aminoethyl)piperidine-1-carboxamide (165mg, 0.963mmoles) (prepared as described in Preparative Example 243 above); diisopropyl ethylamine (0.311mL, 0.963mmoles); anhydrous 1,4-dioxane (7.2mL). Physical properties: δ_H (d_6 -DMSO) 1.00 (2H, m, CH_2), 1.50 (1H, m, CH), 1.59 (2H, m,


CH₂), 1.67 (2H, m, CH₂), 2.60 (2H, m, CH₂), 3.48 (2H, m, CH₂), 3.70 (2H, m, CH₂), 5.84 (2H, bs, CONH₂), 6.43 (1H, s, H₆), 7.50 (2H, m, Ar-H), 7.62 (2H, m, Ar-H), 8.30 (1H, s, H₂) and 8.36 ppm (1H, m, NH); δ_c (d₆-DMSO) CH₂: 31.5, 31.5, 34.8, 43.5, 43.5, 43.5; CH: 32.8, 86.8, 127.1, 129.7, 130.3, 131.0, 143.3; CH: 81.3, 131.0, 138.7, 145.1, 146.4, 157.3, 157.8.

EXAMPLE 435: Reactants: 3-Bromo-7-chloro-5-phenylpyrazolo[1,5-a]pyrimidine (174mg, 0.507mmoles) (prepared as described in Preparative Example 129) and 3-(aminomethyl)-1-methylpiperidine (65mg, 0.507mmoles) (prepared as described in Preparative Example 244 above); diisopropyl ethylamine (0.178mL, 1.014mmoles); anhydrous 1,4-dioxane (2.5mL). Physical properties: HRFABMS: m/z 434.0742 (MH⁺). Calcd. for C₁₉H₂₂N₅BrCl: m/z 434.0747; δ_H (CDCl₃) 1.18 (1H, m, CH₂), 1.68 (1H, m, CH₂), 1.80 (1H, m, CH₂), 1.87 (1H, m, CH₂), 1.96 (1H, m, CH), 2.14 (2H, m, CH₂), 2.32 (3H, s, NCH₃), 2.75 (1H, m, CH₂), 2.29 (1H, m, CH₂), 3.42 (2H, m, -NHCH₂CH), 6.36 (1H, s, H₆), 6.64 (1H, bm, NH), 7.41 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.74 (1H, m, Ar-H) and 8.06 ppm (1H, s, H₂); δ_c (CDCl₃) CH₃: 46.6; CH₂: 24.4, 27.9, 46.1, 56.1, 59.6; CH: 36.0, 87.4, 127.1, 130.1, 130.2, 131.6, 143.8; C: 83.2, 132.1, 138.9, 145.6, 146.4, 158.2.

EXAMPLE 436: Reactants: 3-Bromo-7-chloro-5-phenylpyrazolo[1,5-a]pyrimidine (111.4mg, 0.325mmoles) (prepared as described in Preparative Example 129); 4-(aminomethyl)-1-methylpiperidine (50mg, 0.39mmoles) (prepared as described in Preparative Example 245 above); diisopropyl ethylamine (0.1135mL, 0.65mmoles); anhydrous 1,4-dioxane (1.5mL). Physical data: HRFABMS: m/z 434.0735 (MH⁺). Calcd. for C₁₉H₂₂N₅BrCl: m/z 434.0747; δ_H (CDCl₃) 1.42 (2H, m, CH₂), 1.72 (1H, m, CH), 1.82 (2H, m, CH₂), 1.93 (2H, m, CH₂), 2.20 (3H, s, NCH₃), 2.89 (2H, m, CH₂), 3.34 (2H, m, -NHCH₂CH), 6.31 (1H, s, H₆), 6.46 (1H, m, NH), 7.36 (2H, m, Ar-H), 7.46 (1H, m, Ar-H), 7.70 (1H, m, Ar-H) and 8.00 ppm (1H, s, H₂); δ_c (CDCl₃) CH₃: 46.4; CH₂: 30.2, 30.2, 48.0, 55.3, 55.3; CH: 35.4, 87.5, 127.2, 130.2, 130.2, 131.6, 143.8; C: 83.3, 132.2, 138.9, 145.7, 146.4, 158.1.

EXAMPLE 437: Reactants: 3-Bromo-7-chloro-5-phenylpyrazolo[1,5-a]pyrimidine (191mg, 0.557mmoles) (prepared as described in Preparative Example 129); 3-(aminomethyl)benzonitrile (88.3mg, 0.668mmoles) (prepared as described in Preparative Example 246 above); diisopropyl ethylamine (0.192mL, 1.114mmoles); anhydrous 1,4-dioxane (4.5mL). Physical data: HRFABMS: m/z 438.0125 (MH⁺). Calcd. for C₁₉H₁₂N₅BrCl: m/z 438.0121; δ_H (CDCl₃) 4.76 (2H, d, -CH₂NH-), 6.32 (1H, s, H₆), 7.00 (1H, m, -CH₂NH-), 7.40 (2H, m, Ar-H), 7.46 (1H, m, Ar-H), 7.55 (1H, m, Ar-H), 7.67 (2H, m, Ar-H), 7.71 (1H, m, Ar-H), 7.75 (1H, m, Ar-H) and 8.10 ppm (1H, s, H₂); δ_C (CDCl₃) CH₂: 45.5; CH: 88.2, 127.2, 130.0, 130.2, 130.4, 130.6, 131.4, 131.6, 131.9, 144.1; C: 83.8, 113.4, 118.3, 132.0, 137.8, 138.3, 145.6, 145.9, 158.0.

EXAMPLE 438: Reactants: 3-Bromo-7-chloro-5-phenylpyrazolo[1,5-a]pyrimidine (233.5mg, 0.681mmoles) (prepared as described in Preparative Example 129); 4-(aminomethyl)benzonitrile (108mg, 0.817mmoles) (prepared as described in Preparative Example 247 above); diisopropyl ethylamine (0.235mL, 1.362mmoles); anhydrous 1,4-dioxane (5.3mL). Physical data: HRFABMS: m/z 438.0117 (MH⁺) Calcd. for C₂₀H₁₄N₅BrCl: m/z 438.0121; δ_H (CDCl₃) 4.80 (2H, d, CH₂), 6.30 (1H, s, H₆), 7.01 (1H, m, NH), 7.40 (2H, m, Ar-H), 7.47 (1H, m, Ar-H), 7.70 (2H, m, Ar-H), 7.72 (2H, m, Ar-H), 7.80 (1H, m, Ar-H) and 8.10 ppm (1H, s, H₂); δ_C (CDCl₃) CH₂: 45.8; CH: 88.2, 127.2, 127.7, 127.7, 130.2, 130.4, 131.6, 132.9, 132.9, 144.1; C: 83.8, 112.2, 118.4, 132.0, 138.2, 141.5, 145.5, 146.0, 158.0.




EXAMPLE 439:

3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (50mg, 0.146mmoles) (prepared as described in Preparative Example 129) was dissolved in anhydrous 1,4-dioxane (5mL) in a GeneVac Technologies carousel reaction tube. PS-diisopropyl ethylamine resin (161mg, 0.5828mmoles) was






30

added to each tube. A freshly prepared 1M solution of the appropriate amine R_1NH_2 in anhydrous 1,4-dioxane (0.2185mL, 0.2185mmoles) was added to each tube and the tubes were sealed and heated at 70°C for 78h with magnetic stirring in the reaction block. Each tube was filtered and the resin was washed with anhydrous 1,4-dioxane and then dichloromethane. The combined individual filtrates from each tube were evaporated to dryness and the residues were each re-dissolved in anhydrous 1,4-dioxane (5mL) and placed in GeneVac reaction tubes. To each tube was added PS-isocyanate resin (594mg, 0.8742mmoles) and PS-trisamine resin (129mg, 0.4371mmoles) and the tubes were stirred at 25°C for 20h in the reaction block. The resins were filtered off and washed with anhydrous 1,4-dioxane and dichloromethane. The filtrates from each tube were evaporated to dryness and the residues were each chromatographed on a silica gel column using the column size and the eluant shown in Table 36, to give the title compounds.



TABLE 36

Ex.	Structure	MW	FABMS MH ⁺	Yield	Chromatographic Data
440		428.7	428.0	81%	15x2.5cm Dichloromethane; 0.5% Methanol in dichloromethane
441		428.7	428.0	48%	20x2cm Dichloromethane; 1.5% Methanol in dichloromethane
442		428.7	428.0	24%	15x2.5cm Dichloromethane; 1.5% Methanol in dichloromethane

263

443		463.8	463.0	44%	15x2.2cm Dichloromethane; 5% Methanol in dichloromethane
444		434.8	434.1	63%	15x2.5cm 5% Methanol in dichloromethane
445		448.8	448.2	65%	15x2.5cm 5% Methanol in dichloromethane
446		448.8	448.1	40%	15x2.5cm Dichloromethane; 0.5% Methanol in dichloromethane
447		436.7	436.1	72%	15x2.5cm 0.5% Methanol in dichloromethane

264

448		450.8	450.0	53%	20x2cm Dichloromethane; 0.5% Methanol in dichloromethane
449		381.7	381.0	44%	20x2cm 1.5% Methanol in dichloromethane

Additional physical data for the compounds are given below:

EXAMPLE 440: Physical properties: HRFABMS: m/z 428.0272 (MH^+). Calcd. for

5 $C_{19}H_{16}N_5BrCl$: m/z 428.0278; δ_H ($CDCl_3$) 3.28 (2H, dd, $C_5H_4NCH_2CH_2NH-$), 3.94 (2H, ddd, $C_5H_4NCH_2CH_2NH-$), 6.40 (1H, s, H_6), 7.22-7.29 (3H, m, Ar-H), 7.38-7.44 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.68 (1H, ddd, Ar-H), 7.73 (1H, Ar-H), 8.18 (1H, s, H_2) and 8.68ppm (1H, NH); δ_C ($CDCl_3$) CH_2 : 36.4, 41.5; CH: 87.3, 122.1, 123.6, 127.1, 130.1, 130.1, 131.6, 137.0, 143.8, 149.5; C: 83.1, 132.1, 138.9, 145.7, 146.3, 158.0, 158.1.

EXAMPLE 441: Physical properties: HRFABMS: m/z 428.0272 (MH^+). Calcd. for

15 $C_{19}H_{16}N_5BrCl$: m/z 428.0278; δ_H ($CDCl_3$) 3.12 (2H, dd, $C_5H_4NCH_2CH_2NH-$), 3.77 (2H, ddd, $C_5H_4NCH_2CH_2NH-$), 6.40 (1H, s, H_6), 6.59 (1H, m, Ar-H), 7.34 (1H, bm, Ar-H), 7.39-7.45 (2H, m, Ar-H), 7.52 (1H, m, Ar-H), 7.62 (1H, m, Ar-H), 7.75 (1H, m, Ar-H), 8.05 (1H, s, H_2) and 8.63ppm (1H, m, NH); δ_C ($CDCl_3$) CH_2 : 32.7, 43.1; CH: 87.5, 127.2, 130.2, 130.3, 131.6, 136.4, 142.9, 148.3, 149.8; C: 83.5, 132.0, 138.6, 145.6, 145.9, 158.1.

20 **EXAMPLE 442:** Physical properties: HRFABMS: m/z 428.0275 (MH^+). Calcd. for

$C_{19}H_{16}N_5BrCl$: m/z 428.0278; δ_H ($CDCl_3$) 3.13 (2H, dd, $C_5H_4NCH_2CH_2NH-$), 3.80 (2H, ddd, $C_5H_4NCH_2CH_2NH-$), 6.42 (1H, s, H_6), 6.53 (1H, m, Ar-H), 7.23 (2H, m, Ar-H), 7.40-7.46 (2H, m, Ar-H), 7.62 (1H, m, Ar-H), 7.76 (1H, m, Ar-H), 8.07 (1H,

s, H₂) and 8.63ppm (1H, m, NH); δ_c (CDCl₃) CH₂: 34.7, 42.5; CH: 87.4, 124.5, 124.5, 127.2, 130.2, 130.3, 131.6, 144.0, 150.2, 150.2; C: 83.5, 132.0, 138.6, 145.6, 145.9, 146.6, 158.1.

- 5 **EXAMPLE 443:** Physical properties: HRFABMS: m/z 463.1003 (MH⁺). Calcd. for C₂₀H₂₅N₆BrCl: m/z 463.1013; δ_H (CDCl₃) 1.98 (2H, m, =NCH₂CH₂CH₂NH-), 2.43 (3H, s, NCH₃), 2.67 (2H, m, =NCH₂CH₂CH₂NH-), 2.70 (8H, piperazine CH₂), 3.58 (2H, m, =NCH₂CH₂CH₂NH-), 6.32 (1H, s, H₆), 7.37-7.43 (2H, m, Ar-H), 7.50 (1H, m, Ar-H), 7.73 (1H, m, Ar-H), 8.06 (1H, s, H₂) and 8.60ppm (1H, m, NH); δ_c
10 (CDCl₃) CH₃: 46.1; CH₂: 24.1, 42.8, 53.3, 54.6, 54.6, 57.5, 57.5; CH: 87.1, 127.0, 130.0, 130.1, 131.5, 143.4; C: 82.7, 132.1, 139.2, 145.7, 146.7, 158.0.

- EXAMPLE 444:** Physical properties: HRFABMS: m/z 434.0742 (MH⁺). Calcd. for C₁₉H₂₂N₅BrCl: m/z 434.0747; δ_H (CDCl₃) 1.72 (1H, m, CH/CH₂), 1.78-1.90 (2H, m, CH/CH₂), 2.02 (3H, m, CH/CH₂), 2.50 (1H, m, CH/CH₂), 2.45 (3H, s, NCH₃), 2.51 (1H, m, CH/CH₂), 3.23 (1H, m, CH/CH₂), 3.54 (1H, m, CH/CH₂), 3.60 (1H, m, CH/CH₂), 6.32 (1H, s, H₆), 7.38-7.44 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.75 (1H, m, Ar-H), 7.96 (1H, bm, NH) and 8.05 ppm (1H, s, H₂); δ_c (CDCl₃) CH₃:
15 40.7; CH₂: 22.7, 29.3, 30.1, 39.4, 57.0; CH: 64.2, 87.1, 127.1, 130.0, 130.1,
20 131.6, 143.8; C: 82.8, 132.1, 139.1, 145.7, 146.4, 158.0.

- EXAMPLE 445:** Physical properties: HRFABMS: m/z 448.0910 (MH⁺). Calcd. for C₂₀H₂₄N₅BrCl: m/z 448.0904; δ_H (CDCl₃) 1.90 (4H, m, CH₂), 2.00 (4H, m, CH₂), 2.84 (2H, m, CH₂), 2.95 (4H, m, CH₂), 3.51 (2H, m, CH₂), 6.32 (1H, s, H₆), 7.05
25 (1H, bm, NH), 7.37-7.43 (2H, m, Ar-H), 7.50 (1H, m, Ar-H), 7.73 (1H, m, Ar-H) and 8.04 ppm (1H, s, H₂); δ_c (CDCl₃) CH₂: 23.4, 23.4, 24.8, 26.4, 41.8, 53.9, 53.9, 55.2; CH: 87.3, 127.1, 130.1, 130.2, 131.6, 143.7; C: 83.0, 132.0, 138.9, 145.7, 146.3, 158.1.

- 30 **EXAMPLE 446:** Physical properties: HRFABMS: m/z 448.0548 (MH⁺). Calcd. for C₁₉H₂₀N₅OBrCl: m/z 448.0540; δ_H (CDCl₃) 1.94 (2H, m, CH₂), 2.09 (2H, m, CH₂), 2.49 (2H, m, CH₂), 3.45 (2H, m, CH₂), 3.51 (4H, m, CH₂), 6.32 (1H, s, H₆), 7.37-7.44 (3H, m, Ar-H/NH), 7.51 (1H, m, Ar-H), 7.75 (1H, m, Ar-H) and 8.10 ppm

(1H, s, H₂); δ_c (CDCl₃) CH₂: 18.0, 26.3, 30.8, 39.2, 39.9, 47.5; CH: 87.0, 127.1, 130.1, 130.1, 131.6, 144.1; C: 82.9, 132.1, 138.9, 145.6, 146.2, 157.9, 176.2.

EXAMPLE 447: Physical properties: HRFABMS: m/z 436.0532 (MH⁺). Calcd. for

5 C₁₈H₂₀N₅OBrCl: m/z 436.0540; δ_H (CDCl₃) 2.60 (4H, bm, -N(CH₂CH₂)₂O), 2.83 (2H, m, =NCH₂CH₂NH-), 3.57 (2H, m, =NCH₂CH₂NH-), 3.83 (4H, m, -N(CH₂CH₂)₂O), 6.37 (1H, s, H₆), 6.99 (1H, bm, NH), 7.38-7.45 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.75 (1H, m, Ar-H) and 8.09 ppm (1H, s, H₂); δ_c (CDCl₃) CH₂: 38.2, 53.3, 53.3, 56.2, 66.9, 66.9; CH: 87.6, 127.1, 130.1, 130.2, 131.6, 143.9; C: 83.1, 132.1, 138.9, 145.7, 146.2, 158.1.

EXAMPLE 448: Physical properties: HRFABMS: m/z 450.0688 (MH⁺). Calcd. for


C₁₉H₂₂N₅OBrCl: m/z 450.0696; δ_H (CDCl₃) 1.98 (2H, m, =NCH₂CH₂CH₂NH-), 2.58 (4H, m, -N(CH₂CH₂)₂O), 2.67 (2H, m, =NCH₂CH₂CH₂NH-), 3.59 (2H, m, =NCH₂CH₂CH₂NH-), 3.94 (4H, m, -N(CH₂CH₂)₂O), 6.31 (1H, s, H₆), 7.37-7.44 (2H, Ar-H), 7.51 (1H, m, Ar-H), 7.78 (1H, m, Ar-H), 8.08 (1H, s, H₂) and 8.60 ppm (1H, bm, NH); δ_c (CDCl₃) CH₂: 23.7, 42.7, 52.9, 52.9, 58.0, 66.6, 66.6; CH: 87.0, 127.1, 130.0, 130.1, 131.5, 143.6; C: 82.8, 132.1, 139.1, 145.7, 146.7, 158.0.

EXAMPLE 449: Physical properties: HRFABMS: m/z 381.0114 (MH⁺). Calcd.

for C₁₅H₁₅N₄OBrCl: m/z 381.0118; δ_H (CDCl₃) 1.39 (3H, d, CHCH₃), 2.76 (1H, bm, -OH), 3.71 (1H, m, =CHCH₂OH), 3.81 (1H, m, =CHCH₂OH), 3.88 (1H, m, =CHCH₂OH), 6.38 (1H, s, H₆), 7.38 (2H, m, Ar-H), 7.48 (1H, m, Ar-H), 7.68 (1H, m, Ar-H) and 8.02 ppm (1H, s, H₂); δ_c (CDCl₃) CH₃: 16.9; CH₂: 65.0; CH: 50.0, 88.0, 127.1, 130.1, 130.3, 131.4, 143.8; C: 83.0, 132.0, 138.5, 145.6, 146.0, 158.2.

EXAMPLE 450:

267









3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (50mg, 0.146mmoles) (prepared as described in Preparative Example 129) was dissolved in anhydrous 1,4-dioxane (5mL) in a GeneVac Technologies carousel reaction tube. PS-diisopropyl ethylamine resin (161mg, 0.5828mmoles) was added to each tube. A freshly prepared solution of the appropriate amine R_1NH_2 (0.219mmoles) in anhydrous 1,4-dioxane (0.3mL) was added to each tube, with the exception of Example 99-5 in which the amine was dissolved in 10% MeOH in 1,4-dioxane (0.3mL), and the tubes were sealed and heated at 70°C for 74h with magnetic stirring in the reaction block. Each tube was filtered and the resin was washed with anhydrous 1,4-dioxane and then dichloromethane. The combined individual filtrates from each tube were evaporated to dryness and the residues were each re-dissolved in anhydrous 1,4-dioxane (5mL) and placed in GeneVac reaction tubes. To each tube was added PS-isocyanate resin (594mg, 0.8742mmoles) and PS-trisamine resin (129mg, 0.4371mmoles) and the tubes were stirred at 25°C for 20h in the reaction block. The resins were filtered off and washed with anhydrous 1,4-dioxane and dichloromethane. The filtrates from each tube were evaporated to dryness and the residues were each chromatographed on a silica gel column using the column size and the eluant shown in Table 37, to give the title compounds.



TABLE 37

Ex.	Structure	MW	FABMS MH ⁺	Yield	Chromatographic Data
451		381.7	380.9	66%	15x2.5cm; 0.5% Methanol in dichloromethane

268

452		381.7	380.9	60%	20x2cm; 0.5% Methanol in dichloromethane
453		381.7	380.9	69%	15x2.5cm; 0.35% Methanol in dichloromethane
454		381.7	380.9	75%	15x2.5cm; 0.35% Methanol in dichloromethane
455		397.7	397.2	84%	15x2.5cm; 1.5% Methanol in dichloromethane
456		397.7			
457		395.7	395.0	60%	15x2.5cm; 0.35% Methanol in dichloromethane

269

458		395.7	396.3	50%	15x2.5cm; 0.35% Methanol in dichloromethane
459		395.7	396.0	76%	15x2.5cm; 0.35% Methanol in dichloromethane

Additional physical data for the compounds are given below:

EXAMPLE 451: Physical properties: HRFABMS: m/z 381.0115 (MH^+). Calcd. for $C_{15}H_{15}N_4OBrCl$: m/z 381.0118; $[\alpha]_D^{25^\circ C} +1.4^\circ$ ($c=0.25$, MeOH); δ_H ($CDCl_3$) 1.44 (3H, d, $-CHCH_3$), 3.77 3.89 (1H, dd, $CHCH_2OH$), (1H, dd, $CHCH_2OH$), 3.94 (1H, m, $CHCH_2OH$), 6.41 (1H, s, H_6), 6.58 (1H, d, NH), 7.41 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.74 (1H, m, Ar-H) and 8.04 ppm (1H, s, H_2); δ_C ($CDCl_3$) CH_3 : 17.1; CH_2 : 65.5; CH: 49.9, 88.0, 127.1, 130.1, 130.2, 131.6, 143.8; C: 83.2, 132.1, 138.7, 145.6, 145.8, 158.1.

EXAMPLE 452: Physical properties: HRFABMS: m/z 381.0115 (MH^+). Calcd. for $C_{15}H_{15}N_4OBrCl$: m/z 381.0118; $[\alpha]_D^{25^\circ C} +6.5^\circ$ ($c=0.32$, MeOH); δ_H ($CDCl_3$) 1.44 (3H, d, $-CHCH_3$), 3.78 (1H, dd, $CHCH_2OH$), 3.89 (1H, dd, $CHCH_2OH$), 3.96 (1H, m, $CHCH_2OH$), 6.41 (1H, s, H_6), 6.58 (1H, d, NH), 7.41 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.75 (1H, m, Ar-H) and 8.04 ppm (1H, s, H_2); δ_C ($CDCl_3$) CH_3 : 17.1; CH_2 : 65.5; CH: 49.9, 88.0, 127.1, 130.1, 130.3, 131.6, 143.8; C: 83.2, 132.1, 138.6, 145.6, 145.8, 158.1.

EXAMPLE 453: Physical properties: HRFABMS: m/z 381.0115 (MH^+). Calcd. for $C_{15}H_{15}N_4OBrCl$: m/z 381.0118; $[\alpha]_D^{25^\circ C} +9.4^\circ$ ($c=0.27$, MeOH); δ_H ($CDCl_3$) 1.33 (3H, d, CH_3), 2.25 (1H, bs, OH), 3.37 (1H, dd, CH_2), 3.51 (1H, m, CH_2), 4.16 (1H, m, $CHOH$), 6.35 (1H, s, H_6), 6.93 (1H, m, NH), 7.40 (2H, m, Ar-H), 7.50 (1H, m,

Ar-H), 7.70 (1H, m, Ar-H) and 8.04 ppm (1H, s, H₂); δ_C (CDCl₃) CH₃: 20.8; CH₂: 49.2; CH: 65.7, 87.8, 127.1, 130.1, 130.2, 131.2, 143.9; C: 83.1, 132.1, 138.5, 145.6, 146.6, 158.3.

- 5 **EXAMPLE 454:** Physical properties: HRFABMS: m/z 381.0112 (MH⁺). Calcd. for C₁₅H₁₅N₄OBrCl: m/z 381.0118; $[\alpha]_D^{25^\circ C}$ -3.2° (c=0.29, MeOH); δ_H (CDCl₃) 1.32 (3H, d, CH₃), 2.48 (1H, bs, OH), 3.35 (1H, dd, CH₂), 3.49 (1H, m, CH₂), 4.15 (1H, m, CH₂OH), 6.34 (1H, s, H₆), 6.93 (1H, m, NH), 7.39 (2H, m, Ar-H), 7.49 (1H, m, Ar-H), 7.68 (1H, m, Ar-H) and 8.03 ppm (1H, s, H₂); δ_C (CDCl₃) CH₃: 20.8; CH₂: 49.2; CH: 65.7, 87.7, 127.1, 130.1, 130.3, 131.4, 143.9; C: 83.0, 132.0, 138.6, 145.6, 146.6, 158.3.

- 15 **EXAMPLE 455:** Physical properties: HRFABMS: m/z 397.0054 (MH⁺). Calcd. for C₁₅H₁₅N₄O₂BrCl: m/z 397.0067; $[\alpha]_D^{25^\circ C}$ -9.5° (c=0.28, MeOH); δ_H (CDCl₃) 3.18 (2H, bs, OH), 3.47 (1H, dd, CH₂), 3.58 (1H, dd, CH₂), 3.63 (1H, dd, CH₂OH), 3.70 (1H, dd, CH₂OH), 3.98 (1H, m, CH), 6.35 (1H, s, H₆), 7.10 (1H, m, NH), 7.37 (2H, m, Ar-H), 7.46 (1H, m, Ar-H), 7.64 (1H, m, Ar-H) and 8.01 ppm (1H, s, H₂); δ_C (CDCl₃) CH₂: 44.7, 64.0; CH: 69.7, 87.7, 127.0, 130.1, 130.3, 131.3, 143.9; C: 82.9, 132.0, 138.4, 145.4, 146.7, 158.3.

- 20 **EXAMPLE 456:** This enantiomer may be prepared by essentially the same manner as described above.

- 25 **EXAMPLE 457:** Physical properties: HRFABMS: m/z 395.0260 (MH⁺). Calcd. for C₁₆H₁₇N₄OBrCl: m/z 395.0274; $[\alpha]_D^{25^\circ C}$ -34.3° (c=0.28, MeOH); δ_H (CDCl₃) 1.08 (3H, dd, CH₃), 1.78 (1H, m, CH₂), 1.86 (1H, m, CH₂), 2.35 (1H, bs, CH₂OH), 3.71 (1H, m, CH₂NH), 3.81 (1H, dd, CH₂OH), 3.90 (1H, dd, CH₂OH), 6.42 (1H, s, H₆), 6.53 (1H, m, NH), 7.41 (2H, m, Ar-H), 7.51 (1H, Ar-H), 7.75 (1H, m, Ar-H) and 8.04 ppm (1H, s, H₂); δ_C (CDCl₃) CH₃: 10.5; CH₂: 24.5, 63.7; CH: 55.9, 88.0, 127.1, 130.1, 130.2, 131.6, 143.8; C: 83.2, 132.1, 138.6, 145.6, 146.3, 158.1.

EXAMPLE 458: Physical properties: HRFABMS: m/z 395.0274 (MH⁺). Calcd. for C₁₆H₁₇N₄OBrCl: m/z 395.0274; $[\alpha]_D^{25^\circ C}$ +27.5° (c=0.25, MeOH); δ_H (CDCl₃) 1.05


271

(3H, dd, CH₃), 1.76 (1H, m, CH₂), 1.85 (1H, m, CH₂), 2.28 (1H, bs, CH₂OH), 3.67 (1H, m, CHNH), 3.77 (1H, dd, CH₂OH), 3.84 (1H, dd, CH₂OH), 6.49 (1H, s, H₆), 6.66 (1H, m, NH), 7.39 (2H, m, Ar-H), 7.49 (1H, Ar-H), 7.71 (1H, m, Ar-H) and 8.04 ppm (1H, s, H₂); δ_c (CDCl₃) CH₃: 10.5; CH₂: 24.3, 63.3; CH: 56.1, 88.0, 127.1, 130.1, 130.3, 131.5, 143.8; C: 83.0, 132.1, 138.6, 145.6, 146.3, 158.2.

EXAMPLE 459: Physical properties: HRFABMS: m/z 395.0264 (MH⁺). Calcd. for C₁₆H₁₇N₄OBrCl: m/z 395.0274; δ_H (CDCl₃) 1.77 (2H, m, -NHCH₂CH₂CH₂CH₂OH), 1.90 (1H, bm, -NHCH₂CH₂CH₂CH₂OH), 1.93 (2H, m, -NHCH₂CH₂CH₂CH₂OH), 3.54 (2H, m, -NHCH₂CH₂CH₂CH₂OH), 3.77 (2H, m, -NHCH₂CH₂CH₂CH₂OH), 6.37 (1H, s, H₆), 6.72 (1H, m, -NHCH₂CH₂CH₂CH₂OH), 7.41 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.75 (1H, m, Ar-H) and 8.06 ppm (1H, s, H₂); δ_c (CDCl₃) CH₂: 25.7, 29.7, 42.2, 62.2; CH: 87.4, 127.1, 130.1, 130.2, 131.6, 143.8; C: 83.1, 132.1, 138.8, 145.6, 146.3, 158.1.


EXAMPLE 460:

4-[[3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YLAMINO]METHYL]PIPERIDINE-1-CARBOXYLIC ACID AMIDE:



A. 4-[[3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YLAMINO]METHYL]PIPERIDINE-1-CARBOXYLIC ACID *tert*-BUTYL ESTER:


272



3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (300mg, 0.875mmoles) (prepared as described in Preparative Example 129) was dissolved in anhydrous 1,4-dioxane (6.8mL). 4-(aminomethyl)piperidine-1-carboxylic acid *tert*-butyl ester (225mg, 1.05mmoles) and diisopropyl ethylamine (0.3055mL, 1.75mmoles) were added and the mixture was heated at 75°C for 24h. The solution was evaporated to dryness and the residue was chromatographed on a silica gel column (15x5cm) using dichloromethane as the eluant to give 4-[[3-bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-ylamino]methyl]piperidine-1-carboxylic acid *tert*-butyl ester (461.2mg, 100%): FABMS: m/z 520.1 (MH^+); HRFABMS: m/z 520.1111 (MH^+). Calcd. for $C_{23}H_{28}N_5O_2BrCl$: m/z 520.1115; δ_H ($CDCl_3$) 1.30 (2H, m, CH_2), 1.51 (9H, s, -COOC(CH_3)₃), 1.85 (2H, d, CH_2), 1.95 (1H, m, CH), 2.76 (2H, m, CH_2), 3.40 (2H, m, CH_2), 6.37 (1H, s, H_6), 6.55 (1H, m, NH), 7.42 (2H, m, Ar-H), 7.52 (1H, m, Ar-H), 7.76 (1H, m, Ar-H) and 8.07 ppm (1H, s, H_2); δ_C ($CDCl_3$) CH_3 : 28.5, 28.5, 28.5; CH_2 : 29.1, 29.1, 43.5, 43.5, 47.9; CH: 36.3, 87.5, 127.2, 130.2, 130.3, 131.6, 143.9; C: 79.7, 83.3, 132.1, 138.6, 145.4, 146.3, 154.7, 158.1.

B. [3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YL]PIPERIDIN-4-YLMETHYLAMINE:

273



4-[[3-Bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-


ylamino]methyl]piperidine-1-carboxylic acid *tert*-butyl ester (441mg,

5 0.847mmoles) (prepared as described in Example 460, Step A above) was dissolved in methanol (4.5mL) and 10% (v/v) conc. sulfuric acid in 1,4-dioxane (11.46mL) was added. The mixture was stirred at 25°C for 0.5h. The product was worked up as described in Preparative Example 241, step B and chromatographed on a silica gel column (15x5cm) using 8% (10% conc.

10 ammonium hydroxide in methanol)-dichloromethane as the eluant to give [3-bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-yl]piperidin-4-ylmethylamine (314.4mg, 88%): FABMS: m/z 420.0 (MH^+); HRFABMS: m/z 420.0585 (MH^+). Calcd. for $C_{18}H_{20}N_5BrCl$: m/z 420.0591; δ_H ($CDCl_3$) 1.34 (2H, m, CH_2), 1.86 (2H, m, CH_2), 1.91 (1H, m, CH), 2.10 (1H, bm, piperidine-NH), 2.67 (2H, m, CH_2), 3.18 (2H, m, CH_2), 3.38 (2H, m, CH_2), 6.37 (1H, s, H_6), 6.53 (1H, m, NH), 7.42 (2H, m, Ar-H), 7.52 (1H, m, Ar-H), 7.76 (1H, m, Ar-H) and 8.06 ppm (1H, s Ar-H); δ_C ($CDCl_3$) CH_2 : 31.2, 31.2, 46.2, 46.2, 48.4; CH: 36.4, 89.5, 127.1, 130.1, 130.5, 131.6, 143.8; C: 83.2, 132.1, 138.9, 145.6, 146.4, 158.1.

20 C. 4-[[3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YLAMINO]METHYL]PIPERIDINE-1-CARBOXYLIC ACID AMIDE:

274



[3-Bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-yl]piperidin-

4-ylmethylamine (57mg, 0.136mmoles) (prepared as described in Example 460,

5 Step B above) was dissolved in anhydrous dichloromethane (1.2mL) and trimethylsilylisocyanate (0.091mL, 0.679mmoles) was added. The mixture was stirred at 25°C for 2.5h. The mixture was diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate. The organic layer was dried (MgSO₄), filtered and evaporated to dryness. The residue was

10 chromatographed on a silica gel column (30x2.5cm) using 3% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 4-([3-bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-ylamino]methyl)piperidine-1-carboxylic acid amide (53.7mg, 86%): FABMS: m/z 463.1 (MH⁺); HRFABMS: m/z 463.0647 (MH⁺). Calcd. for C₁₉H₂₁N₆OBrCl: m/z 463.0649; δ_H (d₆-DMSO) 1.09 (2H, m, CH₂), 1.63 (2H, m, CH₂), 1.87 (1H, m, CH), 2.60 (2H, m, CH₂), 3.53 (2H, bm, CONH₂), 3.91 (2H, d, CH₂), 6.52 (1H, s, H₆), 7.50 (2H, m, Ar-H), 7.62 (2H, m, Ar-H), 8.33 (1H, s, H₂) and 8.52 ppm (1H, m, NH); δ_C (d₆-DMSO) CH₂: 30.1, 30.1, 44.2, 44.2, 47.7; CH: 36.4, 88.2, 128.1, 130.7, 131.4, 132.1, 147.9; C: 82.1, 132.1, 139.4, 145.7, 147.9, 158.1, 158.8.


20

EXAMPLE 461:

2-{2-[3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YLAMINO]ETHYL}PIPERIDINE-1-CARBOXYLIC ACID AMIDE:


25

275



A. 2-{2-[3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YLAMINO]ETHYL}PIPERIDINE-1-CARBOXYLIC ACID *tert*-BUTYL ESTER:


5



3-Bromo-7-chloro-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidine (400mg, 1.166mmoles) (prepared as described in Preparative Example 129) was dissolved in anhydrous 1,4-dioxane (5.7mL). 2-Aminoethylpiperidine-1-carboxylic acid *tert*-butyl ester (266mg, 1.166mmoles) and diisopropyl ethylamine (0.409mL, 2.33mmoles) were added and the mixture was heated at 75°C for 48h. Additional diisopropyl ethylamine (0.204mL, 1.166mmoles) was added and the heating was continued for a total of 58h. The solution was evaporated to dryness and the residue was chromatographed on a silica gel column (15x5cm) using dichloromethane followed by 0.3% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 2-{[3-bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-ylamino]ethyl}piperidine-1-carboxylic acid *tert*-butyl ester (491.1mg, 79%): FABMS: m/z 534.1 (MH^+); HRESIMS: m/z 534.12797 (MH^+). Calcd. for $C_{24}H_{30}N_5O_2BrCl$: m/z 534.12714; δ_H ($CDCl_3$) 1.50 (1H, m, CH_2), 1.51 (9H, s, $COOC(CH_3)_3$), 1.57 (2H, m, CH_2), 1.68 (2H, m, CH_2), 1.76 (2H, m, CH_2), 2.24 (1H, bm, CH_2), 2.82/3.40/3.54/4.08/4.51 (5H, m, CH/CH_2), 6.34 (1H, s, H_6), 7.41 (2H, m, Ar-H),


7.51 (1H, m, Ar-H), 7.76 (1H, m, Ar-H) and 8.08 ppm (1H, s, H₂); δ_c (CDCl₃) CH₃: 28.5, 28.5, 28.5; CH₂: 19.2, 25.5, 29.2, 29.2, 39.2, 67.1; CH: ~47.4, 87.1, 127.1, 130.1, 130.1, 131.6, 143.9; C: 80.0, 83.0, 132.1, 138.9, 145.7, 146.2, 158.0.

5 B. [3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YL]-(2-PIPERIDIN-2-YLETHYL)AMINE:



- 10 2-[[3-Bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-ylamino]ethyl]piperidine-1-carboxylic acid *tert*-butyl ester (465mg, 0.869mmoles) (prepared as described in Example 461, Step A above) was dissolved in methanol (4.5mL) and 10% (v/v) conc. sulfuric acid in 1,4-dioxane (11.76mL) was added. The mixture was stirred at 25°C for 1.5h. The product was worked up as described in Preparative Example 241, step B and chromatographed on a silica gel column (15x5cm) using 3.5% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give [3-bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-yl]piperidin-2-ylethylamine (365.6mg, 97%): FABMS: *m/z* 434.1 (MH⁺); HRFABMS: *m/z* 434.0726 (MH⁺). Calcd. for C₁₉H₂₂N₅BrCl: *m/z* 434.0747; δ_H (CDCl₃) 1.24 (1H, m, CH₂), 1.41 (1H, m, CH₂), 1.49 (1H, m, CH₂), 1.66 (1H, m, CH₂), 1.73 (1H, m, CH₂), 1.81 (1H, m, CH₂), 1.88 (2H, m, CH₂), 2.68 (1H, m, CH₂), 2.78 (1H, m, CH₂), 3.20 (1H, m, CH), 3.55 (1H, m, CH₂), 3.60 (1H, m, CH₂), 6.32 (1H, s, H₆), 7.41 (2H, m, Ar-H), 7.51 (1H, m, Ar-H), 7.74 (1H, m, Ar-H), 7.78 (1H, m, NH) and 8.05 ppm (1H, s, H₂); δ_c (CDCl₃) CH₂: 24.7, 26.8, 33.1, 35.2, 40.3, 47.0; CH: 55.7, 87.2, 127.1, 130.0, 130.1, 131.5, 143.8; C: 82.9, 132.1, 139.0, 145.7, 146.5, 158.1.
- 15
- 20
- 25

C. 2-{2-[3-BROMO-5-(2-CHLOROPHENYL)PYRAZOLO[1,5-a]PYRIMIDIN-7-YLAMINO]ETHYL}PIPERIDINE-1-CARBOXYLIC ACID AMIDE:




5

[3-Bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-yl]piperidin-2-ylethylamine (200mg, 0.46mmoles) (prepared as described in Example 461, Step B above) was dissolved in anhydrous dichloromethane (2mL) and trimethylsilylisocyanate (0.31mL, 2.3mmoles) was added. The mixture was stirred at 25°C for 1.25h. Additional trimethylsilylisocyanate (0.155mL, 1.15mmoles) was added and the stirring was continued for a total of 3h. The mixture was diluted with dichloromethane and washed with saturated aqueous sodium bicarbonate. The organic layer was dried (MgSO₄), filtered and evaporated to dryness. The residue was chromatographed on a silica gel column (30x2.5cm) using 2% (10% conc. ammonium hydroxide in methanol)-dichloromethane as the eluant to give 2-{2-[3-bromo-5-(2-chlorophenyl)pyrazolo[1,5-a]pyrimidin-7-ylamino]ethyl}piperidine-1-carboxylic acid amide (106.3mg, 48%): FABMS: m/z 477.0 (MH⁺); HRFABMS: m/z 477.0804 (MH⁺). Calcd. for C₂₀H₂₃N₆OBrCl: m/z 477.0805; δ_H (d₆-DMSO) 1.29 (1H, m, CH₂), 1.52 (5H, m, CH₂), 1.72 (1H, m, CH₂), 2.05 (1H, m, CH₂), 2.51 (2H, s, CONH₂), 2.79 (1H, dd, CH), 3.31 (1H, m, CH₂), 3.34 (1H, m, CH₂), 3.76 (1H, m, CH₂), 4.30 (1H, bm, CH₂), 6.42 (1H, s, H₆), 7.50 (2H, m, Ar-H), 7.60 (1H, m, Ar-H), 7.63 (1H, m, Ar-H), 8.29 (1H, s, H₂) and 8.38 ppm (1H, dd, NH); δ_C (d₆-DMSO) CH₂: 18.6, 25.2, 28.2, 38.4, 38.6, 54.8; CH: 46.7, 86.6, 127.1, 129.7, 130.3, 131.0, 143.4; C: 81.2, 131.0, 138.7, 145.1, 146.4, 158.2.

EXAMPLE 462:

278



To a solution of the compound prepared in Example 204 (1.11 g, 2.12 mmol) in anhydrous acetonitrile (20 mL) was added TMSI (1.70 g, 8.52 mmol), dropwise at ambient temperature. After 10 minutes the acetonitrile was removed *in vacuo*. The resulting yellow foam was treated with 2 N HCl solution (7 mL) and then washed immediately with Et₂O (5X). The pH of the aqueous was adjusted to 10 with 50 % NaOH (aq) and the product was isolated by saturation of the solution with NaCl (s) followed by extraction with CH₂Cl₂ (5X) to give the crystalline product (733 mg, 89% yield). MH⁺ = 387; m. p. = 207.5 °C









EXAMPLES 463-472:

By essentially the same procedure set forth in Example 462 only substituting the compounds shown in Column 2 of Table 38, the compounds shown in Column 3 of Table 38 were prepared.









TABLE 38

Ex.	Column 2	Column 3	CMPD
463			MH ⁺ = 403 ¹ H NMR (300MHz, CDCl ₃) δ 8.52 (s, 1H), 8.38 (d, 1H), 8.04 (s, 1H), 7.78 (d, 1H), 7.65 (t, 1H), 6.18 (s, 1H), 4.89 (s, 2H), 3.26 – 3.21 (d, 2H), 2.96 – 2.70 (m, 3H), 2.05 – 1.78 (m, 4H).





279


464			MH ⁺ = 454 m. p. = 175.4 °C
465			Yield = 87 MH ⁺ = 470 m. p. = 220 °C m. pt (hydrochloride salt) = 164.3 °C
466			MH ⁺ = 464 m. p. = 206 °C
467			MH ⁺ = 411 m. p. = 169.5 °C

280

468			$MH^+ = 334$ m. p. = 176.2 °C
469			$MH^+ = 465$ m. p. = 250.4 °C
470			$MH^+ = 387$ m. p. = 68.5 °C
471			$MH^+ = 387$ m. p. = 59.4 °C

281

472			1. mp = 230-232 2. M+H = 396
472 .10			1. mp = 157-160 2. M+H = 427

EXAMPLE 473:Step A:

5


A solution of the sulfonic acid (560 mg, 1.17 mmol) in 5 mL of dry DMF was cooled to 0 °C and SOCl_2 (278 mg, 2.34 mmol) was added. The reaction mixture was brought to RT and stirred overnight. The next day the contents were poured on ice and the pH was carefully adjusted to 8. The product was extracted in to EtOAc and the solvent was removed after drying (Na_2SO_4) to provide 240 mg (41%) of the crude sulfonyl chloride which was used for the next

10

step without further purification. ^1H NMR (CDCl_3) δ 8.20-8.10 (m, 1H), 8.10-7.95 (m, 3H), 7.65 (d, 2H), 7.45-7.35 (m, 1H), 7.35-7.20 (m, 1H), 7.15-7.05 (m, 1H), 6.95 (t, 1H), 4.85 (d, 2H).

Step B:

5




A solution of compound prepared in Example 473, Step A (120 mg, 0.24 mmol) in 10 mL of THF was treated with 2 mL of 1 M MeNH_2 (2.00 mmol) in THF at RT overnight. The solvent was removed and the residue was purified by chromatography (silica, hexane:EtOAc (4:1 \rightarrow 1:1)) to provide 56 mg (48%) of the sulfonamide. ^1H NMR ($\text{DMSO}-d_6$) δ 9.05 (t, J = 9 Hz, 1H), 8.35 (s, 1H), 7.90 (t, J = 7.5 Hz, 1H), 7.75 (d, J = 9 Hz, 2H), 7.62 (d, J = 9 Hz, 2H), 7.55-7.46 (m, 1H), 7.45-7.38 (m, 1H), 7.38-7.25 (m, 1H), 6.50 (s, 1H), 4.80 (d, 2H), 3.30 (s, 3H) LCMS: MH^+ = 492.1

10

15

EXAMPLE 474:



By essentially the same procedure set forth in Example 473, only substituting dimethylamine, the above compound was prepared. ^1H NMR (CDCl_3)


20

δ 8.14 (t, J = 9 Hz, 1H), 8.00 (s, 1H), 7.76 (d, J = 9 Hz, 2H), 7.54 (d, J = 9 Hz, 2H), 7.34-7.44 (m, 1H), 7.26 (t, J = 9 Hz, 1H), 7.14-7.04 (m, 1H), 6.93 (t, J = 6 Hz, 1H), 6.45 (s, 1H), 4.75 (d, 2H), 2.70 (s, 6H)

LCMS: $MH^+ = 504.2$

5

EXAMPLE 475:



A mixture of the compound prepared in Example 129 (300 mg, 0.66 mmol), NaOH (5 g), CH_3OH-H_2O (100 mL, 90:10) was stirred at 25 C for about 15 h. Progress of hydrolysis was checked by TLC. Reaction mixture was concentrated to remove methanol. The concentrate was diluted with 50 mL water, and extracted with ether to remove any un-reacted ester. Aqueous solution, thus obtained, was neutralized with 3 N HCl to pH 4 to obtain free acid, filtered and washed repeatedly with water. The acid was dried under vacuum (270 mg, 93%) and used without further purification.

Example 476-479:

By essentially the same procedure set forth in Example 475 only substituting the compounds in Column 2 of Table 39, the compounds in Column 3 of Table 39 were prepared.

Table 39

Ex.	Column 2	Column 3	CMPD
476			Yield = 82% LCMS: $MH^+ = 365$

477			Yield = 82% LCMS: MH ⁺ = 379
478			Yield = 72% LCMS: MH ⁺ = 393
479			Yield = 70% LCMS: MH ⁺ = 407

+

Additional data for select examples shown below:


Example 476: ¹H NMR (CDCl₃) δ 8.15 (m, 2H), 8.0 (m, 1H), 7.6 (m, 1H), 7.3 (m, 2H), 6.6 (s, 1H), 4.2 (d, 2H).

5 **Example 477:** ¹H NMR (CDCl₃) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 7.0 (t, 1H), 6.5 (s, 1H), 3.8 (dt, 2H), 2.6 (t, 2H).

Example 479: ¹H NMR (CDCl₃) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 3.5 (dt, 2H), 2.4 (t, 2H), 1.8 (m, 4H).

10 **EXAMPLE 480:**

285



A mixture of the acid from Example 475 (85mg, 0.193 mmol) and Et₃N (20 mg, 0.193 mmol) in THF (20 mL) was stirred at 25 C for 15 min. Isobutryl chloroformate (28mg, 0.205 mmol) was added to the reaction mixture and stirred for 10 min followed by addition of NH₄OH solution (0.5 mL). The reaction mixture was stirred for 1 hr and concentrated to dryness. The dry mass was purified by column chromatography.











10 EXAMPLES 481-509:

By essentially the same procedure set forth in Example 480 only substituting the carboxylic acid shown in Column 2 of Table 40 and the amine shown in Column 3 of Table 40, the compounds shown in Column 4 of Table 40 were prepared.













Table 40

Ex.	Column 2	Column 3	Column 4	CMPD
481		CH ₃ NH ₂		Yield = 88% LCMS: MH ⁺ = 454













286













482		$(\text{CH}_3)_2\text{NH}$		Yield=80 % LCMS $\text{MH}^+ = 468$
483		CH_3NH_2		Yield=70 % LCMS $\text{MH}^+ = 454$
484				Yield=75 % LCMS $\text{MH}^+ = 482.1$
485				Yield=71 % LCMS $\text{MH}^+ = 480.1$

287













486				Yield=75 % LCMS MH ⁺ = 494.1
487				Yield=75 % MH ⁺ = 494.1
488				Yield=75 % MH ⁺ = 496.1
489				Yield=75 % LCMS MH ⁺ = 508.1

288
















490				Yield=78 % LCMS MH ⁺ = 524.1
491				Yield=73 % LCMS MH ⁺ = 508.1
492				Yield=73 % LCMS MH ⁺ = 510.1
493				Yield=76 % LCMS MH ⁺ = 526.1

494				Yield=76 % MH ⁺ = 523.1
495				Yield=76 % MH ⁺ = 523.1
496				Yield=51 % LCMS MH ⁺ = 484.1
497				Yield=66 % MH ⁺ = 537.1

290

498				Yield=76 % LCMS MH ⁺ = 551.2
499				Yield=79 % LCMS MH ⁺ = 552.1
500				Yield=80 % MH ⁺ = 549.1
501				Yield=80 % LCMS MH ⁺ = 478.1

291

502				Yield=80 % LCMH ⁺ = 468.1
503				Yield=80 % MH ⁺ = 522.1
504				Yield=82 % LCMS MH ⁺ = 528.1
505				Yield=60 % MH ⁺ = 392
506				Yield=60 % LCMH ⁺ = 448.1

507				Yield=70 % MH ⁺ = 464.1
508				Yield=50 % LCMS MH ⁺ = 436.1
508 .10		CH ₃ NH ₂		Yield = 92 MH ⁺ = 577

Additional data for select examples given below:

Example 481: ¹H NMR (CDCl₃) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (s, 1H), 7.35 (d, 2H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.95 (t, 1H), 6.5 (s, 1H), 6.25 (bs, 1H), 4.7 (d, 2H), 3.0 (d, 3H).

Example 482: ¹H NMR (CDCl₃) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.45 – 7.35 (m, 4H), 7.25 (d, 2H), 7.15 (dd, 1H), 6.7 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.1 (s, 3H), 3.0 (s, 3H).

Example 483: ¹H NMR (CDCl₃) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.8 (bs, 1H), 7.7 (d, 1H), 7.5 – 7.3 (m, 3H), 7.25 (d, 1H), 7.15 (dd, 1H), 6.75 (t, 1H), 6.5 (s, 1H), 6.2 (bs, 1H), 4.7 (d, 2H), 3.0 (d, 3H).

Example 484: ¹H NMR (CDCl₃) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 6.0 (bs, 1H), 4.7 (d, 2H), 4.25 (m, 1H), 1.2 (d, 6H).

Example 485: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.35 (s, 1H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.9 (t, 1H), 6.5 (s, 1H), 6.3 (t, 1H), 4.7 (d, 2H), 2.9 (m, 1H), 0.8 (bt, 2H), 0.6 (bt, 2H).

5 **Example 486:** ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.8 (d, 2H), 7.4 (d, 2H), 7.35 (d, 1H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.9 (t, 1H), 6.5 (s, 1H), 6.2 (t, 1H), 4.7 (d, 2H), 3.3 (dd, 2H), 1.05 (m, 1H), 0.5 (m, 2H), 0.25 (m, 2H).

Example 487: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.85 (t, 1H), 6.5 (s, 1H), 6.2 (bs, 1H), 4.7 (d, 2H), 4.6 (m, 1H), 2.4 (m, 2H), 1.95 (m, 1H), 1.75 (m, 2H).

10 **Example 488:** ^1H NMR (CDCl_3) δ 8.5 (t, 1H), 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 5.9 (bs, 1H), 4.7 (d, 2H), 1.4 (s, 9H).

15 **Example 489:** ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 6.0 (bs, 1H), 4.7 (d, 2H), 4.4 (m, 1H), 2.05 (m, 2H), 1.7 (m, 4H), 1.4 (m, 2H).

Example 490: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.7 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 6.5 (bs, 2H), 4.7 (d, 2H), 4.1 (m, 1H), 3.9 – 3.7 (m, 3H), 3.3 (m, 1H), 2.0 – 1.9 (m, 4H).

20 **Example 491:** ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.45 – 7.35 (m, 5H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.7 (bs, 2H), 3.3 (bs, 2H), 1.7 (bs, 4H), 1.5 (bs, 2H).

Example 492: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.45 – 7.35 (m, 5H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.85 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.8 – 3.4 (bm, 8H).

25 **Example 493:** ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.45 – 7.35 (m, 5H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.80 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 4.0 (m, 2H), 3.6 (m, 2H), 2.8 – 2.45 (m, 4H).

30 **Example 494:** ^1H NMR (CH_3OD) δ 8.15 (s, 1H), 8.0 (dt, 1H), 7.45 – 7.35 (m, 5H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.80 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.7 (bs, 2H), 3.4 (bs, 2H), 2.5 – 2.4 (m, 4H), 2.2 (s, 3H).

Example 495: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.45 – 7.35 (m, 5H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.80 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.75 (bs, 2H), 3.35 (bs, 2H), 2.4 (bs, 2H), 2.3 (s, 3H), 2.2 (bs, 2H).

Example 496: ^1H NMR (CDCl_3) δ 7.95 (s, 1H), 7.9 (dt, 1H), 7.8 (t, 1H), 7.7 (d, 2H), 7.15 (m, 4H), 7.05 (dd, 1H), 6.9 (dd, 1H), 6.2 (s, 1H), 4.5 (d, 2H), 3.6 (t, 2H), 3.3 (dt, 2H).

Example 497: ^1H NMR (CH_3OD) δ 8.1 (s, 1H), 7.9 (dt, 1H), 7.8 (d, 2H), 7.5 (d, 2H), 7.4 (m, 1H), 7.3 (dd, 1H), 7.2 (dd, 1H), 6.4 (s, 1H), 4.7 (d, 2H), 3.5 (t, 2H), 2.7 (m, 2H), 2.6 (bs, 4H), 1.8 (bs, 4H).

Example 498: ^1H NMR (CDCl_3) δ 8.5 (t, 1H), 8.15 (dt, 1H), 8.0 (s, 1H), 7.8 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.7 – 2.5 (m, 4H), 2.35 (s, 3H), 2.2 (m, 1H), 1.9 – 1.6 (m, 6H).

Example 499: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.8 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 4.7 (d, 2H), 3.7 (m, 4H), 3.5 (dt, 2H), 2.6 (t, 2H), 2.5 (m, 4H).

Example 500: ^1H NMR (CH_3OD) δ 8.15 (s, 1H), 7.9 (dt, 1H), 7.8 (d, 2H), 7.45 (d, 2H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.4 (s, 1H), 4.75 (d, 2H), 4.2 (m, 1H), 3.4 – 2.8 (m, 7H), 1.9 – 1.6 (m, 4H).

Example 501: ^1H NMR (CDCl_3) δ 8.05 (dt, 1H), 8.0 (s, 1H), 7.6 (d, 2H), 7.4 (s, 1H), 7.35 (d, 2H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.9 (t, 1H), 6.5 (s, 1H), 6.4 (t, 1H), 4.7 (d, 2H), 4.2 (d, 2H), 2.3 (bs, 1H).

Example 502: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.75 (d, 2H), 7.45 (s, 1H), 7.4 (d, 2H), 7.3 (dd, 1H), 7.1 (dd, 1H), 6.8 (t, 1H), 6.5 (s, 1H), 6.1 (bs, 1H), 4.7 (d, 2H), 3.5 (dq, 2H), 1.2 (t, 3H).

Example 503: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.8 (d, 2H), 7.4 (d, 2H), 7.35 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 6.9 (t, 1H), 6.5 (s, 1H), 6.4 (t, 1H), 4.75 (d, 2H), 4.1 (m, 2H).

Example 504: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.8 (d, 2H), 7.45 (d, 2H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.1 (dd, 1H), 6.8 (t, 1H), 6.6 (t, 1H), 6.5 (s, 1H), 4.7 (d, 1H), 3.6 (m, 2H), 2.8 (t, 2H), 2.6 (q, 2H), 1.3 (t, 3H).


Example 505: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 7.0 (t, 1H), 6.5 (s, 1H), 3.8 (m, 2H), 2.7 (t, 2H), 3.0 (d, 3H).

Example 506: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 7.0 (t, 1H), 6.5 (s, 1H), 3.8 (m, 2H), 3.6 (m, 6H), 3.4 (m, 2H), 2.7 (t, 2H).

Example 507: ^1H NMR (CDCl_3) δ 8.15 (dt, 1H), 8.0 (s, 1H), 7.4 (m, 1H), 7.25 (dd, 1H), 7.15 (dd, 1H), 7.0 (t, 1H), 6.5 (s, 1H), 3.9 (t, 2H), 3.8 (dt, 2H), 3.7 (t, 2H), 2.7 (t, 2H), 2.6 (m, 4H).

Example 508: ^1H NMR (CH_3OD) δ 8.1 (s, 1H), 7.95 (dt, 1H), 7.5 (m, 1H), 7.35 – 7.2 (m, 2H), 6.5 (s, 1H), 3.6 (m, 4H), 3.25 (m, 4H), 2.4 (t, 2H), 2.05 (dt, 2H).

EXAMPLE 509:




A solution of NaOH (59 mg, 1.47 mmol) in 1 mL of water was added to a suspension of $\text{NH}_2\text{OH}\cdot\text{HCl}$ (102 mg, 1.47 mmol) in 10 mL of methanol at 0 °C. After 5 min, the compound prepared in Example 210.10 (208 mg, 0.49 mmol) was added and the reaction mixture was refluxed overnight. The solvent was removed *in vacuo* and the residue was partitioned between water and EtOAc. The EtOAc layer was dried (Na_2SO_4) and the solvent was evaporated. The resulting crude amidoxime was suspended in trimethyl orthoformate containing catalytic amount of PTS acid and refluxed overnight. The solvent was removed and the residue was taken up in EtOAc. The EtOAc layer was washed with aq NaHCO_3 followed by water and brine. The solvent was evaporated and the residue was purified by chromatography (silica, hexane:EtOAc (1:1)) to provide

80 mg (35%) of the oxadiazole. ^1H NMR (CDCl_3) δ 8.75 (s, 1H), 8.20-8.10 (m, 3H), 8.03 (s, 1H), 7.53 (d, $J = 9$ Hz, 2H), 7.45-7.36 (m, 1H), 7.30-7.22 (m, 2H), 7.16-7.08 (m, 1H), 6.80 (t, $J = 5$ Hz, 1H), 6.56 (s, 1H).


LCMS: $\text{MH}^+ = 465.2$

5 **Example 510:**




By essentially the same procedure set forth in Example 509 only substituting the compound prepared in Preparative Example 192, the above compound was prepared. yield = 75; $\text{MH}^+ = 453$; m. p. = 79.3°C.

EXAMPLE 511:




- 15 A mixture of the nitrile (235 mg, 0.56 mmol) and Me_3SnN_3 (343 mg, 1.67 mmol) in 20 mL of dry toluene was refluxed for 2 days under Ar. The solvent was removed *in vacuo* and the residue was dissolved in dry methanol. HCl gas was bubbled through the solution for 15 min and the reaction mixture allowed to stand at overnight at RT. The next day, the solvent was removed, the residue
- 20 was taken in water and the pH was adjusted to 5. The precipitated product was

extracted into EtOAc. Evaporation of the EtOAc layer after drying (Na_2SO_4) provided the residue which was purified by chromatography (silica, DCM:MeOH (98:2→95:5)) to yield 50 mg (19%) of the pure tetrazole. ^1H NMR (CD_3OD) δ 8.10 (s, 1H), 8.00 (d, $J = 9$ Hz, 2H), 7.90 (t, $J = 7$ Hz, 1H), 7.65 (d, $J = 9$ Hz, 2H), 7.50-7.40 (m, 1H), 7.30-7.10 (m, 2H), 6.45 (s, 1H), 4.80 (s, 2H); LCMS: $\text{MH}^+ = 465.0$

EXAMPLE 512:

- 10 By essentially the same procedure set forth in Example 511 only substituting the compound prepared in Example 192, the above compound was prepared. Yield = 64; $\text{MH}^+ = 453$; m. p. = 238.9°C.

EXAMPLE 513:

15







- The compound prepared in Example 157 was dissolved in dioxane (30 mL) and a HCl-dioxane solution (4 M, 30 mL) was added. The reaction mixture was stirred at room temperature for 4 h. The reaction mixture was evaporated under reduced pressure and ethyl acetate (200 mL) was added. The organic
- 20











solution was washed with 1 N sodium hydroxide followed by saturated brine. The organic layer was dried over anhydrous sodium sulfate and evaporated under reduced pressure. $MH^+ = 442.1$

EXAMPLE 514-526:









- 5 By essentially the same procedure set forth in Example 513, only substituting the compounds shown in Column 2 of Table 41, the compounds shown in Column 3 of Table 41 were prepared.





TABLE 41

Ex.	Column 2	Column 3	CMPD
514			$MH^+ = 420.1$
515			$MH^+ = 442.1$
516			$MH^+ = 380.1$

517			MH ⁺ = 406.1
518			MH ⁺ = 380.1
519			MH ⁺ = 394.1
520			MH ⁺ = 366
521			MH ⁺ = 394

300

522			MH ⁺ = 408.1
523			MH ⁺ = 420.1
524			
525			MH ⁺ = 420.1

526			MH ⁺ = 428.1
526.10			









EXAMPLES 528-564:**General procedure for 5-piperidinyl parallel library formation:**

- 5 To a mixture of the starting material (80 mg, 0.21 mmol) shown in Column 2 of Table 42 in anhydrous CH₂Cl₂ (1.5 mL) was added DIPEA (75 μ L, 0.42 mmol) and the appropriate capping reagent (1.1 equiv., 0.23 mmol). After 1 to 2 h, the reaction mixture was applied to 1000 micron preparatory TLC plate and was subsequently developed using a 8 – 10 % EtOH – CH₂Cl₂ as eluent to afford
- 10 the compounds shown in Column 3 of Table 42.









TABLE 42









Ex.	Column 2	Column 3	CMPD
-----	----------	----------	------

302









528			MH ⁺ = 608 m. p. = 230.1 °C
529			Yield = 82 MH ⁺ = 614 m. p. = 235.4 °C
530			MH ⁺ = 486 m. p. = 60.5 °C
531			MH ⁺ = 500 m. p. = 113.6 °C

303









532			$MH^+ = 430$ m. p. = 158.3 – 159.2 °C
533			$MH^+ = 531$ m. p. = 105.9 °C
534			$MH^+ = 486$
535			$MH^+ = 500$









536			MH ⁺ = 430
537			MH ⁺ = 531
538			MH ⁺ = 486 m. p. = 69.6 °C
539			MH ⁺ = 500 m. p. = 82.3 °C

305









540			MH ⁺ = 430 m. p. = 223.6 °C
541			MH ⁺ = 531 m. p. = 118.1 °C
542			MH ⁺ = 455 m. p. = 109 – 110 °C
543			MH ⁺ = 429 m. p. = 111.5 °C









306

544			MH ⁺ = 455
545			MH ⁺ = 429
546			MH ⁺ = 455 m. p. = 80.1 °C
547			MH ⁺ = 429 m. p. = 64.7 °C









548			MH ⁺ = 494 m. p. = 76.5 °C
549			MH ⁺ = 493 m. p. = 83.6 °C
550			MH ⁺ = 465 m. p. = 207.5 °C
551			MH ⁺ = 494



308

552			MH ⁺ = 493
553			MH ⁺ = 465
554			MH ⁺ = 481 m. p. = 102.7 °C
555			MH ⁺ = 494 m. p. = 85.3 °C

556			MH ⁺ = 493 m. p. = 89.1 °C
557			MH ⁺ = 465 m. p. = 83.8 °C
558			Yield = quant. MH ⁺ = 443 m. p. = 98.3 °C (HCl salt)
559			MH ⁺ = 454

310

560			Yield = quant. MH ⁺ = 429 m. p. = 111.5 – 112.6 °C
561			MH ⁺ = 460 m. p. = 122.7 °C
562			MH ⁺ = 460 m. p. = 95.4 °C
563			MH ⁺ = 460

564			$MH^+ = 460$ m. p. = 95.4 °C
-----	---	--	---------------------------------

Additional data for select examples given below.

Example 534: 1H NMR (300MHz, $CDCl_3$) δ 8.66 – 8.62 (s, 1H), 8.62 – 8.58 (d, 1H), 7.95 (s, 1H), 7.72 – 7.68 (d, 1H), 7.36 – 7.31 (dd, 1H), 6.66 – 6.62 (t, 1H), 5.93 (s, 1H), 4.65 – 4.62 (d, 2H), 3.86 – 3.82 (d, 1H), 3.65 – 3.58 (m, 1H), 3.26 – 3.12 (dd, 4H), 3.02 – 2.80 (m, 3H), 2.10 – 2.00 (m, 1H), 1.67 – 1.57 (m, 3H).

Example 535: 1H NMR (300MHz, $CDCl_3$) δ 8.66 – 8.62 (s, 1H), 8.62 – 8.58 (d, 1H), 7.95 (s, 1H), 7.72 – 7.67 (d, 1H), 7.36 – 7.30 (dd, 1H), 6.70 – 6.64 (t, 1H), 5.90 (s, 1H), 4.63 – 4.61 (d, 2H), 3.93 – 3.86 (m, 1H), 3.69 – 3.61 (m, 4H), 3.27 – 3.23 (m, 4H), 3.10 – 3.01 (dd, 1H), 2.93 – 2.84 (m, 2H), 2.08 – 2.03 (m, 1H), 1.90 – 1.57 (m, 4H).

Example 536: 1H NMR (300MHz, $CDCl_3$) δ 8.67 (s, 1H), 8.62 – 8.58 (d, 1H), 7.96 (s, 1H), 7.72 – 7.68 (d, 1H), 7.36 – 7.30 (dd, 1H), 6.79 – 6.72 (t, 1H), 5.96 (s, 1H), 4.86 (br s, 2H), 4.66 – 4.63 (d, 2H), 3.89 – 3.73 (m, 2H), 3.55 – 3.32 (m, 2H), 3.00 – 2.89 (m, 1H), 2.10 – 1.97 (m, 2H), 1.70 – 1.53 (m, 2H).

Example 537: 1H NMR (300MHz, $CDCl_3$) δ 8.66 (s, 1H), 8.62 – 8.58 (d, 1H), 7.98 (s, 1H), 7.77 – 7.76 (t, 1H), 7.72 – 7.69 (d, 1H), 7.63 – 7.59 (m, 1H), 7.56 (s, 1H), 7.36 – 7.29 (dd, 1H), 6.83 – 6.79 (t, 1H), 5.96 (s, 1H), 4.67 – 4.64 (d, 2H), 3.98 – 3.93 (dd, 1H), 3.79 – 3.68 (m, 2H), 3.37 – 3.28 (m, 1H), 3.03 – 2.94 (m, 1H), 2.12 – 1.99 (m, 1H), 1.76 – 1.56 (m, 3H).

Example 544: 1H NMR (300MHz, $CDCl_3$) δ 8.66 – 8.62 (d, 1H), 8.61 – 8.58 (dd, 1H), 7.95 (s, 1H), 7.72 – 7.67 (d, 1H), 7.36 – 7.30 (dd, 1H), 6.80 – 6.62 (br s, 1H), 5.88 (s, 1H), 4.63 (s, 2H), 3.08 – 2.95 (m, 2H), 2.87 – 2.80 (m, 2H), 2.04 (m,

1H), 1.85 – 1.78 (m, 4H), 1.52 – 1.44 (m, 1H), 0.87 – 0.82 (m, 2H), 0.72 – 0.66 (m, 2H).

Example 545: ^1H NMR (300MHz, CDCl_3) δ 8.66 (s, 1H), 8.62 – 8.58 (br t, 1H), 7.97 (s, 1H), 7.73 – 7.68 (d, 1H), 7.36 – 7.30 (br t, 1H), 6.79 – 6.72 (br t, 1H),
5 5.96 (s, 1H), 4.64 (br s, 2H), 4.59 – 4.46 (br d, 1H), 3.95 – 3.74 (br m, 1H), 3.57 – 3.49 (dd, 1H), 3.10 – 3.01 (dd, 1H), 2.86 – 2.70 (m, 2H), 2.13 (s, 3H), 2.06 – 2.00 (m, 2H), 1.65 – 1.48 (m, 2H).


Example 551: ^1H NMR (300MHz, CDCl_3) δ 8.67 (s, 1H), 8.63 – 8.59 (d, 1H), 7.96 (s, 1H), 7.74 – 7.69 (d, 1H), 7.36 – 7.30 (dd, 1H), 6.69 – 6.64 (t, 1H), 5.95
10 (s, 1H), 4.67 – 4.63 (d, 2H), 3.85 3.65 (m, 1H), 3.75 – 3.65 (m, 1H), 3.25 – 3.18 (dd, 1H), 3.03 – 2.90 (m, 2H), 2.81 (s, 6H), 2.03 – 1.95 (m, 1H), 1.89 – 1.68 (m, 3H).

Example 552: ^1H NMR (300MHz, CDCl_3) δ 8.67 (s, 1H), 8.62 – 8.59 (d, 1H), 7.95 (s, 1H), 7.74 – 7.69 (d, 1H), 7.36 – 7.31 (dd, 1H), 6.67 – 6.60 (t, 1H), 5.98
15 (s, 1H), 4.67 – 4.63 (d, 2H), 3.92 – 3.86 (m, 1H), 3.85 – 3.75 (m, 1H), 3.40 – 3.30 (dd, 1H), 3.27 – 3.16 (m, 1H), 3.10 – 2.86 (m, 2H), 2.10 – 1.78 (m, 3H), 1.40 – 1.30 (d, 6H).

Example 553: ^1H NMR (300MHz, CDCl_3) δ 8.67 (s, 1H), 8.62 (br s, 1H), 7.96 (s, 1H), 7.74 – 7.69 (d, 1H), 7.36 – 7.31 (dd, 1H), 6.70 – 6.66 (t, 1H), 5.98 (s, 1H),
20 4.67 – 4.63 (d, 2H), 3.88 – 3.81 (m, 1H), 3.71 – 3.65 (m, 1H), 3.20 – 3.11 (dd, 1H), 3.02 – 2.91 (m, 1H), 2.90 – 2.80 (m, 4H), 2.01 – 1.80 (m, 3H).

Example 559: ^1H NMR (300MHz, CDCl_3) δ 8.66 – 8.60 (d, 1H), 8.50 – 8.44 (dd, 1H), 8.01 (s, 1H), 7.93 (m, 1H), 7.48 – 7.40 (dd, 1H), 6.08 (s, 1H), 4.80 – 7.74 (s, 2H), 4.32 – 4.19 (br d, 2H), 3.10 – 2.86 (m, 2H), 1.95 – 1.68 (m, 4H).

Example 563: ^1H NMR (300MHz, CDCl_3) δ 8.66 (s, 1H), 8.62 – 8.58 (d, 1H), 7.96 (s, 1H), 7.73 – 7.68 (d, 1H), 7.36 – 7.30 (dd, 1H), 6.96 – 6.86 (br s, 1H), 6.79 – 6.74 (t, 1H), 6.00 (s, 1H), 4.67 – 4.64 (d, 2H), 4.37 – 4.30 (dd, 1H), 4.22 – 4.13 (m, 1H), 3.97 – 3.86 (dd, 1H), 3.73 – 3.64 (m, 1H), 3.17 – 3.14 (d, 3H), 3.07 – 2.99 (m, 1H), 2.20 – 1.97 (m, 2H), 1.68 – 1.48 (m, 2H).

GENERAL PROCEDURE 1: Procedure for the amide formation parallel synthesis:


5

Parallel synthesis was conducted in polypropylene 96-well reaction blocks with removable top seal and fixed bottom seal. Each reaction well was fitted with a 20 micron polypropylene bottom frit and the maximum volume was 3 mL.

- Collection block was not fitted with bottom frit. To each reaction well was added a solution of an amine (0.021 mmol) dissolved in a DMF-THF-MeCN mixture (4:3:3 v/v, 0.95 mL), EDC resin (P-EDC, Polymer Laboratories Ltd., 43 mg, 0.063 mmol), 1-hydroxybenzotriazole (HOBt, 5.67 mg, 0.042 mmol) and a solution of a carboxylic acid in dimethylformamide (1 M, 0.0315 mL, 0.0315 mmol). The reaction mixture was agitated at room temperature for 16 h. The crude product solution was filtered into a reaction well loaded with trisamine resin (P-NH₂, Argonaut Tech. Inc., 30 mg, 0.126 mmol) and isocyanate resin (P-NCO, Argonaut Tech. Inc., 35 mg, 0.063 mmol). The reaction mixture was agitated at room temperature for 16 h and filtered into the collection block. The product solution was evaporated under reduced pressure to afford the desired amide product.

GENERAL PROCEDURE 2: Procedure for the sulfonamide formation parallel synthesis


314



Parallel synthesis was conducted in polypropylene 96-well reaction blocks with removable top seal and fixed bottom seal. Each reaction well was fitted with

- 5 a 20 micron polypropylene bottom frit and the maximum volume was 3 mL. Collection block was not fitted with bottom frit. To each reaction well was added a solution of an amine (0.021 mmol) dissolved in a DMF-THF-MeCN mixture (3:2:2 v/v, 0.95 mL), DIEA resin (P-DIEA, Argonaut Tech. Inc., 18 mg, 0.063 mmol) and a solution of a sulfonyl chloride in dimethylformamide (1 M, 0.0315 mL, 0.0315 mmol). The reaction mixture was agitated at room temperature for
- 10 16 h. The crude product solution was filtered into a reaction well loaded with trisamine resin (P-NH₂, Argonaut Tech. Inc., 30 mg, 0.126 mmol) and isocyanate resin (P-NCO, Argonaut Tech. Inc., 35 mg, 0.063 mmol). The reaction mixture was agitated at room temperature for 16 h and filtered into the
- 15 collection block. The product solution was evaporated under reduced pressure to afford the desired sulfonamide product.

GENERAL PROCEDURE 3: Procedure for the urea formation parallel synthesis



20


Parallel synthesis was conducted in polypropylene 96-well reaction blocks with removable top seal and fixed bottom seal. Each reaction well was fitted with

a 20 micron polypropylene bottom frit and the maximum volume was 3 mL.

Collection block was not fitted with bottom frit. To each reaction well was added a solution of an amine (0.021 mmol) dissolved in a DMF-MeCN mixture (1:1 v/v, 0.95 mL) and a solution of an isocyanate in dichloromethane (0.33 M, 0.126 mL, 0.042 mmol). The reaction mixture was agitated at room temperature for 16 h.

The crude product solution was filtered into a reaction well loaded with trisamine resin (P-NH₂, Argonaut Tech. Inc., 30 mg, 0.126 mmol) and isocyanate resin (P-NCO, Argonaut Tech. Inc., 35 mg, 0.063 mmol). The reaction mixture was agitated at room temperature for 16 h and filtered into the collection block. The product solution was evaporated under reduced pressure to afford the desired urea product.

GENERAL PROCEDURE 4: Procedure for the reductive alkylation parallel synthesis




Parallel synthesis was conducted in polypropylene 96-well reaction blocks with removable top seal and fixed bottom seal. Each reaction well was fitted with a 20 micron polypropylene bottom frit and the maximum volume was 3 mL.

Collection block was not fitted with bottom frit. To each reaction well was added a solution of an amine (0.021 mmol) dissolved in AcOH-DCE mixture (1:99 v/v, 0.5 mL), a solution of an aldehyde or ketone in dichloroethane (1 M, 0.147 mL, 0.147 mmol), and a solution of tetramethylammonium triacetoxyborohydride (11 mg, 0.042 mmol) dissolved in AcOH-DCE mixture 1:99 v/v, 0.5 mL). The reaction mixture was agitated at room temperature for 3 days. The crude product solution was filtered into a reaction well loaded with sulfonic acid resin Lanterns (P-SO₃H, Mimotopes Pty Ltd., 0.3 mmol). The reaction mixture was agitated at room temperature for 2 h and decanted. The product resin Lanterns

were washed with methanol (1 mL) for three times. A solution of ammonia in methanol (2 M, 1.2 mL) was added. The reaction mixture was agitated at room temperature for 30 min. and filtered into the collection block. The product solution was evaporated under reduced pressure to afford the desired tertiary amine product.

GENERAL PROCEDURE 5: Procedure for the parallel synthesis of 7,N-substituted pyrazolo[1,5a]pyrimidines



To 3-bromo-7-chloro-5-(2-chloro-phenyl)-pyrazolo[1,5-a]pyrimidine (9.0 mg, 0.03 mmol) in tetrahydrofuran were added di-*iso*-propylethylamine (12 μ L, 0.07), followed by cyclopropylmethylethylamine (70 μ L, .07 mmol; 1M solution in DMF). The reaction mixture was heated to 70 °C for 36 h and then cooled to rt.


The mixture was treated with (P-NCO, Argonaut Tech. Inc 70 mg, 0.12 mmol), and P-CO₃⁻ (Argonaut Tech. Inc 70 mg, 0.24 mmol) and shaken at rt for 12-18 h. The solution was filtered and evaporated to dryness to provide the product. observed m/z 375.21.

GENERAL PROCEDURE 6: Procedure for the parallel synthesis of 5,N-substituted pyrazolo[1,5a]pyrimidines

General protocols:

Parallel synthesis was performed in a 96 well polypropylene blocks as described elsewhere. In the instance that heating was required, reactions were conducted in 2.5 mL glass tubes individually sealed with a polypropylene mat and heating achieved by a 96 well heat transfer block.

317

**STEP A:**

- 5 To the 3-bromo-5-chloro-7-N-Boc-alkylamino-pyrazolo[1,5-a]pyrimidine (17 mg, 0.04 mmol) in *p*-dioxane were added DIEA (9 μ L, 0.05), followed by cyclopropyl-methylamine (80 μ L, .08 mmol; 1M solution in isopropanol). The reaction mixture was heated to 90 °C for 36 h and then cooled to rt. The mixture was treated with **P**-NCO (Argonaut Tech. Inc. 70 mg, 0.12 mmol) and **P**-CO₃⁻ (Argonaut Tech. Inc. 70 mg, 0.24 mmol) and shaken at rt for 12-18 h. The solution was filtered and evaporated to dryness to provide the product.

STEP B(acidic):

- 15 The product from STEP A was taken up in 35% TFA/DCM and agitated for 4 h followed by concentration under high vacuum. The residue was treated with 10% HCl(aq) in MeOH agitated for 2 h and then concentrated to give the desired product. . observed *m/z* 375.21.


STEP B(basic):

- 20 The product from step A was taken up in EtOH and treated with Ambersep[®] 900-OH ion exchange resin (Acros, 100mg), heated at reflux for 48 h with gently stirring. The reaction mixture was cooled to rt, filtered and concentrated to provide the desired product.


EXAMPLE 565:

- 25 By utilizing the procedure set forth in General Procedure 1 and the compound from Example 462 shown below, the compounds with the observed *m/z* shown in Table 43 were prepared.


318

**EXAMPLE 566:**

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 471 shown below, the compounds shown in Table 44 with the observed m/z were prepared.


**EXAMPLE 567:**

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 515 shown below, the compounds shown in Table 45 with the observed m/z were prepared.


**EXAMPLE 568:**

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 513 shown below, the compounds shown in Table 46 with the observed m/z were prepared.


319

**EXAMPLE 569:**

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 526 shown below, the compounds shown in Table 47
5 with the observed m/z were prepared.


**EXAMPLE 570:**

By utilizing the procedure set forth in General Procedure 1 and the
10 compound from Example 524 shown below, the compounds shown in Table 48 with the observed m/z were prepared.

**EXAMPLE 571:**

320


By utilizing the procedure set forth in General Procedure 1 and the compound from Example 525 shown below, the compounds shown in Table 49 with the observed m/z were prepared.



5

EXAMPLE 572:


By utilizing the procedure set forth in General Procedure 1 and the compound from Example 526.10 shown below, the compounds shown in Table 50 with the observed m/z were prepared.



10

EXAMPLE 573:

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 518 shown below, the compounds shown in Table 51 with the observed m/z were prepared.




15

EXAMPLE 574:


321

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 519 shown below, the compounds shown in Table 52 with the observed m/z were prepared.



5 **EXAMPLE 575:**


By utilizing the procedure set forth in General Procedure 1 and the compound from Example 520 shown below, the compounds shown in Table 53 with the observed m/z were prepared.



10

EXAMPLE 576:

By utilizing the procedure set forth in General Procedure 1 and the compound from Example 522 shown below, the compounds shown in Table 54 with the observed m/z were prepared.




15

EXAMPLE 577:

322


By utilizing the procedure set forth in General Procedure 1 and the compound from Example 523 shown below, the compounds shown in Table 55 with the observed m/z were prepared.



5

EXAMPLE 578:

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 462 shown below, the compounds shown in Table 56 with the observed m/z were prepared.




10

EXAMPLE 579:


By utilizing the procedure set forth in General Procedure 2 and the compound from Example 471 shown below, the compounds shown in Table 57 with the observed m/z were prepared.

15

**EXAMPLE 580:**

323


By utilizing the procedure set forth in General Procedure 2 and the compound from Example 515 shown below, the compounds shown in Table 58 with the observed m/z were prepared.



5

EXAMPLE 581:

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 513 shown below, the compounds shown in Table 59 with the observed m/z were prepared.




10

EXAMPLE 582:


By utilizing the procedure set forth in General Procedure 2 and the compound from Example 513 shown below, the compounds shown in Table 60 with the observed m/z were prepared.

15


324

**EXAMPLE 583:**

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 524 shown below, the compounds shown in Table 61 with the observed m/z were prepared.


**EXAMPLE 584:**

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 525 shown below, the compounds shown in Table 62 with the observed m/z were prepared.

**EXAMPLE 585:**

325


By utilizing the procedure set forth in General Procedure 2 and the compound from Example 526.10 shown below, the compounds shown in Table 63 with the observed m/z were prepared.



5

EXAMPLE 586:

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 518 shown below, the compounds shown in Table 64 with the observed m/z were prepared.




10

EXAMPLE 587:

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 519 shown below, the compounds shown in Table 65 with the observed m/z were prepared.


15



EXAMPLE 588:

326

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 520 shown below, the compounds shown in Table 67 with the observed m/z were prepared.




5

EXAMPLE 589:

By utilizing the procedure set forth in General Procedure 2 and the compound from Example 521 shown below, the compounds shown in Table 68 with the observed m/z were prepared.


10

**EXAMPLE 590:**


By utilizing the procedure set forth in General Procedure 2 and the compound from Example 523 shown below, the compounds shown in Table 69 with the observed m/z were prepared.

15


327

**EXAMPLE 591:**

By utilizing the procedure set forth in General Procedure 3 and the
5 compound from Example 462 shown below, the compounds shown in Table 70
with the observed m/z were prepared.

**EXAMPLE 592:**

10 By utilizing the procedure set forth in General Procedure 3 and the
compound from Example 471 shown below, the compounds shown in Table 71
with the observed m/z were prepared.




15

EXAMPLE 593:

328


By utilizing the procedure set forth in General Procedure 3 and the compound from Example 513 shown below, the compounds shown in Table 72 with the observed m/z were prepared.



5

EXAMPLE 594:

By utilizing the procedure set forth in General Procedure 3 and the compound from Example 524 shown below, the compounds shown in Table 73 with the observed m/z were prepared.




10

EXAMPLE 595:


By utilizing the procedure set forth in General Procedure 3 and the compound from Example 524 shown below, the compounds shown in Table 74 with the observed m/z were prepared.

15


329

**EXAMPLE 596:**

By utilizing the procedure set forth in General Procedure 3 and the compound from Example 519 shown below, the compounds shown in Table 75 with the observed m/z were prepared.


**EXAMPLE 597:**

By utilizing the procedure set forth in General Procedure 3 and the compound from Example 520 shown below, the compounds shown in Table 76 with the observed m/z were prepared.

**EXAMPLE 598:**

330


By utilizing the procedure set forth in General Procedure 3 and the compound from Example 521 shown below, the compounds shown in Table 77 with the observed m/z were prepared.



5

EXAMPLE 599:


By utilizing the procedure set forth in General Procedure 3 and the compound from Example 523 shown below, the compounds shown in Table 78 with the observed m/z were prepared.



10

EXAMPLE 600:


By utilizing the procedure set forth in General Procedure 4 and the compound from Example 462 shown below, the compounds shown in Table 79 with the observed m/z were prepared.




15

EXAMPLE 601:

By utilizing the procedure set forth in General Procedure 4 and the compound from Example 471 shown below, the compounds shown in Table 80 with the observed m/z were prepared.


**EXAMPLE 602:**

By utilizing the procedure set forth in General Procedure 4 and the compound from Example 525 shown below, the compounds shown in Table 81 with the observed m/z were prepared.


**EXAMPLE 603:**

By utilizing the procedure set forth in General Procedure 4 and the compound from Example 526.10 shown below, the compounds shown in Table 82 with the observed m/z were prepared.


332

**EXAMPLE 604:**

By utilizing the procedure set forth in General Procedure 4 and the compound from Example 521 shown below, the compounds shown in Table 83 with the observed m/z were prepared.


**EXAMPLE 605:**

By utilizing the procedure set forth in General Procedure 4 and the compound from Example 523 shown below, the compounds shown in Table 84 with the observed m/z were prepared.

**EXAMPLE 606:**


By utilizing the procedure set forth in General Procedure 5 and the compound from Preparative Example 81 shown below, the compounds shown in Table 85 with the observed m/z were prepared.

333

**EXAMPLE 607:**

By utilizing the procedure set forth in General Procedure 6 and the compound from Preparative Example 196, the compounds shown in Table 86

5 with the observed m/z were prepared.

**ASSAY:**

BACULOVIRUS CONSTRUCTIONS: Cyclins A and E were cloned into

10 pFASTBAC (Invitrogen) by PCR, with the addition of a GluTAG sequence (EYMPME) at the amino-terminal end to allow purification on anti-GluTAG affinity columns. The expressed proteins were approximately 46kDa (cyclin E) and 50kDa (cyclin A) in size. CDK2 was also cloned into pFASTBAC by PCR, with the addition of a haemagglutinin epitope tag at the carboxy-terminal end
15 (YDVPDYAS). The expressed protein was approximately 34kDa in size.

ENZYME PRODUCTION: Recombinant baculoviruses expressing cyclins A, E and CDK2 were infected into SF9 cells at a multiplicity of infection (MOI) of 5, for 48 hrs. Cells were harvested by centrifugation at 1000 RPM for 10 minutes.

20 Cyclin-containing (E or A) pellets were combined with CDK2 containing cell pellets and lysed on ice for 30 minutes in five times the pellet volume of lysis buffer containing 50mM Tris pH 8.0, 0.5% NP40, 1mM DTT and protease/phosphatase inhibitors (Roche Diagnostics GmbH, Mannheim, Germany). Mixtures were stirred for 30-60 minutes to promote cyclin-CDK2
25 complex formation. Mixed lysates were then spun down at 15000 RPM for 10

minutes and the supernatant retained. 5ml of anti-GluTAG beads (for one liter of SF9 cells) were then used to capture cyclin-CDK2 complexes. Bound beads were washed three times in lysis buffer. Proteins were competitively eluted with lysis buffer containing 100-200ug/mL of the GluTAG peptide. Eluate was

5 dialyzed overnight in 2 liters of kinase buffer containing 50mM Tris pH 8.0, 1mM DTT, 10mM MgCl₂, 100uM sodium orthovanadate and 20% glycerol. Enzyme was stored in aliquots at -70°C.

IN VITRO KINASE ASSAY: CDK2 kinase assays (either cyclin A or E-

10 dependent) were performed in low protein binding 96-well plates (Corning Inc, Corning, New York). Enzyme was diluted to a final concentration of 50 ug/ml in kinase buffer containing 50mM Tris pH 8.0, 10mM MgCl₂, 1mM DTT, and 0.1mM sodium orthovanadate. The substrate used in these reactions was a biotinylated peptide derived from Histone H1 (from Amersham, UK). The substrate was
15 thawed on ice and diluted to 2 uM in kinase buffer. Compounds were diluted in 10%DMSO to desirable concentrations. For each kinase reaction, 20 ul of the 50 ug/ml enzyme solution (1 ug of enzyme) and 20 ul of the 1 uM substrate solution were mixed, then combined with 10 ul of diluted compound in each well for testing. The kinase reaction was started by addition of 50 ul of 4 uM ATP and 1
20 uCi of ³³P-ATP (from Amersham, UK). The reaction was allowed to run for 1 hour at room temperature. The reaction was stopped by adding 200 ul of stop buffer containing 0.1% Triton X-100, 1mM ATP, 5mM EDTA, and 5 mg/ml streptavidine coated SPA beads (from Amersham, UK) for 15 minutes. The SPA beads were then captured onto a 96-well GF/B filter plate (Packard/Perkin Elmer
25 Life Sciences) using a Filtermate universal harvester (Packard/Perkin Elmer Life Sciences.). Non-specific signals were eliminated by washing the beads twice with 2M NaCl then twice with 2 M NaCl with 1% phosphoric acid. The radioactive signal was then measured using a TopCount 96 well liquid scintillation counter (from Packard/Perkin Elmer Life Sciences).




30

IC₅₀ DETERMINATION: Dose-response curves were plotted from inhibition data generated, each in duplicate, from 8 point serial dilutions of inhibitory





compounds. Concentration of compound was plotted against % kinase activity, calculated by CPM of treated samples divided by CPM of untreated samples.

To generate IC₅₀ values, the dose-response curves were then fitted to a standard sigmoidal curve and IC₅₀ values were derived by nonlinear regression analysis. The thus-obtained IC₅₀ values for the compounds of the invention are shown in **Table 87**. These kinase activities were generated by using cyclin A or cyclin E using the above-described assay.





Table 87

CMPD	Example	IC ₅₀ (μM)
	1	0.020 0.029
	3	0.032 0.024
	4	0.011


336

	5	0.021
	8	0.003
	6	0.064 0.029
	7	0.01 0.006

337

	10	0.042
	12	0.17
	16	0.62
	1	5.6









338









	3	0.14
---	---	------

As demonstrated above by the assay values, the compounds of the present invention exhibit excellent CDK inhibitory properties.









While the present invention has been described with in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.











TABLE 43











Ex.	Compound	m/z	Ex.	Compound	m/z
4301		457.25	4305		487.27
4302		471.26	4306		487.27
4303		477.26	4307		487.27
4304		483.27	4308		494.27











Ex.	Product	m/z	Ex.	Product	m/z
4309		499.27	4313		509.28
4310		500.27	4314		512.28
4311		503.28	4315		513.28
4312		505.28	4316		513.28











341











Ex.	Product	m/z	Ex.	Product	m/z
4317		518.28	4321		523.29
4318		518.28	4322		523.29
4319		519.29	4323		523.29
4320		521.29	4324		525.29

Product	1.Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4325 2. 527.29			1. 4330 2. 533.29
	1. 4326 2. 527.29			4331
	1. 4327 2. 532.29			1. 4332 2. 533.29
	1. 4328 2. 532.29			1. 4333 2. 533.29
	1. 4329 2. 532.29			1. 4334 2. 537.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4335 2. 537.3		1. 4340 2. 541.3
	1. 4336 2. 537.3		1. 4341 2. 543.3
	1. 4337 2. 539.3		1. 4342 2. 546.3
	1. 4338 2. 539.3		1. 4343 2. 547.3
	1. 4339 2. 539.3		1. 4344 2. 547.3

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4345 2. 549.3			1. 4350 2. 553.3
	1. 4346 2. 550.3			1. 4351 2. 557.31
	1. 4347 2. 551.3			1. 4352 2. 557.31
	1. 4348 2. 551.3			1. 4353 2. 558.31
	1. 4349 2. 551.3			1. 4354 2. 561.31

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4355 2. 561.31			1. 4360 2. 569.31
	1. 4356 2. 561.31			1. 4361 2. 574.32
	1. 4357 2. 561.31			1. 4362 2. 573.32
	1. 4358 2. 561.31			1. 4363 2. 573.32
	1. 4359 2. 569.31			1. 4364 2. 575.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4365 2. 575.32		1. 43700 2. 585.32
	1. 43666 2. 575.32		1. 4371 2. 583.32
	1. 4367 2. 574.32		1. 4372 2. 585.32
	1. 43688 2. 583.32		1. 4373 2. 585.32
	1. 4369 2. 583.32		1. 4374 2. 597.33


































Product	1. Ex. 2. m/z
 <chem>BrC1=CN2C(=N1)C(=CN2)C3=CC=CC=C3N(C3)C4CCN(CC4)C(=O)C5CCCCC5</chem>	1. 4375 2. 499.27
 <chem>BrC1=CN2C(=N1)C(=CN2)C3=CC=CC=C3N(C3)C4CCN(CC4)C(=O)C5=CC=CC=C5</chem>	1. 4376 2. 493.27
 <chem>BrC1=CN2C(=N1)C(=CN2)C3=CC=CC=C3N(C3)C4CCN(CC4)C(=O)C5=C(C)C(=C(C)C)C(C)=C5</chem>	1. 4377 2. 535.29











TABLE 44











Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4401 2. 471.3			1. 4406 2. 487.27
	1. 4402 2. 475.26			1. 4407 2. 487.27
	1. 4403 2. 483.27			1. 4408 2. 487.27
	1. 4404 2. 481.26			1. 4409 2. 494.3
	1. 4405 2. 485.27			1. 4410 2. 496.27











Product	1. Ex 2. m/z		Product	1. Ex. 2. m/z
	1. 4411 2. 499.27			1. 4416 2. 507.28
	2. 4412 2. 499.27			1. 4417 2. 509.3
	1. 4413 2. 500.27			1. 4418 2. 512.28
	1. 4414 2. 503.28			1. 4419 2. 513.28
	1. 4415 2. 505.28			1. 4420 2. 513.28











Product	1. x. 2. m/z		Product	1. Ex. 2. m/z
	1. 4421 2. 513.28			1. 4426 2. 521.29
	1. 4422 2. 513.28			1. 4428 2. 521.29
	1. 4423 2. 518.28			1. 4428 2. 523.29
	1. 4424 2. 518.28			1. 4429 2. 523.29
	1. 4425 2. 519.3			1. 4430 2. 523.29









Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4431 2. 523.29			1. 4436 2. 527.29
	1. 4432 2. 525.29			1. 4437 2. 532.29
	1. 4433 2. 525.3			1. 4438 2. 532.29
	1. 4434 2. 527.29			1. 4439 2. 532.29
	1. 4435 2. 527.29			1. 4440 2. 531.29

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4441 2. 533.3			1. 4446 2. 537.3
	1. 4442 2. 531.29			1. 4447 2. 537.3
	1. 4443 2. 533.29			1. 4448 2. 539.3
	1. 4444 2. 535.29			1. 4449 2. 539.3
	1. 4445 2. 537.3			1. 4450 2. 539.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4451 2. 541.3		1. 4456 2. 549.3
	1. 4451 2. 543.3		1. 4457 2. 550.3
	1. 4453 2. 546.3		1. 4458 2. 550.3
	1. 4454 2. 547.3		1. 4459 2. 551.3
	1. 4455 2. 547.3		1. 4460 2. 551.3
















Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4461 2. 549.3			1. 4466 2. 561.31
	1. 4462 2. 553.3			1. 4467 2. 561.31
	1. 4463 2. 557.3			1. 4468 2. 561.31
	1. 4464 2. 557.31			1. 4469 2. 561.31
	1. 4465 2. 558.31			1. 4470 2. 562.3
















Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4471 2. 569.31			1. 4476 2. 572.31
	1. 4472 2. 569.31			1. 4477 2. 573.32
	1. 4473 2. 572.31			1. 4478 2. 574.32
	1. 4474 2. 572.31			1. 4479 2. 576.32
	1. 4475 2. 575.32			1. 4480 2. 583.32
















Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4481 2. 583.32			1. 4486 2. 585.32
	1. 4482 2. 583.32			1. 4487 2. 597.33
	1. 4483 2. 585.32			1. 4488 2. 499.27
	1. 4484 2. 585.32			
	1. 4485 2. 585.3			
















357

TABLE 45
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4501 2. 512.28		1. 4506 2. 540.3		1. 4511 2. 551.3
	1. 4502 2. 526.29		1. 4507 2. 542.3		1. 4512 2. 554.3
	1. 4503 2. 532.29		1. 4508 2. 542.3		1. 4513 2. 554.3
	1. 4504 2. 538.3		1. 4509 2. 542.3		1. 4514 2. 555.31
	1. 4505 2. 538.3		1. 4510 2. 549.3		1. 4515 2. 558.31
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4516 2. 560.31		1. 4521 2. 568.31		1. 4526 2. 574.32
	1. 4517 2. 562.31		1. 4522 2. 568.31		1. 4527 2. 576.32
	1. 4518 2. 564.31		1. 4523 2. 568.31		1. 4528 2. 576.32
	1. 4519 2. 567.31		1. 4524 2. 573.32		1. 4529 2. 578.32
	1. 4520 2. 568.31		1. 4525 2. 573.32		1. 4530 2. 578.32

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4531 2. 578.32			1. 4536 2. 582.32			1. 4541 2. 588.32
	1. 4532 2. 578.32			1. 4537 2. 582.32			1. 4542 2. 588.32
	1. 4533 2. 580.32			1. 4538 2. 587.32			1. 4543 2. 588.32
	1. 4534 2. 580.32			1. 4539 2. 587.32			1. 4544 2. 588.32
	1. 4535 2. 582.32			1. 4540 2. 587.32			1. 4545 2. 590.32











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4546 2. 588.32		1. 4551 2. 592.33		1. 4556 2. 593.33
	1. 4547 2. 588.32		1. 4552 2. 592.33		1. 4557 2. 596.33
	1. 4548 2. 588.32		1. 4553 2. 592.33		1. 4558 2. 596.33
	1. 4549 2. 588.32		1. 4554 2. 594.33		1. 4559 2. 598.33
	1. 4550 2. 590.32		1. 4555 2. 594.33		1. 4560 2. 601.33

361

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4561 2. 602.33		1. 4665 2. 605.33		1. 4571 2. 612.34
	1. 4562 2. 602.33		1. 4567 2. 606.33		1. 4572 2. 612.34
	1. 4563 2. 604.33		1. 4568 2. 606.33		1. 4573 2. 613.34
	1. 4564 2. 603.3		1. 4569 2. 606.33		1. 4574 2. 616.34
	1. 4565 2. 605.33		1. 4570 2. 608.33		1. 4575 2. 616.34











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4576 2. 616.34		1. 4581 2. 624.34		1. 4586 2. 630.35
	1. 4577 2. 616.34		1. 4582 2. 629.35		1. 4587 2. 630.35
	1. 4578 2. 616.34		1. 4583 2. 629.35		1. 4588 2. 630.35
	1. 4579 2. 616.34		1. 4584 2. 630.35		1. 4589 2. 631.35
	1. 4580 2. 624.34		1. 4585 2. 630.35		1. 4590 2. 638.35











363











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4591 2. 638.35		1. 4596 2. 652.36
	1. 4592 2. 640.35		1. 4597 2. 486.27
	1. 4593 2. 640.35		1. 4598 2. 554.3
	1. 4594 2. 640.35		1. 4599 2. 548.3
	1. 4595 2. 640.35		1. 45100 2.











364











TABLE 46

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4601 2. 512.28			1. 4606 2. 540.3
	1. 4602 2. 526.29			1. 4607 2. 542.3
	1. 4603 2. 532.29			1. 4608 2. 542.3
	1. 4604 2. 538.3			1. 4609 2. 542.3
	1. 4605 2. 536.3			1. 4610 2. 547.3











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4611 2. 551.3		1. 4616 2. 560.31
	1. 4612 2. 552.3		1. 4617 2. 562.31
	1. 4613 2. 552.3		1. 4618 2. 562.3
	1. 4614 2. 555.31		1. 4619 2. 567.31
	1. 4615 2. 558.31		1. 4620 2. 568.31











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4621 2. 568.31		1. 4626 2. 574.32
	1. 4622 2. 568.31		1. 4627 2. 576.32
	1. 4623 2. 568.31		1. 4628 2. 576.32
	1. 4624 2. 573.32		1. 4629 2. 578.32
	1. 4625 2. 573.32		1. 4630 2. 578.32











Product	1. Ex. 2. m/x	Product	1. Ex. 2. m/z
	1. 4631 2. 578.32		1. 4636 2. 582.32
	1. 4632 2. 578.32		1. 4637 2. 582.32
	1. 4633 2. 580.32		1. 4638 2. 587.32
	1. 4634 2. 580.32		1. 4639 2. 585.3
	1. 4635 2. 582.32		1. 4640 2. 585.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4641 2. 588.32		1. 4646 2. 592.33
	1. 4642 2. 588.32		1. 4647 2. 592.33
	1. 4643 2. 588.32		1. 4648 2. 592.33
	1. 4644 2. 588.32		1. 4649 2.
	1. 4645 2. 590.32		1. 4650 2. 594.33











369






Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4651 2. 594.33		1. 4656 2. 602.33
	1. 4652 2. 596.33		1. 4657 2. 602.33
	1. 4653 2. 596.3		1. 4658 2. 604.33
	1. 4654 2. 598.33		1. 4659 2. 603.3
	1. 4655 2. 601.33		1. 4660 2. 605.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4661 2. 605.33		1. 4666 2. 612.34
	1. 4662 2. 606.33		1. 4667 2. 612.34
	1. 4663 2. 606.33		1. 4668 2. 613.34
	1. 4664 2. 606.33		1. 4669 2. 616.34
	1. 4665 2. 608.33		1. 4670 2. 616.34

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4671 2. 616.34		1. 4676 2. 624.34
	1. 4672 2. 616.34		1. 4677 2. 629.35
	1. 4673 2. 616.34		1. 4678 2. 629.35
	1. 4674 2. 616.34		1. 4679 2. 630.35
	1. 4675 2. 624.34		1. 4680 2. 630.35
















372

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4681 2. 630.35		1. 4686 2. 638.35
	1. 4682 2. 630.35		1. 4687 2. 640.35
	1. 4683 2. 630.35		1. 4688 2. 640.35
	1. 4684 2. 631.35		4689 2.
	1. 4685 2. 638.35		1. 4690 2. 640.35
















Product	1. Ex. 2. m/z
	1. 4691 2. 652.36
	1. 4692 2. 484.3
	1. 4693 2. 554.3
	1. 4694 2. 548.3
	1. 4695 2. 590.32

374
















TABLE 47
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4701 2. 498.27		1. 4706 2. 528.29		1. 4711 2. 544.3
	1. 4702 2. 512.28		1. 4707 2. 528.29		1. 4712 2. 546.3
	1. 4703 2. 518.28		1. 4708 2. 535.29		1. 4713 2. 548.3
	1. 4704 2. 524.29		1. 4709 2. 540.3		1. 4714 2. 550.3
	1. 4705 2. 528.29		1. 4710 2. 541.3		1. 4715 2. 553.3

375

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4716 2. 554.3		1. 4721 2. 562.31		1. 4726 2. 568.31
	1. 4717 2. 554.3		1. 4722 2. 562.31		1. 4727 2. 568.31
	1. 4718 2. 554.3		1. 4723 2. 564.31		1. 4728 2. 568.31
	1. 4719 2. 559.31		1. 4724 2. 564.31		1. 4729 2. 573.32
	1. 4720 2. 559.31		1. 4725 2. 566.31		1. 4730 2. 574.32

376

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4731 2. 574.32		1. 4736 2. 580.32		1. 4741 2. 588.32
	1. 4732 2. 576.32		1. 4737 2. 582.32		1. 4742 2. 590.32
	1. 4733 2. 578.32		1. 4738 2. 584.32		1. 4743 2. 591.33
	1. 4734 2. 578.32		1. 4739 2. 585.32		1. 4744 2. 592.33
	1. 4735 2. 580.32		1. 4740 2. 588.32		1. 4745 2. 592.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4746 2. 592.33		1. 4751 2. 602.33		1. 4756 2. 624.34
	1. 4747 2. 598.33		1. 4752 2. 615.34		1. 4757 2. 624.34
	1. 4748 2. 598.33		1. 4753 2. 615.34		1. 4758 2. 626.34
	1. 4749 2. 602.33		1. 4754 2. 616.34		1. 4759 2. 624.34
	1. 4750 2. 602.33		1. 4755 2. 616.34		1. 4760 2. 626.34

378












































Product	1. Ex. 2. m/z
	1. 4761 2. 638.35
	1. 4762 2. 472.26
	1. 4763 2. 534.29











TABLE 48











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4801 2. 498.27		1. 4806 2. 528.29
	1. 4802 2. 512.28		1. 4807 2. 528.29
	1. 4803 2. 524.29		1. 4808 2. 528.29
	1. 4804 2. 524.29		1. 4809 2. 540.3
	1. 4805 2. 526.29		1. 4810 2. 540.3

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4811 2. 544.3			1. 4816 2. 562.31
	1. 4812 2. 548.3			1. 4817 2. 562.31
	1. 4813 2. 553.3			1. 4818 2. 564.31
	1. 4814 2. 554.3			1. 4819 2. 564.31
	1. 4815 2. 560.31			1. 4820 2. 568.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4821 2. 568.31		1. 4826 2. 574.32
	1. 4822 2. 573.32		1. 4828 2. 576.32
	1. 4821 2. 574.32		1. 4829 2. 578.32
	1. 4824 2. 574.32		1. 4829 2. 580.32
	1. 4825 2. 574.32		1. 4830 2. 584.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4831 2. 588.32		1. 4836 2. 616.34
	1. 4832 2. 602.33		1. 4837 2. 624.34
	1. 4833 2. 602.33		1. 4838 2. 626.34
	1. 4834 2. 602.33		1. 4839 2. 626.34
	1. 4835 2. 610.34		1. 4840 2. 564.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4841 2. 535.29		1. 4846 2. 568.31
	1. 4842 2. 541.3		1. 4847 2. 578.32
	1. 4843 2. 559.31		1. 4848 2. 580.32
	1. 4844 2. 559.31		1. 4849 2. 580.32
	1. 4845 2. 564.31		1. 4850 2. 584.32

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 4851 2. 590.32			1. 4856 2. 616.34
	1. 4852 2. 591.33			1. 4857 2. 624.34
	1. 4853 2. 610.34			1. 4858 2. 626.34
	1. 4854 2. 615.34			1. 4859 2. 537.3
	1. 4855 2. 615.34			1. 4860 2. 550.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4861 2. 554.3		1. 4866 2. 587.32
	1. 4862 2. 566.31		1. 4867 2. 588.32
	1. 4863 2. 566.31		1. 4868 2. 592.33
	1. 4864 2. 578.32		1. 4869 2. 592.33
	1. 4865 2. 582.32		1. 4870 2. 573.32







































































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4871 2. 616.34		1. 4876 2. 594.33
	1. 4872 2. 616.34		1. 4877 2. 598.33
	1. 4873 2. 624.34		1. 4878 2. 592.33
	1. 4874 2. 638.35		1. 4879 2. 569.31
	1. 4875 2. 546.3		1. 4880 2. 583.32
















TABLE 49
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4901 2. 490.27		1. 4906 2. 518.28		1. 4911 2. 529.29
	1. 4902 2. 504.28		1. 4907 2. 520.29		1. 4912 2. 532.29
	1. 4903 2. 510.28		1. 4908 2. 520.29		1. 4913 2. 532.29
	1. 4904 2. 516.28		1. 4909 2. 520.29		1. 4914 2. 533.29
	1. 4905 2. 516.28		1. 4910 2. 527.29		1. 4915 2. 536.29

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4916 2. 538.3		1. 4921 2. 546.3		1. 4926 2. 552.3
	1. 4917 2. 540.3		1. 4922 2. 546.3		1. 4927 2. 554.3
	1. 4918 2. 542.3		1. 4923 2. 546.3		1. 4928 2. 554.3
	1. 4919 2. 545.3		1. 4924 2. 551.3		1. 4929 2. 556.31
	1. 4920 2. 546.3		1. 4925 2. 551.3		1. 4930 2. 556.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. m/z
	1. 4931 2. 556.31		1. 4936 2. 560.31		1. 494 2. 566.
	1. 4932 2. 556.31		1. 4937 2. 560.31		1. 494 2. 566.
	1. 4933 2. 558.31		1. 4938 2. 565.31		1. 494 2. 566.
	1. 4934 3. 558.31		1. 4939 2. 565.3		1. 494 2. 566.
	1. 4935 2. 560.31		1. 4940 2. 565.31		1. 494 2. 568.

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4946 2. 570.31		1. 4951 2. 571.31		1. 4956 2. 580.32
	1. 4947 2. 570.31		1. 4952 2. 574.32		1. 4957 2. 580.32
	1. 4948 2. 570.31		1. 4953 2. 576.32		1. 4958 2. 582.32
	1. 4949 2. 572.31		1. 4954 2. 576.32		1. 4959 2. 583.32
	1. 4950 2. 572.31		1. 4955 2. 579.32		1. 4960 2. 583.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4961 2. 583.32		1. 4966 2. 590.32		1. 4971 2. 594.33
	1. 4962 2. 584.32		1. 4967 2. 590.32		1. 4972 2. 594.33
	1. 4963 2. 584.32		1. 4968 2. 591.3		1. 4973 2. 594.33
	1. 4964 2. 584.32		1. 4969 2. 594.33		1. 4974 2. 594.33
	1. 4965 2. 586.32		1. 4970 2. 594.33		1. 4975 2. 602.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 4976 2. 602.33		1. 4981 2. 608.33		1. 4986 2. 616.34
	1. 4977 2. 607.33		1. 4982 2. 608.33		1. 4987 2. 618.34
	1. 4978 2. 607.33		1. 4983 2. 608.33		1. 4988 2. 618.34
	1. 4979 2. 605.33		1. 4984 2. 609.33		1. 4089 2. 618.34
	1. 4980 2. 608.33		1. 4985 2. 616.34		1. 4990 2. 618.34














































Product	1. Ex. 2. m/z
	1. 4991 2. 630.35
	1. 4992 2. 464.26
	1. 4993 2. 532.29
	1. 4994 2. 526.29
	1. 4995 2. 568.31











TABLE 50











Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5001 2. 476.26			1. 5006 2. 504.28
	1. 5002 2. 490.27			1. 5007 2. 506.28
	1. 5003 2. 496.27			1. 5008 2. 506.28
	1. 5004 2. 502.28			1. 5009 2. 506.28
	1. 5005 2. 502.28			1. 5010 2. 513.28











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5011 2. 515.28		1. 5016 2. 524.29
	1. 5012 2. 518.28		1. 5017 2. 526.29
	1. 5013 2. 518.28		1. 5018 2. 528.29
	1. 5014 2. 519.29		1. 5019 2. 531.29
	1. 5015 2. 522.29		1. 5020 2. 532.29











Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5021 2. 532.29			1. 5026 2. 538.3
	1. 5022 2. 532.29			1. 5027 2. 540.3
	1. 5023 2. 532.29			1. 5028 2. 540.3
	1. 5024 2. 537.3			1. 5029 2. 542.3
	1. 5025 2. 537.3			1. 5030 2. 542.3











Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5031 2. 542.3			1. 5036 2. 546.3
	1. 5032 2. 542.3			1. 5037 2. .546.3
	1. 5033 2. 544.3			1. 5038 2. 551.3
	1. 5034 2. 544.3			1. 5039 2. 551.3
	1. 5035 2. .546.3			1. 5040

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5041 2. 552.3		1. 5046 2. 556.31
	1. 5042 2. 552.3		1. 5047 2. 556.31
	1. 5043 2. 552.3		1. 5048 2. 556.31
	1. 5044 2. 552.3		1. 5049 2. 558.31
	1. 5045 2. 554.3		1. 5050 2. 558.31

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5051 2. 558.31			1. 5056 2. 566.31
	1. 5052 2. 560.31			1. 5057 2. 566.31
	1. 5053 2. 562.31			1. 5058 2. 568.31
	1. 5054 2. 562.31			1. 5059 2. 569.31
	1. 5055 2. 565.31			1. 5060 2. 569.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5061		1. 5066 2. 576.32
	1. 5062 2. 570.31		1. 5067 2. 576.32
	1. 5063 2. 570.31		1. 5068 2. 577.32
	1. 5064 2. 570.31		1. 5069 2. 580.32
	1. 5065 2. 572.31		1. 5070 2. 580.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5071 2. 580.32		1 5076 2. 588.32
	1. 5072 2. 580.32		1. 5077 2. 593.33
	1 5073 2. 580.32		1 5078 2. 593.33
	1 5074 2. 580.32		1. 5079 2. 594.33
	1 5075 2. 588.32		1. 5080 2. 594.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5081 2. 594.33		1. 5086 2. 602.33
	1. 5082 2. 594.33		1. 5087 2. 604.33
	1. 5083 2. 594.33		1. 5088 2. 604.33
	1. 5084 2. 595.33		1. 5089 2. 604.33
	1. 5085 2. 602.33		1. 5090 2. 604.33



















































Product	1. Ex. 2. m/z
	1. 5091 2. 616.34
	1. 5092 2. 450.25
	1. 5093 2. 518.28
	1. 5094 2. 512.28
	1. 5095 2. 554.3

TABLE 51

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5101 2. 448.8		1. 5106 2. 476.8		1. 5111 2. 490.8
	1. 5102 2. 462.8		1. 5107 2. 478.8		1. 5112 2. 494.8
	1. 5103 2. 468.8		1. 5108 2. 478.8		1. 5113 2. 498.8
	1. 5104 2. 474.7		1. 5109 2. 478.8		1. 5114 2. 503.8
	1. 5105 2. 474.7		1. 5110 2. 490.8		1. 5115 2. 504.8

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5116 2. 504.8		1. 5121 2. 514.8		1. 5126 2. 524.8
	1. 5117 2. 504.9		1. 5122 2. 514.8		1. 5127 2. 524.9
	1. 5118 2. 510.8		1. 5123 2. 519.2		1. 5128.2 2. 524.9
	1. 5119 2. 512.8		1. 5124 2. 519.2		1. 5129 2. 524.9
	1. 5120 2. 512.8		1. 5125 2. 523.8		1. 5130 2. 526.9

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5131 2. 528.8		1. 5136 2. 552.8		1. 5141 2. 567.8
	1. 5132 2. 530.9		1. 5137 2. 553.7		1. 5142 2. 574.9
	1. 5133 2. 534.8		1. 5138 2. 553.7		1. 5143 2. 576.9
	1. 5134 2. 538.9		1. 5139 2. 560.9		1. 5144 2. 576.9
	1. 5135 2. 552.8		1. 5140 2. 566.9		1. 5145 2. 514.8
























































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5146 2. 485.8		1. 5151 2. 519.2		1. 5156 2. 534.8
	1. 5147 2. 491.8		1. 5152 2. 523.8		1. 5157 2. 540.9
	1. 5148 2. 509.8		1. 5153 2. 528.8		1. 5158 2. 541.8
	1. 5149 2. 509.8		1. 5154 2. 530.9		1. 5159 2. 541.8
	1. 5150 2. 514.8		1. 5155 2. 530.9		1. 5160 2. 560.9











TABLE 52











Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5201 2. 462.8			1. 5206 2. 490.8
	1. 5202 2. 476.8			1. 5207 2. 492.8
	1. 5203 2. 482.8			1. 5208 2. 492.8
	1. 5204 2. 488.8			1. 52009 2. 492.9
	1. 5205 2. 488.8			1. 5210 2. 504.8











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5211 2. 504.8		1. 5216 2. 518.9
	2. 5212 2. 508.8		1. 5217 2. 518.9
	1. 5213 2. 512.8		1. 5218 2. 524.9
	1. 5214 2. 517.8		1. 5219 2. 526.9
	1. 5215 2. 518.9		1. 5220 2. 526.9

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5221 2. 528.8		1. 5226 2. 538.8
	1. 5222 2. 528.8		1. 5227 2. 538.9
	1. 5223 2. 533.3		1. 5228 2. 538.9
	1. 5224 2. 533.3		1. 5229 2. 538.9
	1. 5225 2. 537.9		1. 5230 2. 540.9

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5231 2. 542.9			1. 5236 2. 566.8
	1. 5232 2. 544.9			1. 5237 2. 567.7
	1. 5233 2. 548.9			1. 5238 2. 567.7
	1. 5234 2. 552.9			1. 5239 2. 574.9
	1. 5235 2. 566.8			1. 5240 2. 581

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5241 2. 533.3		1. 5246 2. 548.9
	1. 5242 2. 537.9		1. 5247 2. 554.9
	1. 5243 2. 542.8		1. 5248 2. 555.9
	1. 5244 2. 544.9		1. 5249 2. 555.9
	1. 5245 2. 544.9		1. 5250 2. 574.9

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5251 2. 579.9		1. 5256 2. 501.8
	1. 5252 2. 579.9		1. 5257 2. 514.8
	1. 5253 2. 581		1. 5258 2. 518.9
	1. 5254 2. 588.9		1. 5259 2. 530.9
	1. 5255 2. 590.9		1. 5260 2. 530.9

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5261 2. 538.9		1. 5266 2. 556.9
	1. 5262 2. 542.9		1. 5267 2. 556.9
	1. 5263 2. 547.3		1. 5268 2. 537.9
	1. 5264 2. 551.9		1. 5269 2. 566.9
	1. 5265 2. 552.9		1. 5270 2. 581







































































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5271 2. 581.7		1. 5276 2. 559.3
	1. 5272 2. 581.7		1. 5277 2. 562.9
	1. 5273 2. 588.9		1. 5278 2. 556.9
	1. 5274 2. 602.9		1. 5279 2. 499.8
	1. 5275 2. 510.8		1. 5280 2. 499.8
















TABLE 53

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5301 2. 434.7		1. 5306 2. 462.8		1. 5311 2. 476.8
	1. 5302 2. 448.8		1. 5307 2. 464.8		1. 5312 2. 480.8
	1. 5303 2. 454.8		1. 5308 2. 464.8		1. 5313 2. 484.8
	1. 5304 2. 460.7		1. 5309 2. 464.8		1. 5314 2. 489.8
	1. 5305 2. 460.7		1. 5310 2. 476.8		1. 5315 2. 490.8

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5316 2. 490.8		1. 5321 2. 500.8		1. 5326 2. 510.8
	1. 5317 2. 490.8		1. 5322 2. 500.8		1. 5327 2. 510.8
	1. 5318 2. 496.8		1. 5323 2. 505.2		1. 5328 2. 510.8
	1. 5319 2. 498.8		1. 5324 2. 505.2		1. 5329 2. 510.8
	1. 5320 2. 498.8		1. 5325 2. 509.8		1. 5330 2. 512.8

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5331 2. 514.8		1. 5336 2. 538.8		1. 5341 2. 553.8
	1. 5332 2. 516.9		1. 5337 2. 539.6		1. 5342 2. 560.9
	1. 5333 2. 520.8		1. 5338 2. 539.6		1. 5343 2. 562.9
	1. 5334 2. 524.9		1. 5339 2. 546.9		1. 5344 2. 562.9
	1. 5335 2. 538.8		1. 5340 2. 552.9		1. 5345 2. 500.8

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5346 2. 471.7		1. 5351 2. 505.2		1. 5356 2. 520.8
	1. 5347 2. 477.8		1. 5352 2. 509.8		1. 5357 2. 526.8
	1. 5348 2. 495.8		1. 5353 2. 514.8		1. 5358 2. 527.8
	1. 5349 2. 495.8		1. 5355 2. 516.9		1. 5359 2. 527.8
	1. 5350 2. 500.8		1. 5356 2. 516.9		1. 5360 2. 546.9

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5361 2. 551.8		1. 5366 2. 473.8		1. 5371 2. 510.8
	1. 5362 2. 551.8		1. 5367 2. 486.8		1. 5372 2. 514.8
	1. 5363 2. 552.9		1. 5368 2. 490.8		1. 5373 2. 519.2
	1. 5364 2. 560.9		1. 5369 2. 502.8		1. 5374 2. 523.8
	1. 5365 2. 562.9		1. 5370 2. 502.8		1. 5375 2. 524.8












































































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5376 2. 528.8		1. 5381 2. 553.7		1. 5386 2. 531.2
	1. 5377 2. 528.8		1. 5382 2. 553.7		1. 5387 2. 534.8
	1. 5378 2. 509.8		1. 5383 2. 560.9		1. 5388 2. 528.8
	1. 5379 2. 538.9		1. 5384 2. 574.9		1. 5389 2. 538.8
	1. 5380 2. 552.9		1. 5385 2. 482.8		1. 5390 2. 471.7
















TABLE 54

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5401 2. 463.3		1. 5406 2. 491.3		1. 5411 2. 505.3
	1. 5402 2. 477.3		1. 5407 2. 493.3		1. 5412 2. 509.3
	1. 5403 2. 483.3		1. 5408 2. 493.3		1. 5413 2. 513.3
	1. 5404 2. 489.3		1. 5409 2. 493.3		1. 5414 2. 518.3
	1. 5405 2. 489.3		1. 5410 2. 505.3		1. 5415 2. 519.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5416 2. 519.3		1. 5121 2. 529.3		1. 5426 2. 539.3
	1. 5417 2. 519.3		1. 5422 2. 529.3		1. 5427 2. 539.3
	1. 5418 2. 525.3		1. 5423 2. 533.3		1. 5428 2. 539.3
	1. 5419 2. 527.3		1. 5424 2. 533.3		1. 5429 2. 539.3
	1. 5420 2. 527.3		1. 5425 2. 538.3		1. 5430 2. 541.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5431 2. 543.3		1. 5436 2. 567.3		1. 5441 2. 582.3
	1. 5432 2. 545.3		1. 5437 2. 567.3		1. 5442 2. 589.3
	1. 5433 2. 549.3		1. 5438 2. 567.3		1. 5443 2. 591.3
	1. 5434 2. 553.3		1. 5439 2. 575.3		1. 5444 2. 591.3
	1. 5435 2. 567.3		1. 5440 2. 581.3		1. 5445 2. 529.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5446 2. 500.3		1. 5451 2. 533.3		1. 5456 2. 549.3
	1. 5447 2. 830.5		1. 54525 2. 719.4		1. 5457 2. 555.3
	1. 5448 2. 524.3		1. 5453 2. 543.3		1. 5458 2. 556.3
	1. 5449 2. 524.3		1. 5454 2. 545.3		1. 5459 2. 556.3
	1. 5450 2. 529.3		1. 5455 2. 545.3		1. 5460 2. 575.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5461 2. 580.3		1. 5466 2. 502.3		1. 5471 2. 539.3
	1. 5462 2. 580.3		1. 5467 2. 515.3		1. 5472 2. 543.3
	1. 5463 2. 581.3		1. 5468 2. 519.3		1. 5473 2. 547.3
	1. 5464 2. 589.3		1. 5469 2. 531.3		1. 5474 2. 552.3
	1. 5465 2. 591.3		1. 5470 2. 531.3		1. 5475 2. 553.3





























































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5476 2. 557.3		1. 5481 2. 581.3		1. 5486 2. 581.3
	1. 5477 2. 557.3		1. 5482 2. 589.3		1. 5487 2. 891.5
	1. 5478 2. 538.3		1. 5483 2. 511.3		1. 5488 2. 500.3
	1. 5479 2. 567.3		1. 5484 2. 559.3		1. 5489 2. 500.3
	1. 5480 2. 581.3		1. 5485 2. 563.3		1. 5490 2. 501.3

TABLE 55

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5501 2. 488.81		1. 5506 2. 516.87		1. 5511 2. 530.87
	1. 5502 2. 502.84		1. 5507 2. 518.84		1. 5512 2. 534.84
	1. 5503 2. 508.87		1. 5508 2. 518.84		1. 5513 2. 538.87
	1. 5504 2. 514.81		1. 5509 2. 518.88		1. 5514 2. 543.85
	1. 5505 2. 514.81		1. 5510 2. 530.87		1. 5515 2. 544.9

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5516 2. 544.9		1. 5521 2. 554.87		1. 5526 2. 564.87
	1. 5517 2. 544.92		1. 5522 2. 554.87		1. 5527 2. 564.91
	1. 5518 2. 550.89		1. 5523 2. 559.29		1. 5528 2. 564.91
	1. 5519 2. 552.9		1. 5524 2. 559.29		1. 5529 2. 564.91
	1. 5520 2. 552.9		1. 5525 2. 563.88		1. 5530 2. 566.93

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5531 2. 568.9		1. 5536 2. 600.95		1. 5541
	1. 5532 2. 574.91		1. 5537 2. 606.99		1. 5542 2. 525.83
	1. 5533 2. 578.94		1. 5538 2. 614.97		1. 5543 2. 581.9
	1. 5534 2. 592.85		1. 5539 2. 616.95		1. 5544 2. 581.9
	1. 5535 2. 592.85		1. 5540 2. 616.95		1. 5545 2. 605.92

















































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5546 2. 573.32		1. 5551 2. 525.83
	1. 5547 2. 582.88		1. 5552 2. 526.82
	1. 5548 2. 607.76		1. 5553 2. 573.92
	1. 5549 2. 607.76		
	1. 5550 2. 536.86		











TABLE 56

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5601 2. 481.26			1. 5606 2. 548.3
	1. 5602 2. 495.27			1. 5607 2. 554.3
	1. 5603 2. 535.29			1. 5608 2. 554.3
	1. 5604 2. 543.3			1. 5609 2. 555.31
	1. 5605 2. 543.3			1. 5610 2. 559.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5611 2. 565.31		1. 5616 2. 579.32
	1. 5612 2. 565.31		1. 5617
	1. 5613 2. 565.31		1. 5618
	1. 5614 2. 569.31		1. 5619
	1. 5615 2. 571.31		1. 5620






Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5621 2. 615.34			1. 5626 2. 547.3
	2. 5622 2. 680.37			1. 5627 2. 547.3
	1. 5623 2. 529.29			1. 5628 2. 554.3
	1. 5624 2. 543.3			1. 5629 2. 555.31
	1. 5625 2. 547.3			1. 5630 2. 559.31











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5631 2. 561.31		1. 5636 2. 565.31
	1. 5632 2. 563.31		1. 5637 2. 579.32
	1. 5633 2. 563.31		1. 5638 2. 581.32
	1. 5634 2. 563.31		1. 5639 2. 581.32
	1. 5635 2. 563.31		1. 5640 2. 585.32











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5641 2. 587.32		1. 5646
	1. 5642 2. 589.32		1. 5647 2. 597.33
	1. 5643 2. 589.32		1. 5648 2. 597.33
	1. 5644 2. 593.33		1. 5649 2. 597.33
	1. 5645 2. 597.33		1. 5650 2. 597.33











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5651 2. 597.33		1. 5656 2. 607.33
	1. 5652 2. 599.33		1. 5657 2. 607.33
	1. 5653 2. 603.33		1. 5658 2. 607.33
	1. 5654 2. 605.33		1. 5659 2. 621.34
	1. 5655 2. 607.33		1. 5660 2. 631.35

438

Product	1. Ex. 2. m/z
	1. 5661 2. 631.35
	1. 5662 2. 631.35
	1. 5663 2. 642.35
	1. 5664 2. 665.37
	1. 5665 2. 571.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5711 2. 559.31		1. 5716 2. 563.31
	1. 5712 2. 559.31		1. 5717 2. 565.31
	1. 5713 2. 561.31		1. 5718 2. 565.31
	1. 5714 2. 562.31		1. 5719 2. 565.31
	1. 5715 2. 563.31		1. 5720 2. 565.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5721 2. 569.31		1. 5726 2. 581.32
	1. 5722 2. 571.31		1. 5727 2. 585.32
	1. 5723 2. 579.32		1. 5728 2. 587.32
	1. 5724 2. 579.32		1. 5729 2. 589.32
	1. 5725 2. 581.32		1. 5730 2. 589.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5731 2. 591.33		1. 5736 2. 597.33
	1. 5732 2. 597.33		1. 5737 2. 597.33
	1. 5733 2. 595.33		1. 5738 2. 596.33
	1. 5734 2. 597.33		1. 5739 2. 597.33
	1. 5735 2. 597.33		1. 5740 2. 581.32









































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5741 2. 605.33		1. 5746 2. 613.34
	1. 5742 2. 607.33		1. 5747 2. 615.34
	1. 5743 2. 607.33		1. 5748 2. 621.34
	1. 5744 2. 607.33		1. 5749 2. 631.35
	1. 5745 2. 613.34		1. 5750 2. 680.37

TABLE 58

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5801 2. 550.3		1. 5806 2. 598.33
	1. 5802 2. 590.32		10 5807 2. 602.33
	1. 5803 2. 598.33		1. 5808 2. 602.33
	1. 5804 2. 598.33		1. 5809 2. 603.33
	1. 5805 2. 598.33		1. 5810 2. 609.33

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 5811 2. 609.33			1. 5816 2. 618.34
	1. 5812 2. 610.34			1. 5817 2. 624.34
	1. 5813 2. 612.34			1. 5818 2. 626.34
	1. 5814 2. 614.34			1. 5819 2. 634.35
	1. 5815 2. 614.34			1. 5820 2. 632.35

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5821 2. 636.35		1. 5826 2. 648.36
	1. 5822 2. 640.35		1. 5827 2. 654.36
	1. 5823 2. 642.35		1. 5828 2. 660.36
	1. 5824 2. 644.35		1. 5829 2. 670.37
	1. 5825 2. 644.35		1. 5830 2. 676.37



















Product	1. Ex. 2. m/z
	1. 5831 2. 735.4
	1. 5832 2. 522.29
	1. 5833 2. 584.32

TABLE 59

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5901 2. 536.29		1. 5906 2. 598.33		1. 5911 2. 603.33
	1. 5902 2. 550.3		1. 5907 2. 598.33		1. 5912 2. 609.33
	1. 5903 2. 590.32		1. 5908 2. 602.33		1. 5913 2. 609.33
	1. 5904 2. 598.33		1. 5909 2. 602.33		1. 5914 2. 610.34
	1. 5905 2. 598.33		1. 5910 2. 602.33		1. 5915 2. 612.34





























































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 5916 2. 614.34		1. 5921 2. 618.34		1. 5926 2. 620.34
	1. 5917 2. 614.34		1. 5922 2. 620.34		1. 5927 2. 624.34
	1. 5918 2. 616.34		1. 5923 2. 620.34		1. 5928 2. 626.34
	1. 5919 2. 618.34		1. 5924 2. 620.34		1. 5929 2. 626.34
	1. 5920 2. 618.34		1. 5925 2. 620.34		1. 5930 2. 632.35

TABLE 60

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6001 2. 536.29		1. 6006 2. 588.32		1. 6011 2. 595.33
	1. 6002 2. 576.32		1. 6007 2. 588.32		1. 6012 2. 595.33
	1. 6003 2. 584.32		1. 6008 2. 588.32		1. 6013 2. 596.33
	1. 6004 2. 584.32		1. 6009 2. 589.32		1. 6014 2. 598.33
	1. 6005 2. 584.32		1. 6010 2. 595.33		1. 6015 2. 600.33

451

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6016 2. 600.33		1. 6021 2. 606.33		1. 6026 2. 612.34
	1. 6017 2. 602.33		1. 6022 2. 606.33		1. 6027 2. 620.34
	1. 6018 2. 604.33		1. 6023 2. 606.33		1. 6028 2. 620.34
	1. 6019 2. 604.33		1. 6024 2. 606.33		1. 6029 2. 622.34
	1. 6020 2. 604.33		1. 6025 2. 606.33		1. 6030 2. 626.34

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6031 2. 628.35		1. 6036 2. 638.35		1. 6041 2. 648.36
	1. 6032 2. 630.35		1. 6037 2. 638.35		1. 60425 2. 648.36
	1. 6033 2. 630.35		1. 6038 2. 638.35		1. 6043 2. 648.36
	1. 6034 2. 634.35		1. 6039 2. 622.34		1. 6044 2. 648.36
	1. 6035 2. 638.35		1. 6040 2. 646.36		1. 6045 2. 654.36

453
















Product	1. Ex. 2. m/z
	1. 6046 2. 654.36
	1. 6047 2. 656.36
	1. 6048 2. 662.36
	1. 6049 2. 721.4
	1. 6050 2. 570.31

TABLE 61

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6101 2. 584.32		1. 6106 2. 588.32
	1. 6102 2. 584.32		1. 6107 2. 595.33
	1. 6103 2. 584.32		1. 6108 2. 600.33
	1. 6104 2. 588.32		1. 6109 2. 602.33
	1. 6105 2. 588.32		1. 6110 2. 604.33
























































Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 6111 2. 604.33			1. 6116 2. 606.33
	1. 6112 2. 604.33			1. 6117 2. 606.33
	1. 6113 2. 606.33			1. 6118 2. 648.36
	1. 6114 2. 606.33			1. 6119 2. 721.4
	1. 6115 2. 606.33			1. 6120 2. 570.31

TABLE 62

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6201 2. 514.28		1. 6206 2. 576.32		1. 6211 2. 581.32
	1. 6202 2. 528.29		1. 6207 2. 576.32		1. 6212 2. 587.32
	1. 6203 2. 568.31		1. 6208 2. 580.32		1. 6213 2. 587.32
	1. 6204 2. 576.32		1. 6209 2. 580.32		1. 6214 2. 587.32
	1. 6205 2. 576.32		1. 6210 2. 580.32		1. 6215 2. 588.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6216 2. 590.32		1. 6221 2. 596.33		1. 6226 2. 598.33
	1. 6217 2. 592.33		1. 6222 2. 596.33		1. 6227 2. 596.33
	1. 6218 2. 592.33		1. 6223 2. 598.33		1. 6228 2. 602.33
	1. 6219 2. 592.33		1. 6224 2. 596.33		1. 6229 2. 604.33
	1. 6220 2. 594.33		1. 6225 2. 598.33		1. 6230 2. 604.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6231 2. 610.34		1. 6236 2. 614.34		1. 6241 2. 630.35
	1. 6232 2. 610.34		1. 6237 2. 618.34		1. 6242 2.
	1. 6233 2. 612.34		1. 6238 2. 622.34		1. 6243 2. 630.35
	1. 6234 2. 612.34		1. 6239 2. 622.34		1. 6244 2. 630.35
	1. 6235 2. 614.34		1. 6240 2. 626.34		1. 6245 2. 630.35

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6246 2. 630.35		1. 6251 2. 636.35		1. 6256 2. 646.36
	1. 6247 2. 630.35		1. 6252 2. 638.35		1. 6257 2. 646.36
	1. 6248 2. 630.35		1. 6253 2. 640.35		1. 6258 2. 648.36
	1. 6249 2. 630.35		1. 6254 2. 640.35		1. 6259 2. 654.36
	1. 6250 2. 632.35		1. 6255 2. 638.35		1. 6260 2. 713.39

















































Product	1. Ex. 2. m/z
	1. 6261 2. 500.27
	1. 6262 2. 562.31
	1. 6263 2. 604.33

TABLE 63

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6301 2. 500.27		1. 6306 2. 562.31		1. 6311 2. 567.31
	1. 6302 2. 514.28		1. 6307 2. 562.31		1. 6312 2. 573.32
	1. 6303 2. 554.3		1. 6308 2. 566.31		1. 6313 2. 573.32
	1. 6304 2. 562.31		1. 6309 2. 566.31		1. 6314 2. 573.32
	1. 6305 2. 562.31		1. 6310 2. 566.31		1. 6315 2. 574.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6316 2. 576.32		1. 6321 2. 582.32		1. 6326 2. 584.32
	1. 6317 2. 578.32		1. 6322 2. 582.32		1. 6327 2. 584.32
	1. 6318 2. 578.32		1. 6323 2. 584.32		1. 6328 2. 588.32
	1. 6319 2. 580.32		1. 6324 2. 582.32		1. 6329 2. 590.32
	1. 6320 2. 582.32		1. 6325 2. 584.32		1. 6330 2. 596.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6331 2. 598.33		1. 6336 2. 606.33		1. 6341 2. 616.34
	1. 6332 2. 598.33		1. 6337 2. 608.33		1. 6342 2. 616.34
	1. 6333 2. 600.33		1. 6338 2. 608.33		1. 6343 2. 616.34
	1. 6334 2. 600.33		1. 6339 2. 612.34		1.6344 2. 616.34
	1. 6335 2. 604.33		1. 6340 2. 616.34		1. 6345 2. 616.34

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6346 2. 616.34		1. 6351 2. 622.34		1. 6356 2. 626.34
	1. 6347 2. 616.34		1. 6352 2. 624.34		1. 6357 2. 632.35
	1. 6348 2. 616.34		1. 6353 2. 626.34		1. 6358 2. 632.35
	1. 6349 2. 618.34		1. 6354 2. 626.34		1. 6359 2. 634.35
	1. 6350 2. 600.33		1. 6355 2. 626.34		1. 6360 2. 640.35





































































Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 6361 2. 650.36			1. 6366 2. 486.27
	1. 6362 2. 650.36			1. 6367 2. 548.3
	1. 6363 2. 661.36			1. 6368 2. 590.32
	1. 6364 2. 684.38			
	1. 6365 2. 699.38			

TABLE 64

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6401 2. 474.26		1. 6406 2. 536.29		1. 6411 2. 541.3
	1. 6402 2. 488.27		1. 6407 2. 536.29		1. 6412 2. 547.3
	1. 6403 2. 528.29		1. 6408 2. 540.3		1. 6413 2. 547.3
	1. 6404 2. 536.29		1. 6409 2. 540.3		1. 6414 2. 548.3
	1. 6405 2. 536.29		1. 6410 2. 540.3		1. 6415 2. 550.3
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6416 2. 552.3		1. 6421 2. 556.31		1. 6426 2. 558.31
	1. 6417 2. 552.3		1. 6122 2. 558.31		1. 6427 2. 564.31
	1. 6418 2. 554.3		1. 6423 2. 558.31		1. 6428 2. 572.31
	1. 6419 2. 556.31		1. 6424 2. 558.31		1. 6429 2. 572.31
	1. 6420 2. 556.31		1. 6425 2. 558.31		1. 6430 2. 574.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6431 2. 574.32		1. 6436 2. 586.32		1. 6441 2. 590.32
	1. 6432 2. 578.32		1. 6437 2. 590.32		1. 6442 2. 590.32
	1. 6433 2. 580.32		1. 6438 2. 590.32		1. 6443 2. 590.32
	1. 6434 2. 582.32		1. 6439 2. 590.32		1. 6444 2. 590.32
	1. 6435 2. 582.32		1. 6440 2. 589.32		1. 6445 2. 590.32
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6446 2. 592.33		1. 6451 2. 600.33		1. 6456 2. 614.34
	1. 6447 2. 574.32		1. 6452 2. 600.33		1. 6447 2. 673.37
	1. 6448 2. 598.33		1. 6453 2. 606.33		1. 6458 2. 460.25
	1. 6449 2. 600.33		1. 6454 2. 606.33		6459 2.
	1. 6450 2. 600.33		1. 6455 2. 608.33		1. 6460 2. 564.31
















470














TABLE 65

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6501 2. 488.27		1. 6506 2. 550.3		1. 6511 2. 561.31
	1. 6502 2. 502.28		1. 6507 2. 554.3		1. 6512 2. 561.31
	1. 6503 2. 542.3		1. 6508 2. 554.3		1. 6213 2. 561.31
	1. 6504 2. 550.3		1. 6509 2. 554.3		1. 6514 2. 562.31
	1. 6505 2. 550.3		1. 6510 2. 555.31		1. 6515 2. 564.31

471
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6516 2. 566.31		1. 6521 2. 570.31		1. 6526 2. 576.32
	1. 6517 2. 566.31		1. 6522 2. 572.31		1. 6527 2. 578.32
	1. 6518 2. 568.31		1. 6523 2. 572.31		1. 6528 2. 586.32
	1. 6519 2. 570.31		1. 6524 2. 572.31		1. 6529 2. 586.32
	1. 6520 2. 570.31		1. 6525 2. 572.31		1. 6530 2. 588.32
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6531 2. 588.32		1. 6536 2. 600.33		1. 6541 2. 604.33
	1. 6532 2. 592.33		1. 6537 2. 604.33		1. 6542 2. 604.33
	1. 6533 2. 594.33		1. 6538 2. 604.33		1. 6543 2. 604.33
	1. 6534 2. 596.33		1. 6539 2. 604.33		1. 6544 2. 606.33
	1. 6535 2. 596.33		1. 6540 2. 604.33		1. 6545 2. 588.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6546 2. 612.34		1. 6551 2. 620.34		1. 6556 2. 474.26
	1. 6547 2. 614.34		1. 6552 2. 620.34		1. 6557 2. 536.29
	1. 6548 2. 614.34		1. 6553 2. 622.34		1. 6558 2. 578.32
	1. 6549 2. 614.34		1. 6554 2. 628.35		
	1. 6550 2. 614.34		1. 6555 2. 687.38		

474

TABLE 67

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6701 2. 474.26		1. 6706 2. 526.29		1. 6711 2. 536.29
	1. 6702 2. 514.28		1. 6707 2. 526.29		1. 6712 2. 538.3
	1. 6703 2. 522.29		1. 6708 2. 526.29		1. 6713 2. 538.3
	1. 6704 2. 522.29		1. 6709 2. 533.29		1. 6714 2.
	1. 6705 2. 522.29		1. 6710 2. 534.29		1. 6715 2. 542.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6716 2. 542.3		1. 6721 2. 550.3		1. 6726 2. 564.31
	1. 6717 2. 542.3		1. 6722 2. 558.31		1. 6727 2. 566.31
	1. 6718 2. 544.3		1. 6723 2. 558.31		1. 6728 2. 568.31
	1. 6719 2. 544.3		1. 6724 2. 560.31		1. 6729 2. 568.31
	1. 6720 2. 544.3		1. 6725 2. 560.31		1. 6730 2. 572.31

476
























































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6731 2. 576.32		1. 6736 2. 576.32		1. 6741 2. 586.32
	1. 6732 2. 576.32		1. 6737 2. 576.32		1. 6742 2. 586.32
	1. 6733 2. 576.32		1. 6738 2. 576.32		1. 6743 2. 592.33
	1. 6734 2. 576.32		1. 6739 2. 584.32		1. 6744 2. 592.33
	1. 6735 2. 576.32		1. 6740 2. 586.32		1. 6745 2. 600.33











TABLE 68

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 6801 2. 488.27			1. 6806 2. 550.3
	1. 6802 2. 502.28			1. 6807 2. 554.3
	1. 6803 2. 542.3			1. 6808 2. 554.3
	1. 6804 2. 550.3			1. 6809 2. 554.3
	1. 68058 2. 550.3			1. 6810 2. 555.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6811 2. 561.31		1. 6816 2. 566.31
	1. 6812 2. 561.31		1. 6817 2. 568.31
	1. 6813 2. 562.31		1. 6818 2. 570.31
	1. 6814 2. 564.31		1. 6819 2. 570.31
	1. 6815 2. 566.31		1. 6820 2. 570.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6821 2. 571.31		1. 6826 2. 576.32
	1. 6822 2. 572.31		1. 6827 2. 578.32
	1. 6823 2. 572.31		1. 6828 2. 586.32
	1. 6824 2. 572.31		1. 6829 2. 586.32
	1. 6825 2. 572.31		1. 6830 2. 588.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6831 2. 588.32		1. 6836 2. 600.33
	1. 6832 2. 592.33		1. 6837 2. 602.33
	1. 6833 2. 614.34		1. 6838 2. 604.33
	1. 6834 2. 596.33		1. 6839 2. 604.33
	1. 6835 2. 596.33		1. 6840 2. 604.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6841 2. 604.33		1. 6846 2. 606.33
	1. 6842 2. 604.33		1. 6847 2. 588.32
	1. 6843 2. 605.33		1. 6848 2. 610.34
	1. 6844 2. 604.33		1. 6849 2. 612.34
	1. 6845 2. 604.33		1. 6850 2. 594.33



















































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6851 2. 614.34		1. 6856 2. 620.34
	1. 6852 2. 614.34		1. 6856 2. 628.35
	1. 6853 2. 614.34		1. 6857 2. 687.38
	1. 6854 2. 620.34		1. 6859 2. 474.26
	1. 6855 2. 620.34		1. 6860 2. 536.29











TABLE 69

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6901 2. 514.28		1. 6906 2. 580.32
	1. 6902 2. 568.31		1. 6907 2. 580.32
	1. 6903 2. 576.32		1. 6908 2. 580.32
	1. 6904 2. 576.32		1. 6909 2. 581.32
	1. 6905 2. 576.32		1. 6910 2. 587.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6911 2. 587.32		1. 6916 2. 594.33
	1. 6912 2. 588.32		1. 6917 2. 596.33
	1. 6913 2. 590.32		1. 6918 2. 596.33
	1. 6914 2. 592.33		1. 6919 2. 596.33
	1. 6915 2. 592.33		1. 6920 2. 598.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6921 2. 598.33		1. 6926 2. 604.33
	1. 6922 2. 598.33		1. 6927 2. 612.34
	1. 6923 2. 598.33		1. 6928 2. 612.34
	1. 6924 2. 598.33		1. 6929 2. 614.34
	1. 6925 2. 602.33		1. 6930 2. 614.34

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6931 2. 618.34		1. 6936 2. 630.35
	1. 6932 2. 620.34		1. 6937 2. 630.35
	1. 6933 2. 622.34		1. 6938 2. 630.35
	1. 6934 2. 622.34		1. 6939 2. 630.35
	1. 6935 2. 626.34		1. 6940 2. 630.35

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 6941 2. 630.35			1. 6946 2. 636.35
	1. 6942 2. 630.35			1. 6947 2. 638.35
	1. 6943 2. 630.35			1. 6948 2. 640.35
	1. 6944 2. 632.35			1. 6949 2. 640.35
	1. 6945 2. 614.34			1. 6950 2. 640.35









































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 6951 2. 640.35		1. 6956 2. 675.23
	1. 6952 2. 646.36		1. 6957 2. 713.39
	1. 6953 2. 646.36		1. 6958 2. 500.27
	1. 6954 2. 648.36		1. 6959 2. 562.31
	1. 6955 2. 654.36		1. 6960 2. 604.33

TABLE 70

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7001 2. 472.26		1. 7006 2. 522.29
	1. 7002 2. 488.27		1. 7007 2. 520.29
	1. 7003 2. 488.27		1. 7008 2. 520.29
	1. 7004 2. 500.27		1. 7009 2. 526.29
	1. 7005 2. 520.29		1. 7010 2. 533.29

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7011 2. 536.29		1. 7016 2. 538.3
	1. 7012 2. 536.29		1. 7017 2. 540.3
	1. 7013 2. 536.29		1. 7018 2. 542.3
	1. 7014 2. 536.29		1. 7019 2. 544.3
	1. 7015 2. 536.29		1. 7020 2. 544.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7021 2. 548.3		1. 7026 2. 566.31
	1. 7022 2. 552.3		1. 7027 2. 568.31
	1. 7023 2. 552.3		1. 7028 2. 572.31
	1. 7024 2. 556.31		1. 7029 2. 576.32
	1. 7025 2. 558.31		1. 7030 2. 576.32





























Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7031 2. 576.32		1. 7036 2. 598.33
	1. 7032 2. 576.32		1. 7037 2. 598.33
	1. 7033 2. 590.32		1. 7038 2. 610.34
	1. 7034 2. 486.27		
	1. 7035 2. 594.33		

TABLE 71

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7101 2. 474.26		1. 7106 2. 522.29
	1. 7102 2. 488.27		1. 7107 2. 522.29
	1. 7103 2. 500.27		1. 7108 2. 526.3
	1. 7104 2. 522.29		1. 7109 2. 533.29
	1. 7105 2. 522.29	 Chiral	1. 7110 2. 536.29

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7111 2. 536.29		1. 7116 2. 538.3
	1. 7112 2. 536.29		1. 7117 2. 540.3
	1. 7113 2. 536.29		1. 7118 2. 542.3
	1. 7114 2. 536.29		1. 7119 2. 542.3
	1. 7115 2. 536.3		1. 7120 2. 542.3

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7121 2. 546.3			1. 7126 2. 558.31
	1. 7122 2. 550.3			1. 7127 2. 550.3
	1. 7123 2. 552.3			1. 7128 2. 566.31
	1. 7124 2. 552.3			1. 7129 2. 565.31
	1. 7125 2. 556.31			1. 7130 2. 572.31


















































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7131 2. 576.32		1. 7136 2. 590.32
	1. 7132 2. 576.32		1. 7137 2. 590.32
	1. 7133 2. 576.32		1. 7138 2. 598.33
	1. 7134 2. 584.32		1. 7139 2. 598.33
	1. 7135 2. 585.3		

TABLE 72

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7201 2. 529.29			1. 7206 2. 577.32
	1. 7202 2. 543.3			1. 7207 2. 577.32
	1. 7203 2. 543.3			1. 7208 2. 577.32
	1. 7204 2. 555.31			1. 7209 2. 581.32
	1. 7205 2. 577.32			1. 7210 2. 581.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7211 2. 588.32		1. 7216 2. 591.33
	1. 7212 2. 588.32		1. 7217 2. 593.33
 <i>Optical</i>	1. 7213 2. 591.33		1. 7218 2. 593.33
 <i>Optical</i>	1. 7214 2. 591.33		1. 7219 2. 593.33
	1. 7215 2. 591.33		1. 7220 2. 595.33

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7221 2. 597.33		1. 7226 2. 603.33
	1. 7222 2. 597.33		1. 7227 2. 603.33
	1. 7223 2. 597.33		1. 7228 2. 607.33
	1. 7224 2. 599.33		1. 7229 2. 607.14
	1. 7225 2. 599.33		1. 7230 2. 611.34

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7231 2. 613.34			1. 7236 2. 631.35
	1. 7232 2. 605.33			1. 7237 2. 631.35
	1. 7233 2. 621.34			1. 7238 2. 631.35
	1. 7234 2. 623.34			1. 7239 2. 631.35
	1. 7235 2. 627.34			1. 7240 2. 631.35






































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7241 2. 639.35		1. 7246 2. 653.36
	1. 7242 2. 641.35		1. 7247 2. 665.37
	1. 7243 2. 645.35		
	1. 7244 2. 541.3		
	1. 7245 2. 649.36		

TABLE 73

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7301 2. 529.29		1. 7306 2. 567.31		1. 7311 2. 577.32
	1. 7302 2. 541.3		1. 7307 2. 567.31		1. 7312 2. 577.32
	1. 7303 2. 563.13		1. 7308 2. 574.32		1. 7313 2. 579.32
	1. 7304 2. 563.31		1. 7309 2. 574.32		1. 7314 2. 579.32
	1. 7305 2. 563.31		1. 7310 2. 577.32		1. 7315 2. 579.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7316 2. 581.32		1. 7321 2. 585.32		1. 7326 2. 599.33
	1. 7317 2. 583.32		1. 7322 2. 589.32		1. 7327 2. 591.33
	1. 7318 2. 583.32		1. 7323 2. 591.33		1. 7328
	1. 7319 2. 583.32		1. 7324 2. 593.33		1. 7329 2. 613.34
	1. 7320 2. 585.32		1. 7325 2. 597.33		1. 7330 2. 617.34












































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7331 2. 599.33		1. 7336 2. 617.34		1. 73412. 635.35
	1. 7332 2. 591.33		1. 7337 2. 617.34		1. 7342 2. 639.35
	1. 7333 2. 609.33		1. 7338 2. 617.34		1. 7343 2. 639.35
	1. 7334 2. 613.34		1. 7339 2. 625.34		
	1. 7335 2. 617.34		1. 7340 2. 631.35		

TABLE 74

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7401 2. 515.28			1. 7406 2. 563.31
	1. 7402 2. 529.29			1. 7407 2. 563.31
	1. 7403 2. 541.3			1. 7408 2. 567.31
	1. 7404 2. 563.13			1. 7409 2. 567.31
	1. 7405 2. 563.31			1. 7410 2. 574.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7411 2. 574.32		1. 7416 2. 579.32
	1. 7412 2. 577.14		1. 7417 2. 579.32
	1. 7413 2. 577.32		1. 7418 2. 579.32
	1. 7414 2. 577.32		1. 7419 2. 581.32
	1. 7415 2. 577.32		1. 7420 2. 583.32

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7421 2. 583.32			1. 7426 2. 593.33
	1. 7422 2. 585.32			1. 7427 2. 597.33
	1. 7423 2. 585.32			1. 7428 2. 591.33
	1. 7424 2. 589.32			1. 7429 2. 609.33
	1. 7425 2. 591.33			1. 7430 2. 613.34




















































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7431 2. 617.34		1. 7436 2. 639.35
	1. 7432 2. 617.34		1. 7437 2. 639.35
	1. 7433 2. 617.34		1. 7438 2. 651.36
	1. 7434 2. 627.34		
	1. 7435 2. 631.35		

TABLE 75

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7501 2. 481.26		1. 7506 2. 529.29		1. 7511 2. 540.3
	1. 7502 2. 495.27		1. 7507 2. 529.29		1. 7512 2. 540.3
	1. 7503 2. 495.27		1. 7508 2. 529.29		1. 7513 2. 543.3
	1. 7504 2. 507.28		1. 7509 2. 533.29		1. 7514 2. 543.3
	1. 7505 2. 529.29		1. 7510 2. 533.29		1. 7515 2. 543.3
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7516 2. 543.3		1. 7521 2. 549.3		1. 7526 2. 555.31
	1. 7517 2. 545.3		1. 7522 2. 549.3		1. 7527 2. 557.31
	1. 7518 2. 545.3		1. 7523 2. 549.3		1. 7528 2. 559.31
	1. 7519 2. 545.3		1. 7524 2. 551.3		1. 7529 2. 560.31
	1. 7520 2. 547.3		1. 7525 2. 551.3		1. 7530 2. 573.32
















511













Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7531 2. 579.32		1. 7536 2. 583.32		1. 7541 2. 605.33
	1. 7532 2. 583.32		1. 7537 2. 591.33		1. 7542 2. 605.33
	1. 7533 2. 583.32		1. 7538 2. 593.33		1. 7543 2. 614.34
	1. 7534 2. 583.32		1. 7539 2. 597.33		
	1. 7535 2. 583.32		1. 7540 2. 601.33		

512

TABLE 76

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7601 2. 453.25		1. 7606 2. 501.28		1. 7611 2. 512.28
	1. 7602 2. 467.26		1. 7607 2. 501.28		1. 7612 2. 512.28
	1. 7603 2. 467.26		1. 7608 2. 501.28		1. 7613 2. 515.28
	1. 7604 2. 479.26		1. 7609 2. 505.28		1. 7614 2. 515.28
	1. 7605 2. 501.28		1. 7610 2. 505.28		1. 7615 2. 515.28

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7616 2. 515.28		1. 7621 2. 521.29		1. 7626 2. 527.29
	1. 7617 2. 517.28		1. 7622 2. 521.29		1. 7627 2. 529.29
	1. 7618 2. 517.28		1. 7623 2. 521.29		1. 7628 2. 541.3
	1. 7619 2. 517.28		1. 7624 2. 523.29		1. 7629 2. 545.3
	1. 7620 2. 519.29		1. 7625 2. 523.29		1. 7630 2. 547.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7631 2. 551.3		1. 7636 2. 555.31		1. 7641 2. 577.32
	1. 7632 2. 555.31		1. 7637 2. 563.31		1. 7642 2. 577.32
	1. 7633 2. 555.31		1. 7638 2. 565.31		
	1. 7634 2. 555.31		1. 7639 2. 569.31		
	1. 7635 2. 555.31		1. 7640 2. 573.32		

515

TABLE 77









































Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7701 2. 481.26			1. 7706 2. 529.29
	1. 7702 2. 495.27			1. 7707 2. 529.29
	1. 7703 2. 495.27			1. 7708 2. 529.29
	1. 7704 2. 507.28			1. 7709 2. 533.29
	1. 7705 2. 529.29			1. 7710 2. 533.29











TABLE 78

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7801 2. 507.28			1. 7806 2. 555.31
	1. 7802 2. 521.29			1. 7807 2. 555.31
	1. 7803 2. 521.29			1. 7808 2. 555.31
	1. 7804 2. 533.29			1. 7809 2. 559.31
	1. 7805 2. 555.31			1. 7810 2. 559.31

517

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7811 2. 566.31		1. 7816 2. 569.31
	1. 7812 2. 566.31		1. 7817 2. 571.31
	1. 7813 2. 569.31		1. 7818 2. 571.31
	1. 7814 2. 569.31		1. 7819 2. 571.31
	1. 7815 2. 569.31		1. 7820 2. 573.32

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7821 2. 575.32		1. 7826 2. 581.32
	1. 7822 2. 575.32		1. 7827 2. 583.32
	1. 7823 2. 575.32		1. 7828 2. 585.32
	1. 7824 2. 577.32		1. 7829 2. 585.32
	1. 7825 2. 577.32		1. 7830 2. 589.32

Product	1. Ex. 2. m/z.	Product	1. Ex. 2. m/z.
	1. 7831 2. 591.33		1. 7836 2. 609.33
	1. 7832 2. 599.33		1. 7837 2. 609.33
	1. 7833 2. 601.33		1. 7838 2. 609.33
	1. 7834 2. 605.33		1. 7839 2. 609.33
	1. 7835 2. 609.33		1. 7840 2. 617.34
















































Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7841 2. 619.34			1. 7846 2. 631.35
	1. 7842 2. 623.34			1. 7847 2. 643.35
	1. 7843 2. 519.29			
	1. 7844 2. 627.34			
	1. 7845 2. 631.35			











TABLE 79

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7901 2. 477.26		1. 7906 2. 504.28
	1. 7902 2. 485.27		1. 7907 2. 505.28
	1. 7903 2. 485.27		1. 7908 2. 507.28
	1. 7904 2. 486.27		1. 7909 2. 509.28
	1. 7905 2. 495.27		1. 7910 2. 509.28

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7911 2. 509.28		1. 7916 2. 523.29
	1. 7912 2. 513.28		1. 7917 2. 525.29
	1. 7913 2. 515.28		1. 7918 2. 529.29
	1. 7914 2. 519.29		1. 7919 2. 529.29
	1. 7915 2. 523.29		1. 7920 2. 530.29

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7921 2. 530.29			1. 7926 2. 547.3
	1. 7922 2. 535.29			1. 7927 2. 547.3
	1. 7923 2. 537.3			1. 7928 2. 547.3
	1. 7924 2. 547.3			1. 7929 2. 547.3
	1. 7925 2. 547.3			1. 7930 2. 549.3

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 7931 2. 555.31			1. 7936 2. 567.31
	1. 7932 2. 555.31			1. 7937 2. 567.31
	1. 7933 2. 557.31			1. 7938 2. 569.31
	1. 7934 2. 557.31			1. 7939 2. 571.31
	1. 7935 2. 557.31			1. 7940 2. 571.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7941 2. 608.33		1. 7946 2. 471.26
	1. 7942 2. 621.34		1. 7947 2. 473.26
	1. 7943 2. 443.24		1. 7948 2. 483.27
	1. 7944 2. 457.25		1. 7949 2. 485.27
	1. 7945 2. 471.26		1. 7950 2. 489.27



















































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 7951 2. 499.27		1. 7956 2. 536.29
	1. 7952 2. 513.28		1. 7957 2. 539.3
	1. 7953 2. 514.28		1. 7958 2. 541.3
	1. 7954 2. 525.29		1. 7959 2. 547.3
	1. 7955 2. 525.29		1. 7960 2. 572.31

TABLE 80











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8001 2. 459.25		1. 8006 2. 477.26
	1. 8002 2. 459.25		1. 8007 2. 485.27
	1. 8003 2. 469.26		1. 8008 2. 485.27
	1. 8004 2. 469.26		1. 8009 2. 485.27
	1. 8005 2. 473.26		1. 8010 2. 486.27

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8011 2. 493.27		1. 8016 2. 495.27
	1. 8012 2. 495.27		1. 8017 2. 495.27
	1. 8013 2. 495.27		1. 8018 2. 502.28
	1. 8014 2. 495.27		1. 8019 2. 502.28
	1. 8015 2. 495.27		1. 8020 2. 502.28

Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 8021 2. 505.28			1. 8026 2. 509.28
	1. 8022 2. 505.28			1. 8027 2. 509.28
	1. 8023 2. 507.28			1. 8028 2. 513.28
	1. 8024 2. 507.28			1. 8029 2. 513.28
	1. 8025 2. 507.28			1. 8030 2. 513.28





Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8031 2. 515.28		1. 8036 2. 523.29
	1. 8032 2. 519.29		1. 8037 2. 525.29
	1. 8033 2. 521.29		1. 8038 2. 529.29
	1. 8034 2. 523.29		1. 8039 2. 529.29
	1. 8035 2. 523.29		1. 8040 2. 530.29

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8041 2. 530.29		1. 8046 2. 547.3
	1. 8042 2. 535.29		1. 8047 2. 544.3
	1. 8043 2. 537.3		1. 8048 2. 547.3
	1. 8044 2. 547.3		1. 8049 2. 547.3
	1. 8045 2. 547.3		1. 8050 2. 547.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8051 2. 548.3		1. 8056 2. 569.31
	1. 8052 2. 548.3		1. 8057 2. 569.31
	1. 8053 2. 547.3		1. 8058 2. 571.31
	1. 8054 2. 555.31		1. 8059 2. 608.33
	1. 8055 2. 553.3		1. 8060 2. 621.34
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8061 2. 443.24		1. 8066 2. 473.26
	1. 8062 2. 457.25		1. 8067 2. 483.27
	1. 8063 2. 471.26		1. 8068 2. 485.27
	1. 8064 2. 471.26		1. 8069 2. 483.27
	1. 8065 2. 469.26		1. 8070 2. 499.27

534

Product	1. Ex. 2. m/z
	1. 8071 2. 505.2 8
	1. 8072 2. 537.3
	1. 8073 2. 537.3
	1. 8074 2. 547.3

535

TABLE 81

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8101 2. 510.28		1. 8106 2. 528.29		1. 8111 2. 542.3
	1. 8102 2. 518.28		1. 8107 2. 537.3		1. 8112 2. 542.3
	1. 8103 2. 518.28		1. 8108 2. 538.3		1. 8113 2. 548.3
	1. 8104 2. 519.29		1. 8109 2. 540.3		1. 8114 2. 552.3
	1. 8105 2. 526.29		1. 8110 2. 540.3		1. 8115 2. 554.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8116 2. 556.31		1. 8121 2. 580.32		1. 8126 2. 588.32
	1. 8117 2. 558.31		1. 8122 2. 580.32		1. 8127 2. 590.32
	1. 8118 2. 562.31		1. 8123 2. 580.32		1. 8128 2. 590.32
	1. 8119 2. 580.32		1. 8125 2. 580.32		1. 8129 2. 604.33
	1. 8120 2. 580.32		1. 8126 2. 588.32		1. 8130 2. 612.34

537













































Product	1. Ex. 2. m/z
	1. 8131 2. 654.36
	1. 8132 2. 476.26
	1. 8133 2. 580.32
	1. 8134 2. 580.32











TABLE 82











Product	1. Ex. 2. m/z		Product	1. Ex. 2. m/z
	1. 8201 2. 488.27			1. 8206 2. 512.28
	1. 8202 2. 496.27			1. 8207 2. 514.28
	1. 8203 2. 504.28			1. 8208 2. 523.29
	1. 8204 2. 504.28			1. 8209 2. 524.29
	1. 8205 2. 505.28			1. 8210 2. 526.29











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8211 2. 526.29		1. 8216 2. 534.29
	1. 8212 2. 528.29		1. 8217 2. 537.3
	1. 8213 2. 528.29		1. 8218 2. 538.3
	1. 8214 2. 528.29		1. 8219 2. 542.3
	1. 8215 2. 532.29		1. 8220 2. 542.3

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8221 2. 544.3		1. 8226 2. 551.3
	1. 8222 2. 544.3		1. 8227 2. 554.3
	1. 8223 2. 548.3		1. 8228 2. 556.31
	1. 8224 2. 548.3		1. 8229 2. 566.31
	1. 8225 2. 551.3		1. 8230 2. 566.31

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8231 2. 566.31		1. 8236 2. 568.31
	1. 8232 2. 566.31		1. 8237 2. 574.32
	1. 8233 2. 566.31		1. 8238 2. 574.32
	1. 8243 2. 566.31		1. 8239 2. 576.32
	1. 8235 2. 566.31		1. 8240 2. 576.32
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8241 2. 576.32		1. 8246 2. 590.32
	1. 8242 2. 586.32		1. 8247 2. 598.33
	1. 8243 2. 586.32		1. 8248 2. 627.34
	1. 8244 2. 588.32		1. 8249 2. 640.35
	1. 8245 2. 590.32		1. 8250 2. 462.25

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8251 2. 476.26		1. 8256 2. 504.28
	1. 8252 2. 490.27		1. 8257 2. 504.28
	1. 8253 2. 492.27		1. 8258 2. 508.28
	1. 8254 2. 492.27		1. 8259 2. 530.29
	1. 8255 2. 502.28		1. 8260 2. 533.29
















Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8261 2. 544.3		1. 8266 2. 560.31
	1. 8262 2. 546.3		1. 8267 2. 566.31
	1. 8263 2. 546.3		1. 8268 2. 591.33
	1. 8264 2. 558.31		1. 8269 2. 595.33
	1. 8265 2. 558.31		1. 8270 2. 555.31
















545
















TABLE 83

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8301 2. 465.882		1. 8306 2. 491.79		1. 8311 2. 501.825
	1. 8302 2. 465.885		1. 8307 2. 491.792		1. 8312 2. 501.833
	1. 8303 2. 475.827		1. 8308 2. 492.787		1. 8313 2. 510.822
	1. 8304 2. 479.89		1. 8309 2. 499.846		1. 8314 2. 510.821
	1. 8305 2. 483.829		1. 8310 2. 501.826		1. 8315 2. 511.841






546

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8316 2. 513.855		1. 8321 2. 519.783		1. 8326 2. 531.804
	1. 8317 2. 515.832		1. 8322 2. 519.781		1. 8327 2. 531.812
	1. 8318 2. 515.832		1. 8323 2. 525.813		1. 8328 2. 535.83
	1. 8319 2. 515.837		1. 8324 2. 529.806		1. 8329 2. 535.831
	1. 8320 2. 519.782		1. 8325 2. 529.81		1. 8330 2. 541.788

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8331 2. 543.821		1. 8336 2. 554.727		1. 8341 2. 561.838
	1. 8332 2. 553.797		1. 8337 2. 554.738		1. 8342 2. 564.714
	1. 8333 2. 553.796		1. 8338 2. 554.738		1. 8343 2. 564.704
	1. 8334 2. 553.795		1. 8339 2. 554.727		1. 8344 2. 564.72
	1. 8335 2. 554.73		1. 8340 2. 561.836		1. 8345 2. 573.823











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8346 2. 573.735		1. 8351 2. 529.808		1. 8356 2. 479.856
	1. 8347 2. 575.839		1. 8352 2. 614.763		1. 8357 2. 511.842
	1. 8348 2. 577.818		1. 8353 2. 628.727		1. 8358 2. 520.859
	1. 8349 2. 577.814		1. 8354 2. 449.857		1. 8359 2. 533.903
	1. 9350 2. 585.818		1. 8355 2. 463.867		1. 8360 2. 545.823











549

Product	1. Ex. 2. m/z
	1. 8361 2. 547.909
	1. 8362 2. 510.817
	1. 8363 2. 542.833
	1. 8364 2. 573.815
	1. 8365 2. 547.798











550

TABLE 84











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8401 2. 491.883		1. 8406 2. 509.829
	1. 8402 2. 491.883		1. 8407 2. 517.795
	1. 8403 2. 501.823		1. 8408 2. 517.799
	1. 8404 2. 501.827		1. 8409 2. 518.796
	1. 8405 2. 505.89		1. 8410 2. 525.851











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8411 2. 527.831		1. 8416 2. 539.859
	1. 8412 2. 527.833		1. 8417 2. 541.84
	1. 8413 2. 536.833		1. 8418 2. 541.843
	1. 8414 2. 536.829		1. 8419 2. 541.843
	1. 8415 2. 537.845		1. 8420 2. 546.786











552

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8421 2. 546.792		1. 8426 2. 555.817
	1. 8422 2. 546.79		1. 8427 2. 557.81
	1. 8423 2. 547.854		1. 8428 2. 557.813
	1. 8424 2. 551.826		1. 8429 2. 561.839
	1. 8425 2. 555.819		1. 8430 2. 561.833

553

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8431 2. 564.841		1. 8436 2. 579.794
	1. 8432 2. 567.795		1. 8437 2. 580.738
	1. 8433 2. 569.823		1. 8438 2. 580.732
	1. 8434 2. 579.797		1. 8439 2. 580.733
	1. 8435 2. 579.796		1. 8440 2. 580.734

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8441 2. 580.726		1. 8446 2. 590.726
	1. 8442 2. 581.805		1. 8447 2. 590.724
	1. 8443 2. 587.844		1. 8448 2. 599.834
	1. 8444 2. 587.835		1. 8449 2. 599.749
	1. 8445 2. 590.723		1. 8450 2. 601.854

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8451 2. 603.831		1. 8456 2. 640.779
	1. 8452 2. 603.83		1. 8457 2. 654.746
	1. 8453 2. 611.83		1. 8458 2. 475.871
	1. 8454 2. 555.824		1. 8459 2. 489.879
	1. 8455 2. 585.817		1. 8460 2. 503.885








































Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8461 2. 505.866		1. 8466 2. 559.919
	1. 8462 2. 507.828		1. 8467 2. 573.925
	1. 8463 2. 537.859		1. 8468 2. 579.876
	1. 8464 2. 546.877		1. 8469 2. 536.831
	1. 8465 2. 559.921		1. 8470 2. 568.843









TABLE 85

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8501 2. 367.2		1. 8506 2. 443.1
	1. 8502 2. 381.2		1. 8507 2. 323.1
	1. 8503 2. 443.2		1. 8508 2. 365.2
	1. 8504 2. 457.3		1. 8509 2. 379.21
			1. 8510 2. 381.21











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8511 2. 381.21		1. 8516 2. 409.22
	1. 8512 2. 393.22		1. 8517 2. 411.23
	1. 8513 2. 395.22		1. 8518 2. 421.23
	1. 8514 2. 397.22		1. 8519 2. 421.23
	1. 8515 2. 407.22		1. 8520 2. 425.23

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8521 2. 429.24		1. 8526 2. 445.24
	1. 8522 2. 439.24		1. 8527 2 .444.24
	1. 8523 2. 443.24		1. 8528 2. 444.24
	1. 8524 2. 443.24		1. 8529 2. 446.25
	1. 8525 2. 445.24		1. 8530 2. 457.25











560

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8531 2. 480.26		1. 8536 2. 431.24
	1. 8532 2. 483.27		1. 8537 2. 429.24
	1. 8533 2. 483.27		1. 8538 2. 397.22
			1. 8539 2. 411.23
			1. 8540 2. 411.23











561











Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8541 2. 432.24		1. 8547 2. 478.26
	1. 8542 2. 433.24		1. 8548 2. 498.27
	1. 8543 2. 438.24		1. 8549 2. 365.1
	1. 8545 2. 464.26		1. 8549 2. 337.1
	1. 8546 2. 466.26		1. 8550 2. 351.1











563

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8611 2. 417.23		1. 8616 2. 434.24
	1. 8612 2. 415.23		1. 8617 2. 446.25
	1. 8613 2. 428.24		1. 8618 2. 460.25
	1. 8614 2. 430.24		1. 8619 2. 459.25
	1. 8615 2. 434.24		1. 8620 2. 474.26











564

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8621 2. 377.21		1. 8626 2. 427.23
	1. 8622 2. 406.22		1. 8627 2. 431.24
	1. 8623 2. 412.23		1. 8628 2. 446.25
	1. 8624 2. 420.23		1. 8629 2. 446.25
	1. 8625 2. 426.23		1. 8630 2. 407.2



Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
 <p>Chiral</p>	1. 8631 2. 407.2		1. 8636 2. 475.3
	1. 8632 2. 409.2		1. 8637 2. 419.2
 <p>Chiral</p>	1. 8633 2. 421.2		1. 8638 2. 448.2
	1. 8634 2. 439.2	 <p>Chiral</p>	1. 8639 2. 379.2
	1. 8635 2. 455.3		1. 8640 2. 437.2

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8641 2. 415.23		1. 8646 2. 423.23
	1. 8642 2. 443.24		1. 8647 2. 430.24
	1. 8643 2. 416.23		1. 8648 2. 437.24
	1. 8644 2. 417.23		1. 8649 2. 439.24
	1. 8645 2. 423.23		1. 8650 2. 466.26

567

Product	1. Ex. 2. m/z	Product	1. Ex. 2. m/z
	1. 8651 2. 423.23		1. 8656 2. 445.24
	1. 8652 2. 425.23		1. 8657 2. 445.24
	1. 8653 2. 437.24		1. 8658 2. 451.25
	1. 8654 2. 437.24		1. 8659 2. 476.26
	1. 8655 2. 437.24		1. 8660 2. 479.26


568

Product	1. Ex. 2. m/z		
	1. 8661 2. 477.26		
	1. 8662 2. 479.26		

CLAIMS

What is claimed is:


1. A compound represented by the structural formula:



5

or a pharmaceutically acceptable salt or solvate of said compound,
wherein:



R is H, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, cycloalkyl,
cycloalkylalkyl, alkenylalkyl, alkynylalkyl, heterocyclyl, heterocyclylalkyl,
10 heteroarylalkyl (including N-oxide of said heteroaryl), $-(\text{CHR}^5)_n$ -aryl, $-(\text{CHR}^5)_n$ -



wherein each of said alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocyclyl, and
15 heteroaryl can be unsubstituted or optionally substituted with one or more
moieties which can be the same or different, each moiety being independently
selected from the group consisting of halogen, alkyl, aryl, cycloalkyl,
heterocyclylalkyl, CF_3 , OCF_3 , CN, $-\text{OR}^5$, $-\text{NR}^5\text{R}^{10}$, $-\text{C}(\text{R}^4\text{R}^5)_p\text{-R}^9$,
 $-\text{N}(\text{R}^5)\text{Boc}$, $-(\text{CR}^4\text{R}^5)_p\text{OR}^5$, $-\text{C}(\text{O}_2)\text{R}^5$, $-\text{C}(\text{O})\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{SO}_3\text{H}$, $-\text{SR}^{10}$,
20 $-\text{S}(\text{O}_2)\text{R}^7$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$, $-\text{N}(\text{R}^5)\text{S}(\text{O}_2)\text{R}^7$, $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^7$ and $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{NR}^5\text{R}^{10}$;


R^2 is selected from the group consisting of R^9 , alkyl, alkenyl, alkynyl, CF_3 ,
heterocyclyl, heterocyclylalkyl, halogen, haloalkyl, aryl, arylalkyl, heteroarylalkyl,
alkynylalkyl, cycloalkyl, heteroaryl, alkyl substituted with 1-6 R^9 groups which can
be the same or different and are independently selected from the list of R^9
25 shown below, aryl substituted with 1-3 aryl or heteroaryl groups which can be the

- same or different and are independently selected from phenyl, pyridyl, thiophenyl, furanyl and thiazolo groups, aryl fused with an aryl or heteroaryl group, heteroaryl substituted with 1-3 aryl or heteroaryl groups which can be the same or different and are independently selected from phenyl, pyridyl,
- 5 thiophenyl, furanyl and thiazolo groups, heteroaryl fused with an aryl or



- wherein one or more of the aryl and/or one or more of the heteroaryl in
- 10 the above-noted definitions for R² can be unsubstituted or optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, -CN, -OR⁵, -SR⁵, -S(O₂)R⁶, -S(O₂)NR⁵R⁶, -NR⁵R⁶, -C(O)NR⁵R⁶, CF₃, alkyl, aryl and OCF₃;

- R³ is selected from the group consisting of H, halogen, -NR⁵R⁶, -OR⁶,
- 15 -SR⁶, -C(O)N(R⁵R⁶), alkyl, alkynyl, cycloalkyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl and heteroarylalkyl,



- wherein each of said alkyl, cycloalkyl, aryl, arylalkyl, heterocyclyl,
- 20 heterocyclylalkyl, heteroaryl and heteroarylalkyl for R³ and the heterocyclyl moieties whose structures are shown immediately above for R³ can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, CF₃, CN, -OCF₃,

$-(CR^4R^5)_pOR^5$, $-OR^5$, $-NR^5R^6$, $-(CR^4R^5)_pNR^5R^6$, $-C(O_2)R^5$, $-C(O)R^5$, $-C(O)NR^5R^6$, $-SR^6$, $-S(O_2)R^6$, $-S(O_2)NR^5R^6$, $-N(R^5)S(O_2)R^7$, $-N(R^5)C(O)R^7$ and $-N(R^5)C(O)NR^5R^6$, with the proviso that no carbon adjacent to a nitrogen atom on a heterocyclyl ring carries a $-OR^5$ moiety;

5 R^4 is H, halo or alkyl;

R^5 is H, alkyl, aryl or cycloalkyl;

R^6 is selected from the group consisting of H, alkyl, alkenyl, aryl, arylalkyl, arylalkenyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl, wherein each of said alkyl, aryl, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl can be unsubstituted or optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, heterocyclylalkyl, CF_3 , OCF_3 , CN, $-OR^5$, $-NR^5R^{10}$, $-C(R^4R^5)_pR^9$, $-N(R^5)Boc$, $-(CR^4R^5)_pOR^5$, $-C(O_2)R^5$, $-C(O)R^5$, $-C(O)NR^5R^{10}$, $-SO_3H$, $-SR^{10}$, $-S(O_2)R^7$, $-S(O_2)NR^5R^{10}$, $-N(R^5)S(O_2)R^7$, $-N(R^5)C(O)R^7$ and $-N(R^5)C(O)NR^5R^{10}$;

R^{10} is selected from the group consisting of H, alkyl, aryl, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl, wherein each of said alkyl, aryl, arylalkyl, cycloalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, and heteroarylalkyl can be unsubstituted or optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, heterocyclylalkyl, CF_3 , OCF_3 , CN, $-OR^5$, $-NR^4R^5$, $-C(R^4R^5)_pR^9$, $-N(R^5)Boc$, $-(CR^4R^5)_pOR^5$, $-C(O_2)R^5$, $-C(O)NR^4R^5$, $-C(O)R^5$, $-SO_3H$, $-SR^5$, $-S(O_2)R^7$, $-S(O_2)NR^4R^5$, $-N(R^5)S(O_2)R^7$, $-N(R^5)C(O)R^7$ and $-N(R^5)C(O)NR^4R^5$;

or optionally (i) R^5 and R^{10} in the moiety $-NR^5R^{10}$, or (ii) R^5 and R^6 in the moiety $-NR^5R^6$, may be joined together to form a cycloalkyl or heterocyclyl moiety, with each of said cycloalkyl or heterocyclyl moiety being unsubstituted or optionally independently being substituted with one or more R^9 groups;

R^7 is selected from the group consisting of alkyl, cycloalkyl, aryl, arylalkenyl, heteroaryl, arylalkyl, heteroarylalkyl, heteroarylalkenyl, and

heterocyclyl, wherein each of said alkyl, cycloalkyl, heteroarylalkyl, aryl, heteroaryl and arylalkyl can be unsubstituted or optionally independently substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, CF_3 , OCF_3 , CN , $-\text{OR}^5$, $-\text{NR}^5\text{R}^{10}$, $-\text{CH}_2\text{OR}^5$, $-\text{C}(\text{O}_2)\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{C}(\text{O})\text{R}^5$, $-\text{SR}^{10}$, $-\text{S}(\text{O}_2)\text{R}^{10}$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$, $-\text{N}(\text{R}^5)\text{S}(\text{O}_2)\text{R}^{10}$, $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^{10}$ and $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{NR}^5\text{R}^{10}$;

R^8 is selected from the group consisting of R^6 , $-\text{OR}^6$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(=\text{N}-\text{CN})-\text{NH}_2$, $-\text{C}(=\text{NH})-\text{NHR}^5$, heterocyclyl, and $-\text{S}(\text{O}_2)\text{R}^7$;

R^9 is selected from the group consisting of halogen, $-\text{CN}$, $-\text{NR}^5\text{R}^{10}$, $-\text{C}(\text{O}_2)\text{R}^6$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{10}$, $-\text{OR}^6$, $-\text{SR}^6$, $-\text{S}(\text{O}_2)\text{R}^7$, $-\text{S}(\text{O}_2)\text{NR}^5\text{R}^{10}$, $-\text{N}(\text{R}^5)\text{S}(\text{O}_2)\text{R}^7$, $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^7$ and $-\text{N}(\text{R}^5)\text{C}(\text{O})\text{NR}^5\text{R}^{10}$;

m is 0 to 4;

n is 1 to 4; and

p is 1 to 4,

with the proviso that when R^2 is phenyl, R^3 is not alkyl, alkynyl or halogen, and


that when R^2 is aryl, R is not $\begin{array}{c} \text{~} \\ \text{~} \\ \text{~} \end{array} \text{---}(\text{CHR}^5)_n\text{---NR}^5\text{R}^8$, and with the further proviso that when R is arylalkyl, then any heteroaryl substituent on the aryl of said arylalkyl contains at least three heteroatoms.

2. The compound of claim 1, wherein R is $-(\text{CHR}^5)_n\text{-aryl}$, $-(\text{CHR}^5)_n\text{-heteroaryl}$, alkyl, cycloalkyl, heterocyclyl, or heteroarylalkyl (including N-oxide of said heteroaryl), wherein each of said alkyl, aryl, cycloalkyl, heterocyclyl and heteroaryl can be unsubstituted or optionally substituted with one or more moieties as stated in claim 1;

R^2 is halogen, alkyl, haloalkyl, CN , cycloalkyl, heterocyclyl or alkynyl;

R^3 is H, lower alkyl, aryl, heteroaryl, cycloalkyl, $-\text{NR}^5\text{R}^6$,

573



- wherein said alkyl, aryl, heteroaryl, cycloalkyl and the heterocyclyl structures shown immediately above for R^3 are optionally substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of halogen, CF_3 , OCF_3 , lower alkyl, CN, $-C(O)R^5$, $-S(O_2)R^5$, $-C(=NH)-NH_2$, $-C(=CN)-NH_2$, hydroxyalkyl, alkoxy carbonyl, $-SR^5$, and OR^5 , with the proviso that no carbon adjacent to a nitrogen atom on a heterocyclyl ring carries a $-OR^5$ moiety;

R^4 is H or lower alkyl;


R^5 is H, lower alkyl or cycloalkyl;

n is 1 to 2; and

p is 1 or 2.

3. The compound of claim 2, wherein R is hydroxyalkyl, $-(CHR^5)_n$ -aryl, or $-(CHR^5)_n$ -heteroaryl, wherein each of said aryl and heteroaryl is unsubstituted or substituted with one or more groups which can be the same or different, each group being independently selected from the group consisting of heteroaryl, amine, heterocyclyl, $-C(O)N(R^5R^6)$, $-S(O_2)R^5$, $-S(O_2)N(R^5R^6)$, alkoxy and halo.
4. The compound of claim 2, wherein R^2 is Br, Cl, CF_3 , CN, lower alkyl, cyclopropyl, alkynyl, alkyl substituted with $-OR^6$ or tetrahydrofuranyl.
5. The compound of claim 2, wherein R^3 is H, lower alkyl, aryl, heteroaryl, cycloalkyl,

574




wherein each of said alkyl, aryl, heteroaryl, cycloalkyl and the heterocyclyl


- 5 structures shown immediately above for R^3 are optionally substituted with one or more moieties which moieties can be the same or different, each moiety being independently selected from the group consisting of halogen, CF_3 , OCF_3 , lower alkyl, CN and OR^5 , with the proviso that no carbon adjacent to a nitrogen atom on a heterocyclyl ring carries a $-OR^5$ moiety.
- 10 6. The compound of claim 2, wherein R^4 is H or lower-alkyl.
7. The compound of claim 2, wherein R^5 is H.
8. The compound of claim 2, wherein n is 1.
9. The compound of claim 1, wherein p is 1.
10. The compound of claim 2, wherein R is benzyl or hydroxyalkyl.
- 15 11. The compound of claim 2, wherein R is pyrid-3-ylmethyl, wherein said pyridyl may be unsubstituted or optionally independently substituted with one or more moieties as stated in claim 1.
12. The compound of claim 2, wherein R is pyrid-4-ylmethyl, wherein said pyridyl may be unsubstituted or optionally independently substituted with one or
- 20 more moieties as stated in claim 1.
13. The compound 2, wherein R is the N-oxide of pyrid-2-ylmethyl, pyrid-3-ylmethyl, or pyrid-4-ylmethyl, wherein each of said pyridyl may be unsubstituted or optionally independently substituted with one or more moieties as stated in claim 1.

14. The compound of claim 4, wherein said R^2 is Br.
15. The compound of claim 4, wherein said R^2 is Cl.
16. The compound of claim 4, wherein R^2 is ethyl.
17. The compound of claim 4, wherein R^2 is cyclopropyl.
- 5 18. The compound of claim 4, wherein R^2 is ethynyl.
19. The compound of claim 2, wherein R^3 is lower alkyl, cycloalkyl, heterocyclyl, aryl or $-N(R^5R^6)$.
20. The compound of claim 19, wherein R^3 is isopropyl.
21. The compound of claim 19, wherein R^3 is cyclohexyl or norbornyl wherein
- 10 each of said cyclohexyl or norbornyl can be unsubstituted or substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of alkyl and hydroxyalkyl.
22. The compound of claim 19, wherein R^3 is unsubstituted phenyl.
23. The compound of claim 19, wherein R^3 is a phenyl substituted with one or
- 15 moieties which can be the same or different, each moiety being independently selected from the group consisting of F, Br, Cl and CF_3 .
24. The compound of claim 19, wherein R^5 of said $-N(R^5R^6)$ is H or hydroxyalkyl, and R^6 of said $-N(R^5R^6)$ is selected from the group consisting of
- 20 alkyl, hydroxyalkyl, cycloalkyl and methylenedioxy, wherein each of said alkyl and cycloalkyl can be unsubstituted or substituted with one or more moieties which can be the same or different, each moiety being independently selected from the group consisting of amine, ethoxycarbonyl, amide, hydroxyalkyl, hydroxy,
- 25 The compound of claim 19, wherein R^5 and R^6 of said $-N(R^5R^6)$ are joined together to form a heterocyclyl moiety, wherein said heterocyclyl moiety can be
- 25 unsubstituted or optionally independently substituted with one or more groups which can be the same or different, each group being selected from the group consisting of hydroxyalkyl, amide, $-C(O)R^5$, $>C(CH_3)_2$, $-S(O_2)R^5$, $-S(O_2)N(R^5R^6)$, $-C(=NH)N(R^5R^6)$ and $-C(=N-CN)N(R^5R^6)$.
26. The compound of claim 25, wherein said heterocyclyl moiety formed by R^5
- 30 and R^6 is a pyrrolidine or piperidine ring.
27. A compound of the formula:


577




578




579




581







584




585




586



587




588




5

589




590



or a pharmaceutically acceptable salt or solvate thereof.


5 28. A compound of the formula:




The image displays 15 chemical structures of pyrazolo[1,5-a]pyrimidine derivatives, arranged in a grid. Each structure features a pyrazolo[1,5-a]pyrimidine core with a bromine atom at the 3-position and an amino group at the 6-position. The structures are differentiated by their substituents at the 2-position and the 4-position of the pyrimidine ring.

- Structure 1 (top left):** 2-(furan-2-yl)-6-((pyridin-2-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 2 (top left, second):** 2-(furan-2-yl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 3 (top middle):** 2-(cyclohexyl)-6-((1H-indol-3-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 4 (top right):** 2-(2-fluorophenyl)-6-((1H-indol-3-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 5 (second row, left):** 2-phenyl-6-((1H-indol-3-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 6 (second row, middle):** 2-phenyl-6-((4-nitrophenylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 7 (second row, right):** 2-phenyl-6-((4-nitrophenylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 8 (third row, left):** 2-(tetrahydro-2H-pyran-4-yl)-6-((pyridin-2-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 9 (third row, middle):** 2-(tetrahydro-2H-pyran-4-yl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 10 (third row, right):** 2-(2-fluorophenyl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 11 (third row, far right):** 2-(cyclohexyl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 12 (bottom row, left):** 2-(thiophen-2-yl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 13 (bottom row, second):** 2-(cyclohexyl)-6-((pyridin-2-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 14 (bottom row, third):** 2-(cyclohexyl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.
- Structure 15 (bottom row, right):** 2-(cyclohexyl)-6-((pyridin-4-ylmethyl)amino)-3-bromopyrazolo[1,5-a]pyrimidine.


592



593




594




or a pharmaceutically acceptable salt or solvate thereof.

5 29. A compound of the formula:


595




596




597






599



or a pharmaceutically acceptable salt or solvate thereof.

5 30. A compound of the formula:

600



5 or a pharmaceutically acceptable salt or solvate thereof.

31. A method of inhibiting one or more cyclin dependent kinases, comprising administering a therapeutically effective amount of at least one compound of claim 1 to a patient in need of such inhibition.

32. A method of treating one or more diseases associated with cyclin
 10 dependent kinase, comprising administering a therapeutically effective amount of at least one compound of claim 1 to a patient in need of such treatment.

33. The method of claim 32, wherein said cyclin dependent kinase is CDK2.

34. The method of claim 32, wherein said cyclin dependent kinase is mitogen activated protein kinase (MAPK/ERK).

15 35. The method of claim 32, wherein said cyclin dependent kinase is glycogen synthase kinase 3 (GSK3beta).

36. The method of claim 32, wherein said disease is selected from the group consisting of:

5 cancer of the bladder, breast, colon, kidney, liver, lung, small cell lung cancer, esophagus, gall bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma;

leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma and Burkett's lymphoma;

10 acute and chronic myelogenous leukemia, myelodysplastic syndrome and promyelocytic leukemia;

fibrosarcoma, rhabdomyosarcoma;

astrocytoma, neuroblastoma, glioma and schwannomas;

melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma

15 pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma.

37. A method of treating one or more diseases associated with cyclin dependent kinase, comprising administering to a mammal in need of such treatment

20 an amount of a first compound, which is a compound of claim 1, or a pharmaceutically acceptable salt or solvate thereof; and

an amount of at least one second compound, said second compound being an anti-cancer agent;

25 wherein the amounts of the first compound and said second compound result in a therapeutic effect.

38. The method of claim 37, further comprising radiation therapy.

39. The method of claim 37, wherein said anti-cancer agent is selected from the group consisting of a cytostatic agent, cisplatin, doxorubicin, taxotere, taxol, etoposide, CPT-11, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, 30 epothilones, tamoxifen, 5-fluorouracil, methoxtrexate, 5FU, temozolomide, cyclophosphamide, SCH 66336, R115777, L778,123, BMS 214662, Iressa, Tarceva, antibodies to EGFR, Gleevec, intron, ara-C, adriamycin, cytoxan,

- gemcitabine, Uracil mustard, Chlormethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine,
- 5 Fludarabine phosphate, oxaliplatin, leucovirin, ELOXATIN™, Pentostatine, Vinblastine, Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, Mithramycin, Deoxycoformycin, Mitomycin-C, L-Asparaginase, Teniposide 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate,
- 10 Testolactone, Megestrolacetate, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, goserelin, Cisplatin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, CPT-
- 15 11, Anastrozole, Letrozole, Capecitabine, Reloxafine, Droloxafine, or Hexamethylmelamine..
40. A pharmaceutical composition comprising a therapeutically effective amount of at least one compound of claim 1 in combination with at least one pharmaceutically acceptable carrier.
- 20 41. The pharmaceutical composition of claim 38, additionally comprising one or more anti-cancer agents selected from the group consisting of cytostatic agent, cisplatin, doxorubicin, taxotere, taxol, etoposide, CPT-11, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, epothilones, tamoxifen, 5-fluorouracil, methotrexate, 5FU, temozolomide, cyclophosphamide, SCH 66336,
- 25 R115777, L778,123, BMS 214662, Iressa, Tarceva, antibodies to EGFR, Gleevec, intron, ara-C, adriamycin, cytoxan, gemcitabine, Uracil mustard, Chlormethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine,
- 30 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, Pentostatine, Vinblastine, Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, Mithramycin, Deoxycoformycin, Mitomycin-C,

- L-Asparaginase, Teniposide 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrolacetate, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone,
- 5 Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, goserelin, Cisplatin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, CPT-11, Anastrozole, Letrozole, Capecitabine, Reloxafine, Droloxafine, or Hexamethylmelamine.
- 10 42. A compound of claim 1 in purified form.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 03/27555

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/519 //(C07D487/04,239:00,231:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 628 559 A (BEIERSDORF LILLY GMBH) 14 December 1994 (1994-12-14) cited in the application page 3, formula (I) page 4, partial structure (A) page 7, partial structure (L) page 8, line 46 - line 48 ---	27-30
A	US 6 107 305 A (BURSUKER ISIA ET AL) 22 August 2000 (2000-08-22) cited in the application column 1, formula (I) column 1, line 24 - line 25 column 11, line 19 --- -/--	27-30

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

17 December 2003

Date of mailing of the international search report

29/12/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoepfner, W

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 03/27555

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 6 413 974 B1 (BORCHERDING DAVID R ET AL) 2 July 2002 (2002-07-02) cited in the application abstract column 15, formula (I) claim 26 ---	27-30
A	WO 02 50079 A (ULIBARRI GERARD ;LONCHAMPT MARIE-ODILE (FR); THURIEAU CHRISTOPHE ()) 27 June 2002 (2002-06-27) cited in the application page 1, line 1 - line 3 page 5, formula (I) page 8, line 21 -----	27-30

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-26,31-42

Present claims 1-26 and 31-42 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a relatively limited proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

In particular:

the vast number of values for most of the variables, in conjunction with their cascading meanings and especially the presence of open definitions such as "alkyl", "arylalkyl" or "heterocyclyl" and the like render the scope of the invention, for which protection is sought, ill-defined and obscure. Consequently, an exhaustive and complete search is precluded for practical and economical reasons. The search was based upon though not limited to examples and tables given in the description and to the subject-matter of claims 27-30 (cf. Arts. 6, 15 and Rule 33 PCT).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 03/27555

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 31-39 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1-26, 31-42
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 03/27555

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0628559	A	14-12-1994	EP 0628559 A1	14-12-1994
			CA 2125458 A1	11-12-1994
			DE 69430279 D1	08-05-2002
			DE 69430279 T2	06-03-2003
			ES 2174863 T3	16-11-2002
			JP 7002860 A	06-01-1995
			US 5571813 A	05-11-1996
			US 5602136 A	11-02-1997
			US 5602137 A	11-02-1997
<hr/>				
US 6107305	A	22-08-2000	AU 747705 B2	23-05-2002
			AU 1629799 A	05-07-1999
			CA 2314355 A1	24-06-1999
			EP 1043998 A1	18-10-2000
			JP 2002508324 T	19-03-2002
			WO 9930710 A1	24-06-1999
			ZA 9811178 A	07-06-2000
<hr/>				
US 6413974	B1	02-07-2002	NONE	
<hr/>				
WO 0250079	A	27-06-2002	FR 2818278 A1	21-06-2002
			AU 2509602 A	01-07-2002
			CA 2432417 A1	27-06-2002
			EP 1345941 A1	24-09-2003
			WO 0250079 A1	27-06-2002
<hr/>				

THIS PAGE BLANK (USPTO)